The Journal of Laryngology and Otology

EDITED BY
G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

Voice after Laryngectomy	SHEILA DRUMMOND
THE INCIDENCE OF LARYNGEAL CANCER IN SINGAPORE	C. S. Muir
OBSERVATIONS ON "SO-CALLED" LATENT LABYRINTH- ITIS AND THE FISTULA SYMPTOM	A. C. ARULPRAGASAM
SURGICAL TREATMENT OF LEUKOPLAKIA OF THE	J. F. PILBEAM and B. Cohen
EXTENDED LARYNGOPHARYNGECTOMY FOR CARCI- NOMA OF THE LARYNGOPHARYNX AND THE CERVICAL OESOPHAGUS (USE OF A FREE GRAFT OF LARGE INTESTINE)	P. CHRYSOSPATHIS and A. COYAS
EXPERIENCE IN THE USE OF THE NON-DELAYED CHEST FLAP IN THE RECONSTRUCTION OF THE PHARYNX AND CERVICAL OESOPHAGUS FOLLOWING RADICAL SURGERY	SIEGFRIED ZEHM
A STATISTICAL STUDY ON THE NASAL SEPTUM IN / 5,096 CASES IN IRAQ	DAOUD S. ALI
CLINICAL RECORDS— MALIGNANT MELANOMA OF THE NASAL SEPTUM (A CASE REPORT WITH BRIEF REVIEW OF	
LITERATURE)	M. KANNAN KUTTY and T. SREEDHARAN
OAT-CELL CARCINOMA AND PARANASAL SINUSES	R. N. RAYCHOWDHURI
LINGUAL ABSCESS	EUGENE N. MYERS and JOHN B. ROBERTS
Converse to Minimum	

London

Headley Brothers

109 Kingsway WC2

Annual Subscription £5/5/0 net, U.S.A. \$15

Monthly, 12/6 net post free

The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

EDITED BY

H. BATEMAN

ASSISTANT EDITOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of

this most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of fro per article; beyond this authors are expected to pay half the cost. Coloured illustrations will

be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2.

5. Orders for reprints must be sent when returning galley proofs, and for this

purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the Journal of Laryngology. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to The Editor, Journal of

Laryngology, c/o Headley Brothers, 109 Kingsway, London, WC2.

8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.

9. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, Headley Brothers, 109 Kingsway, London, WC2.

United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, Headley Brothers, 109 Kingsway, London, WC2, England.

Tournal of Laryngology and Otology, 1965

CONTENTS

	PAGE
THE EFFECTS OF ENVIRONMENTAL NOISE ON PSEUDO VOICE AFTER LARYNGECTOMY. Sheila Drummond (Melbourne)	193
THE INCIDENCE OF LARYNGEAL CANCER IN SINGAPORE. C. S. Muir (Singapore)	203
Observations on "So-Called" Latent Labyrinthitis and the Fistula Symptom. A. C. Arulpragasam (Jaffna, Ceylon)	214
SURGICAL TREATMENT OF LEUKOPLAKIA OF THE PALATE. J. F. Pilbeam and B. Cohen (Barnet)	225
EXTENDED LARYNGOPHARYNGECTOMY FOR CARCINOMA OF THE LARYNGOPHARYNX AND THE CERVICAL ŒSOPHAGUS (USE OF A FREE GRAFT OF LARGE INTESTINE). P. Chrysospathis and A. Coyas (Athens, Greece)	233
EXPERIENCE IN THE USE OF THE NON-DELAYED CHEST FLAP IN THE RECONSTRUCTION OF THE PHARYNX AND CERVICAL ŒSOPHAGUS FOLLOWING RADICAL SURGERY. Siegfried Zehm (Würzburg, Germany)	237
A STATISTICAL STUDY ON THE NASAL SEPTUM IN 5,096 CASES IN IRAQ. Daoud S. Ali (Baghdad)	244
CLINICAL RECORDS— MALIGNANT MELANOMA OF THE NASAL SEPTUM (A CASE REPORT WITH BRIEF REVIEW OF LITERATURE). M. Kannan Kutty and T. Sreedharan (Calicut, South India)	249
OAT-CELL CARCINOMA AND PARANASAL SINUSES. R. N. Raychowdhuri (Barnsley)	253
LINGUAL ABSCESS. Eugene N. Myers (Boston, Massachusetts) and John B. Roberts (Albuquerque, New Mexico)	256
GENERAL NOTES	259

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$16.00 per year Canada \$16.75 per year Foreign \$17.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D. EDITOR

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.

When the patient needs a hearing aid—whom do you recommend?

INGRAMS, of course, the firm with the different makes to choose from.

INGRAMS, 2 Shepherd Street, London, W.1

FOR SURGERY

NEW SYSTEMIC HAEMOSTATIC HAEMOSTATIC ENSURES A CLEAR FIELD



Clinical studies of DICYNENE have shown its efficacy in the prevention and treatment of haemorrhage due to the rupture of small vessels. DICYNENE exhibits low toxicity. The drug is well tolerated and there are no known contra-indications. For further details write to BAXTER LABORATORIES LTD., High Wycombe, Bucks. Tel. High Wycombe 4612





SEE THE FULL RANGE OF ZEISS EQUIPMENT ON DAILY DEMONSTRATION AT STAND No. 4, LABEX, EARLS COURT, MARCH 29th—APRIL 2nd



Orbenin is recommended for the immediate treatment of all Hospital Staphylococcal infections

Orbenin, an oral penicillin stable to penicillinase, is bactericidal to resistant and sensitive staphylococci



Orbenin (regd.) (cloxacillin sodium) is a product of British research at

Beecham Research Laboratories

"The discovery of a...harmless means of effectively reducing the viscosity of sputum...would be a considerable advance in the management of patients with a variety of chest diseases..."


(Editorial, Brit. med. J., i:930, 1962)



the first mucodissolvent

cutorsadas anomanas and dataraants

superseues enzymes una aetergenis
produces true chemical lysis on contact
liquefies both mucoid and purulent secretions





THE BRITISH DRUG HOUSES LTD. LONDON N.I.

"N-acetyl-L-cysteine has proved to be the most effective agent to date for the liquefaction of mucoid and purulent secretions, apparently regardless of their character or origin."

(J. thorac. cardiov. Surg., 44:830, 1962)

UNIQUE MODE OF ACTION

The active constituent of AIRBRON is acetylcysteine, a derivative of the naturally occurring amino acid, L-cysteine. It acts by opening up the disulphide bonds present in mucus, dramatically reducing the viscosity.

COMPARISON WITH EARLIER AGENTS

In vitro measurement of sputum viscosity, comparing AIRBRON with an enzyme, saline solution and detergent, have been made. These show that AIRBRON produces both a more pronounced and a more prolonged reduction of sputum viscosity than these other agents.²

USE IN TRACHEOSTOMY CARE

According to one investigator ³, the advantages of using acetylcysteine in tracheostomy care are that:

- * Plugging of tracheostomies by secretions is avoided
- * Traumatic bronchitis due to very frequent and vigorous suction is avoided
- * The need for bronchoscopies to remove secretions is obviated
- * Secondary pulmonary infections are avoided

In addition to tracheostomy care, AIRBRON may well have many other uses in laryngology and otology.

AIRBRON may be given by nebulization or by direct instillation into the tracheo-bronchial tree. It is supplied in ampoules of 10 and 30 ml., each in boxes of 3.

REFERENCES

- 1. Ann. N.Y. Acad. Sci., 106:298, 1963
- Paper read at the meeting of the American Medical Association, Chicago, June 1962
- J. thorac. cardiov. Surg., 44:330, 1962
 Amer. Rev. resp. Dis., 90:111, 1964
- 5. Dis. Chest, 46:66, 1964

- 6. J. Pediat., 62:31, 1963
- 7. Sth. med. J., 56:1271, 1963
- 8. Amer. Rev. resp. Dis., 87:775, 1963

9. Amer. Rev. resp. Dis., 89:284, 1964

Full literature and file card available on request

403 B 3345



when the patient must remain alert

Fabahistin®

Mebhydrolin B.P.C.

the non-sedating antihistamine

Children: 1-4 tablets daily · Adults: 2-6 tablets daily



® Registered trade mark of BAYER GERMANY
U.K. Subsidiary FBA PHARMACEUTICALS LTD, Haywards Heath, Sussex

ONLY AMPLIVOX provides a complete audiometric service!



Transistor Audiometers

for the school screening service, ward use, etc.

Speech Audiometry Equipment

via records and tape, including new attachment for use with tape recorders.






Clinical Audiometers

with narrow band masking for the highest accuracy and extended resolution.

Audiometer Booths

for ideal testing conditions in Hospitals, Clinics, Factories. Standard and special sizes available.




PLUS

An efficient and speedy calibration service

AMPLIVOX LTD Medical Acoustic Division

Please write for technical ANII AI V A literature covering your needs 80 New Bond Street London, WI Tel



OTOSPORIN'

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' Drops. 'Otosporin' is an aqueous suspension containing neomycin and polymyxin B which are together effective against practically all the bacteria found in ear infections. In addition, it contains hydrocortisone to reduce inflammation and swelling, thereby providing easier access for the antibiotics.

'OTOSPORIN' BRAND Drops ISSUED IN BOTTLES OF 5 ml.



BURROUGHS WELLCOME & CO. (THE WELLCOME FOUNDATION LTD.) LONDON