

## Molecular hydrogen generated by elemental magnesium supplementation alters rumen fermentation and microbiota in goats

Min Wang<sup>1,2</sup>, Rong Wang<sup>1,3</sup>, XiuMin Zhang<sup>1</sup>, Emilio M. Ungerfeld<sup>4</sup>, Donglei Long<sup>1,3</sup>, HongXiang Mao<sup>1,3</sup>, JinZhen Jiao<sup>1</sup>, Karen A. Beauchemin<sup>5</sup> and Zhiliang Tan<sup>1,2\*</sup>

<sup>1</sup>Key Laboratory for Agro-Ecological Processes in Subtropical Region, National Engineering Laboratory for Pollution Control and Waste Utilization in Livestock and Poultry Production, Institute of Subtropical Agriculture, The Chinese Academy of Sciences, Changsha, Hunan 410125, People's Republic of China

<sup>2</sup>Hunan Co-Innovation Center of Animal Production Safety (CICAPS), Changsha, Hunan 410128, People's Republic of China

<sup>3</sup>College of Animal Science and Technology, Hunan Agricultural University, Changsha 410128, People's Republic of China

<sup>4</sup>Instituto de Investigaciones Agropecuarias (INIA) Carillanca, Temuco 8340422, Chile

<sup>5</sup>Lethbridge Research and Development Centre, Agriculture and Agri-Food Canada, Lethbridge, Alberta T1J 4B1, Canada

(Submitted 8 May 2017 – Final revision received 29 June 2017 – Accepted 24 July 2017 – First published online 20 September 2017)

### Abstract

We tested the hypotheses that supplementation of a diet with elemental Mg increases ruminal dissolved H<sub>2</sub> (dH<sub>2</sub>) in rumen fluid, which in turn alters rumen fermentation and microbial community in goats. In a randomised block design, twenty growing goats were allocated to two treatments fed the same basal diet with 1.45% Mg(OH)<sub>2</sub> or 0.6% elemental Mg. After 28 d of adaptation, we collected total faeces to measure total tract digestibility, rumen contents to analyse fermentation end products and microbial groups, and measured methane (CH<sub>4</sub>) emission using respiration chambers. Ruminal Mg<sup>2+</sup> concentration was similar in both treatments. Elemental Mg supplementation increased dH<sub>2</sub> at 2.5 h post morning feeding (+180%, *P* < 0.001). Elemental Mg supplementation decreased total volatile fatty acid concentration (−8.6%, *P* < 0.001), the acetate:propionate ratio (−11.8%, *P* < 0.03) and fungal copy numbers (−63.6%, *P* = 0.006), and increased propionate molar percentage (+11.6%, *P* < 0.001), methanogen copy numbers (+47.9%, *P* < 0.001), dissolved CH<sub>4</sub> (+35.6%, *P* < 0.001) and CH<sub>4</sub> emissions (+11.7%, *P* = 0.03), compared with Mg(OH)<sub>2</sub> supplementation. The bacterial community composition in both treatments was overall similar. Ruminal dH<sub>2</sub> was negatively correlated with acetate molar percentage and fungal copy numbers (*P* < 0.05), and positively correlated with propionate molar percentage and methanogen copy numbers (*P* < 0.05). In summary, elemental Mg supplementation increased ruminal dH<sub>2</sub> concentration, which inhibited rumen fermentation, enhanced methanogenesis and seemed to shift fermentation pathways from acetate to propionate, and altered microbiota by decreasing fungi and increasing methanogens.

**Key words:** Elemental magnesium: Fermentation pathways: Dissolved hydrogen: Methane: Rumen microbiota

Volatile fatty acids (VFA), carbon dioxide (CO<sub>2</sub>) and molecular H<sub>2</sub> are produced during carbohydrate fermentation by bacteria, protozoa and fungi in the rumen. VFA serve as a major energy source for the host animal<sup>(1)</sup>, with acetate and propionate being the main precursors of fat and glucose, respectively. The main consumers of H<sub>2</sub> are methanogenic Archaea, producing methane (CH<sub>4</sub>) as an end product of fermentation, which keeps a low H<sub>2</sub> partial pressure in the rumen<sup>(2)</sup>. It is understood that the accumulation of H<sub>2</sub>, which typically occurs when methanogenesis is inhibited, could hinder the re-oxidation of reduced electron carriers and adversely affect fermentation and fibre digestion<sup>(3)</sup>.

Besides being used as a substrate for methanogenesis, H<sub>2</sub> is also involved in VFA production<sup>(2,4)</sup>. This is because different

numbers of moles of H<sub>2</sub> (or of reducing equivalent pairs in reduced co-factors), are released or incorporated in the production pathway of each VFA. For example, fermentation of 1 mol of glucose to acetate releases 4 mol of electron pairs, most of which are transferred to protons to form H<sub>2</sub>. In contrast, fermentation of glucose to propionate involves H<sub>2</sub> or metabolic H incorporation<sup>(2)</sup>. When animals are switched from fibrous to starchy diets, or rumen methanogenesis is inhibited, dissolved H<sub>2</sub> (dH<sub>2</sub>) in the rumen elevates, and, as expected from the stoichiometry of acetate and propionate production, fermentation shifts from acetate to propionate<sup>(5–7)</sup>.

However, when the effect of direct addition of H<sub>2</sub> as gas (gaseous H<sub>2</sub>, gH<sub>2</sub>) has been studied in *in vitro* batch cultures<sup>(8–10)</sup> and *in vivo*<sup>(11)</sup>, the results observed in acetate and

**Abbreviations:** CH<sub>4</sub>, methane; CO<sub>2</sub>, carbon dioxide; dCH<sub>4</sub>, dissolved CH<sub>4</sub>; dCO<sub>2</sub>, dissolved carbon dioxide; dH<sub>2</sub>, dissolved H<sub>2</sub>; gCO<sub>2</sub>, gaseous CO<sub>2</sub>; gH<sub>2</sub>, gaseous H<sub>2</sub>; H<sub>2</sub>, molecular hydrogen; OUT, operational taxonomic units; VFA, volatile fatty acid.

\* **Corresponding author:** Z. Tan, fax +86 731 4612685, email zltan@isa.ac.cn

propionate do not fully agree with that rationale. In batch cultures, Broudiscou *et al.*<sup>(12)</sup> found that the effect of an initial 0.5 or 1 kPa gH<sub>2</sub> headspace on acetate and propionate production depended on the inoculum used, and when using an inoculum adapted to a fibrous substrate, gH<sub>2</sub> addition unexpectedly increased acetate and decreased propionate. Similarly, Patra & Yu<sup>(8)</sup> found that including gH<sub>2</sub> in the initial culture headspace unexpectedly resulted in lower propionate molar percentage. Similarly, Qiao *et al.*<sup>(10)</sup> found that increasing gH<sub>2</sub> in the headspace of batch cultures actually increased acetate molar percentage and the acetate:propionate ratio, and decreased propionate molar percentage. Infusion of gH<sub>2</sub> in the rumen of dairy cows did not result in changes in total VFA concentration or VFA profile<sup>(11)</sup>. The effect of gH<sub>2</sub> on dH<sub>2</sub> concentration was not reported in those studies. Given that dH<sub>2</sub> and gH<sub>2</sub> have been shown not to be at equilibrium *in vitro*<sup>(13)</sup> and *in vivo*<sup>(14)</sup>, and that the form of H<sub>2</sub> available to microbes is dH<sub>2</sub> rather than gH<sub>2</sub>, we propose that the role of H<sub>2</sub> on fermentation needs further examination by studying the effect of dH<sub>2</sub> addition.

Our first hypothesis for this study using goats as the experimental ruminant model was that dH<sub>2</sub> could be increased through supplementing elemental Mg to the diet, as elemental Mg would react with water in rumen fluid releasing H<sub>2</sub>: Mg + 2 H<sub>2</sub>O → Mg(OH)<sub>2</sub> + H<sub>2</sub><sup>(15)</sup>. Our second hypothesis was that increased dH<sub>2</sub> would alter rumen fermentation and the microbial community composition.

## Methods

All animal procedures were approved by the Animal Care Committee, Institute of Subtropical Agriculture, the Chinese Academy of Sciences (CAS), Changsha, China.

### Goats and diets

A randomised block design with two treatments was used to investigate the effects of H<sub>2</sub> generated by the reaction of elemental Mg with water in the rumen fluid of goats. In all, twenty growing Xiangdong black male goats (mean initial body weight = 20.3 (SD 3.42) kg) were allocated to ten blocks according to body weight and CH<sub>4</sub> emission (g/kg DM intake) measured in a preliminary experiment (unpublished results). Two treatments were formulated with equal dietary Mg content: basal diet plus 1.45% (DM basis) Mg(OH)<sub>2</sub> powder (99% purity; Beijing Taizejiaye Technology Development Co., Ltd) for the control group and basal diet with 0.60% (DM basis) elemental Mg powder (99% purity; Beijing Taizejiaye Technology Development Co., Ltd) for the hypothesised elevated dH<sub>2</sub> group. In a preliminary experiment (unpublished results), these treatments had been shown to be harmless to the health of goats. Each block of animals contained two goats and each goat within a block was randomly assigned to one of the two dietary treatments.

Goats were kept in individual pens and had free access to fresh water. The diet was formulated to meet 1.4 times maintenance requirements of goats. The composition of the basal diet is shown in Table 1. Dietary forage and concentrate were

**Table 1.** Ingredients and chemical composition of the basal diet offered to goats (g/kg DM)

Items	Diet
<b>Dietary ingredient (g/kg DM)</b>	
<b>Forage</b>	
Rice straw	300
<b>Concentrate</b>	
Soyabeans	54
Maize	304
Wheat bran	290
CaCO <sub>3</sub>	5
Fat	11
NaCl	5
Urea	11
Premix*	20
<b>Chemical composition (g/kg DM)</b>	
DM	960
Organic matter	912
Ash	88.3
Crude protein	150
Starch	262
Neutral detergent fibre	312
Acid detergent fibre	186
Diethyl ether extract	35.0
Gross energy (MJ/kg)	18.2

\* Premix was formulated to provide the following (per kg of premix): 400 g of NaHCO<sub>3</sub>, 2 g of Fe, 1 g of Cu, 0.01 g of Co, 0.05 g of I, 6.6 g of Mn, 4.4 g of Zn, 0.003 g of Se, 333 mg of retinol, 5 mg of cholecalciferol, 838 mg of α-tocopherol.

not mixed, and both forage and concentrate were offered separately, each divided into two equal portions at 08.00 and 17.00 hours. Diets were offered for a 28-d adaptation period before conducting measurements. The elemental Mg and Mg(OH)<sub>2</sub> supplements were mixed with the concentrate fraction immediately before feeding to avoid the reaction with water in the environment, and the concentrate fraction containing the elemental Mg and Mg(OH)<sub>2</sub> supplements was eaten completely within approximately 1 h. During the initial 10 d of adaptation to diets, feed was offered *ad libitum* targeting 5% refusals. The amount of feed allocated daily during the next 18 d of adaptation was adjusted to 100% of the DM intake previously measured in order to minimise feed selection. The refusals, when present, were collected and analysed to determine the actual nutrient intakes.

### Nutrient digestibility

Nutrient digestibility was determined over a 5-d period from days 29 to 33. Total faeces and urine were collected twice daily, weighed and mixed daily. A subsample (approximately 1%) was frozen immediately at -20°C, and another subsample (approximately 1%) was acidified using 10% (w/w) H<sub>2</sub>SO<sub>4</sub> to prevent N loss and then frozen immediately at -20°C. The faeces samples were later dried at 60°C for 48 h in a forced-air oven, and ground through a 1-mm screen. The acidified oven-dried samples were used for total N analysis, whereas non-acidified, oven-dried samples were used for other chemical analyses.

### Rumen sampling

Collection of rumen contents was performed at 0, +2.5 and +6 h relative to the commencement of the morning feeding on days

34 and 35. Rumen contents (300 ml) were collected by oral stomach tubing as described by Wang *et al.*<sup>(16)</sup>, with the initial 100 ml discarded to avoid saliva contamination. The pH of rumen contents was measured immediately after sampling using a portable pH meter (Starter 300; Ohaus Instruments Co. Ltd). Two subsamples of 35 ml each were immediately transferred into 50-ml plastic syringes for measuring dH<sub>2</sub> and dissolved CH<sub>4</sub> (dCH<sub>4</sub>) concentration, as described by Wang *et al.*<sup>(13)</sup>. Two other 35-ml subsamples were immediately frozen at -80°C in liquid N<sub>2</sub> for DNA extraction and subsequent microbial analyses. In addition, 2-ml samples of rumen contents were collected and centrifuged at 15 000 *g* for 10 min at 4°C, and 1.5 ml of supernatant was acidified using 0.15 ml of 25 % (w/v) metaphosphoric acid, and stored at -20°C for subsequent measurement of VFA concentration. The remaining sample of rumen contents was stored at -20°C for the measurements of ammonium (NH<sub>4</sub><sup>+</sup>), Mg<sup>2+</sup> and glucose concentration.

### Methane and carbon dioxide emissions

CH<sub>4</sub> and CO<sub>2</sub> emissions were measured in three plexiglass respiration chambers that permitted the goats to see each other, thereby minimising stress. CH<sub>4</sub> and CO<sub>2</sub> emissions were measured individually for each goat for 48 h using the protocol of Wang *et al.*<sup>(17)</sup> slightly modified. In brief, during seven periods of 2 d each, each block of animals containing two goats assigned to different treatments was randomly assigned to a chamber. One goat from each block was then randomly assigned to a measurement period, and the second goat from that block was placed in the same chamber in the subsequent period. Within the chamber, the goats were restrained with free access to a feed bin and drinking water. Airflow was maintained under negative pressure (flow rate = 35 m<sup>3</sup>/h). Outlet gas from the chamber and ambient gas were connected to a multiport inlet unit of a gas analyser (GGA-30p; Los Gatos Research) for measuring CH<sub>4</sub> and CO<sub>2</sub> concentration. The cycling time to measure CH<sub>4</sub> and CO<sub>2</sub> concentration produced in the chamber was 30 min, with 8 min for analysing gases from each of the three chambers and 6 min for analysing background environmental concentrations of CH<sub>4</sub> and CO<sub>2</sub> in incoming air. Daily CH<sub>4</sub> and CO<sub>2</sub> emissions were calculated using net CH<sub>4</sub> and CO<sub>2</sub> concentrations and flow rate of air through each chamber at 30-min intervals, and differences between chambers were corrected using the methodology described by McGinn *et al.*<sup>(18)</sup>. The chambers were opened twice a day at 08.00 and 18.00 hours to deliver the diets. The chamber cleaning, such as swapping faeces and urine trays, took place during the morning before feeding.

### Sample analysis

All samples of feeds, refusals and faeces were dried and ground to pass through a 1-mm sieve. Contents of DM (method 945.15), organic matter (OM) (method 942.05), crude protein (method 954.01) and diethyl ether extract (method 920.39) were determined according to published methodologies<sup>(19)</sup>. Gross energy was determined using an isothermal automatic calorimeter (5E-AC8018; Changsha Kaiyuan Instruments Co. Ltd). Contents of neutral detergent fibre (NDF) and acid detergent fibre were

determined and expressed inclusive of residual ash<sup>(20)</sup>, and NDF was assayed with the addition of a heat-stable amylase, but without sodium sulphite. The starch content was determined after pre-extraction with ethanol (80%), and glucose released from starch by enzyme hydrolysis was measured using amyloglucosidase<sup>(21)</sup>.

Frozen acidified rumen samples were thawed and centrifuged at 15 000 *g* for 10 min at 4°C, and individual VFA concentrations in the supernatant were measured using GC (Agilent 7890A; Agilent Inc.), according to the method described by Wang *et al.*<sup>(13)</sup>. The estimated net H<sub>2</sub> production relative to the amount of total VFA produced (R<sub>NH2</sub>) was estimated according to the stoichiometric equation developed by Wang *et al.*<sup>(16)</sup>, under the assumption of equal fractional rates of individual VFA absorption. Ammonia and glucose in the supernatant were determined colorimetrically according to the methods of Weatherburn<sup>(22)</sup> and Nelson<sup>(23)</sup>, respectively. The concentration of Mg<sup>2+</sup> in the supernatant was determined by Inductively Coupled Plasma-Optical Emission Spectrometers using Varian 720-ES series (Agilent Inc.).

Dissolved gases were also measured immediately after sampling using the procedure described by Wang *et al.*<sup>(16)</sup> with a slight modification. In brief, a 20-ml syringe containing 10 ml of N<sub>2</sub> gas was connected to a 50-ml plastic syringe containing 35-ml rumen content samples via polyurethane tubing. The N<sub>2</sub> gas was then injected into the 50-ml syringe, and the gases dissolved in the rumen fluid were extracted into the N<sub>2</sub> gas phase by vigorously hand shaking for 5 min. Gaseous H<sub>2</sub> and CH<sub>4</sub> (gCH<sub>4</sub>) concentrations in the gas phase were measured by GC (Agilent 7890A). The dH<sub>2</sub> and dCH<sub>4</sub> concentrations in the original rumen fluid were calculated using equations described by Wang *et al.*<sup>(13)</sup>. Gaseous CO<sub>2</sub> (gCO<sub>2</sub>) was calculated as the total dissolved gas extracted minus dCH<sub>4</sub> extracted, assuming that rumen gases would be composed of CO<sub>2</sub> and CH<sub>4</sub>. Total dissolved CO<sub>2</sub> (dCO<sub>2</sub>) concentration in the original rumen fluid (C<sub>TdCO<sub>2</sub></sub>, mol/l) was calculated by combining equations from Wang *et al.*<sup>(16)</sup> and Hille *et al.*<sup>(24)</sup>, and expressed as follows:

$$\begin{aligned}
 C_{TdCO_2} &= C_{eTdCO_2} + V_g C_{gCO_2} / (22 \cdot 4 V_l) \\
 C_{eTdCO_2} &= \alpha_{CO_2} C_{gCO_2} 10^{(pH - pK_{CO_2})} + \alpha_{CO_2} C_{gCO_2} \\
 C_{gCO_2} &= (V_g - V_N - V_g C_{gCH_4}) / V_g \\
 \alpha_{CO_2} &= \frac{100}{22.4 \exp(-6.8346 + 1.2817(10^4/T) - 3.7668(10^6/T^2) + 2.997(10^8/T^3))}
 \end{aligned}$$

where C<sub>eTdCO<sub>2</sub></sub> is the total dCO<sub>2</sub> concentration (mol/l) in the rumen fluid at equilibrium after extraction; C<sub>gCO<sub>2</sub></sub> the gCO<sub>2</sub> concentration (litres/l) measured in the gas phase of the 20-ml syringe at equilibrium after extraction of dissolved gases; C<sub>gCH<sub>4</sub></sub> the gCH<sub>4</sub> concentration (litres/l) measured in the gas phase of the 20-ml syringe at equilibrium after extraction of dissolved gases; V<sub>g</sub> the gas volume (ml) at equilibrium after extraction of dissolved gases; V<sub>l</sub> the volume of liquid (ml); V<sub>N</sub> the injected N<sub>2</sub> gas volume (10 ml); pK<sub>CO<sub>2</sub></sub> the dissociation constant of bicarbonate and set to be 6.11<sup>(24)</sup>; α<sub>CO<sub>2</sub></sub> the Bunsen absorption coefficient (mol/l·atm) for CO<sub>2</sub><sup>(25)</sup>; and T the temperature in K (273 + temperature in °C).

### Estimation of Gibbs free-energy changes

Gibbs energy changes ( $\Delta G$ ) of fermentation reactions (online Supplementary Table S1) were estimated using measured concentrations of metabolites at 0, +2.5 and +6 h relative to the commencement of the morning feeding. Gibbs energy changes of reactions at standard conditions of 298 K were calculated from standard Gibbs energy of formation of reactants and products, and then adjusted to a rumen temperature of 312 K (39°C) using the van't Hoff equation<sup>(26)</sup>. Gibbs energy changes estimated for actual rumen conditions were subsequently adjusted using measured concentrations of soluble metabolites ( $10^{-\text{pH}}$ , glucose, acetate, propionate and butyrate) and dissolved gases ( $\text{dH}_2$ ,  $\text{dCO}_2$  and  $\text{dCH}_4$ )<sup>(14)</sup>.

### Quantitative real-time PCR analyses

Rumen samples (2 ml) at each of the three sampling time points were pooled and freeze-dried, and then physically disrupted using a bead beater for 1 min. Genomic DNA was extracted using the QIAamp DNA Stool Mini kit (Qiagen) according to the manufacturer's instructions. The quantity of DNA was measured based on absorbance at 260 and 280 nm using a NanoDrop ND 100 (Nano Drop Technologies). The absolute quantification of total bacteria, protozoa, fungi, methanogens and select bacterial species was measured by quantitative real-time PCR (qPCR) using primers validated in our laboratory (online Supplementary Table S2)<sup>(27)</sup>. Quantitative PCR was performed according to the procedures described by Jiao *et al.*<sup>(27)</sup>. Final absolute amounts of target groups or species were estimated by relating the  $C_T$  value to the standard curves and expressed as  $\log_{10}$  copies/DM rumen contents. The abundances of six select rumen bacterial species (*Ruminococcus albus*, *R. flavefaciens*, *Fibrobacter succinogenes*, *Selenomonas ruminantium*, *Ruminobacter amylophilus* and *Prevotella ruminicola*) and of genus *Prevotella* spp. were measured using qPCR and species-specific 16S rRNA gene-targeted primers (online Supplementary Table S2) and expressed relative to the total bacterial DNA. The abundances of each bacterial species and of genus *Prevotella* spp. were determined using the  $\Delta C_T$  method<sup>(28)</sup>.

### High-throughput sequencing and analysis

Extracted purified DNA (50 ng) from each rumen sample was subjected to PCR amplification of the V3–V4 region of 16S rRNA gene using universal bacterial primers 338 F (5'-ACTCCTACG GGAGGCAGCA-3') and 806 R (5'-GGA CTACHVGGGTWT CTAAT-3')<sup>(29)</sup>. PCR was performed using a GeneAmp<sup>®</sup> 9700 thermal cycler (Applied Biosystems). The PCR products were purified using the AxyPrep<sup>™</sup> DNA Gel Extraction Kit (Axygen Biosciences) according to the manufacturer's instructions and quantified using the QuantiFluor<sup>™</sup>-ST system (Promega). Purified PCR products were high-throughput-sequenced using an Illumina MiSeq PE300 instrument at Majorbio Bio-Pharm Technology Co., Ltd, using protocols recommended by procedures of Miseq reagent kits v3 (Illumina). Sequences were quality filtered and demultiplexed using exact matches to the supplied DNA barcodes.

Bacterial phylotypes were identified using *uclust*<sup>(30)</sup> and assigned to operational taxonomic units (OTU) at 97% sequence identity. Taxonomic identity of each phylotype was determined using the Greengenes database<sup>(31)</sup>. The resulting OTU were combined into an OTU table that represented abundance of each OTU in each microbial sample. Alpha diversity of bacterial communities was obtained using Mothur version 1.30.1. Similarity between bacterial communities was assessed using the Bray–Curtis distance metric and visualised using principal coordinates (PCoA) analysis.

### Statistical analyses

The statistical model used included dietary treatment and block as fixed effects and sampling day as random effect. When sampling time was included, the model included dietary treatment and block as fixed effects, sampling time as a repeated-measure variable, and the interaction between dietary treatment and sampling time and sampling day as random effects. The best linear or log linear regression was derived between  $\text{dH}_2$  and concentration of other rumen metabolites using ordinary least squares. Associations between response variables were studied through calculating their Pearson's correlation coefficient ( $r$ ) and statistical significance.  $P \leq 0.05$  was considered significant, and  $0.05 < P \leq 0.1$  was accepted as a tendency to significance. The SPSS 12.0 software was used for the statistical analyses.

### Results

Although no differences were observed for feed intake and digestibility between both treatments, elemental Mg supplementation increased  $\text{CH}_4$  emissions expressed as g/d (+11.7%,  $P = 0.03$ ) and as g/kg OM intake (+9.87%,  $P = 0.03$ ) (Table 2). No differences were observed in ruminal  $\text{Mg}^{2+}$  concentration and pH between the two treatments (Table 3). The pH was negatively correlated ( $r = -0.75$ ,  $P < 0.001$ ) with ruminal  $\text{Mg}^{2+}$  concentration (online Supplementary Fig. S1).

Elemental Mg supplementation increased the average  $\text{dH}_2$  (+95.1%,  $P = 0.02$ ) and  $\text{dCH}_4$  concentrations (+39.0%,  $P < 0.001$ ) (Table 3). Interactions between treatment and time were observed for  $\text{dH}_2$  ( $P = 0.02$ ) and  $\text{dCH}_4$  ( $P < 0.001$ ). Elemental Mg supplementation increased  $\text{dH}_2$  only at +2.5 h (+180%,  $P < 0.001$ ) but not at 0 or +6 h relative to the commencement of the morning feeding, and increased  $\text{dCH}_4$  at +2.5 h (+63.4%,  $P < 0.001$ ) and +6 h (+50.5%,  $P < 0.001$ ) but not at 0 h relative to the commencement of the morning feeding (Fig. 1). At +2.5 h, ruminal  $\text{dH}_2$  was positively correlated with  $\text{dCH}_4$  ( $r = 0.50$ ,  $P = 0.02$ ) (online Supplementary Fig. S2). Furthermore, ruminal  $\text{dH}_2$  concentration varied widely among the 10 goats treated with elemental Mg supplementation, the highest  $\text{dH}_2$  concentration being 17 mmol/l. When this highest  $\text{dH}_2$  point was removed, the Pearson's correlation coefficient between  $\text{dH}_2$  and  $\text{dCH}_4$  was 0.486 ( $P = 0.03$ , data not shown).

Elemental Mg supplementation decreased total VFA molar concentration (−11.6%,  $P < 0.001$ ), acetate molar percentage (−3.80%,  $P < 0.001$ ), the acetate to propionate ratio (−11.8%,  $P < 0.03$ ) and  $R_{\text{NH}_2}$  (−42.1%,  $P < 0.03$ ), and increased propionate molar percentage (+11.6%,  $P < 0.001$ ) (Table 3). Elemental

Mg supplementation slightly increased  $\Delta G$  of glucose fermentation to acetate ( $P=0.03$ ) and to butyrate ( $P=0.04$ ), did not alter  $\Delta G$  of glucose to 2/3 acetate + 4/3 propionate ( $P=0.36$ ) and tended ( $P=0.009$ ) to decrease  $\Delta G$  of glucose + 2H<sub>2</sub> to propionate at +2.5 h relative to the commencement of the morning feeding (Table 4).

**Table 2.** Feed intake and digestibility, methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) emissions in growing goats fed diets with 1.45% Mg(OH)<sub>2</sub> or 0.6% elemental magnesium after 28 d of adaption ( $n$  10) (Mean values with their standard errors)

Items	Mg(OH) <sub>2</sub>		Elemental Mg		P
	Mean	SE	Mean	SE	
<b>Intake (g/d)</b>					
DM	577	17.4	573	17.4	0.46
OM	524	16.2	522	15.9	0.73
CP	86.4	2.60	87.3	2.49	0.25
NDF	251	7.91	247	8.21	0.21
ADF	106	3.5	103	3.9	0.10
Starch	175	5.2	178	5.0	0.11
GE	7.67	0.229	7.81	0.219	0.10
<b>Total tract apparent digestibility (g/kg)</b>					
OM	683	15.3	689	9.2	0.78
CP	779	16.1	797	9.7	0.47
NDF	572	26.7	591	16.9	0.57
Starch	980	5.8	981	5.3	0.65
<b>CH<sub>4</sub> emissions</b>					
g/d	8.50	0.481	9.50	0.584	0.03
g/kg OM intake	16.2	0.85	17.8	0.71	0.03
g/kg OM digested	23.9	1.42	25.9	0.83	0.10
<b>CO<sub>2</sub> emissions</b>					
g/d	363	15.0	388	18.6	0.25

CP, crude protein; NDF, neutral detergent fibre; ADF, acid detergent fibre; GE, gross energy.

Because the hypothesised effect on dH<sub>2</sub> was observed only at +2.5 h relative to the commencement of the morning feeding, we focused on this time point for studying the effects of elemental Mg supplementation on the associations between concentration of dH<sub>2</sub> and other rumen metabolites. Ruminal dH<sub>2</sub> was negatively correlated with total VFA concentration ( $r$  -0.60,  $P=0.005$ ), acetate molar percentage ( $r$  -0.76,  $P<0.001$ ), acetate:propionate ratio ( $r$  -0.60,  $P=0.005$ ) and R<sub>NH2</sub> ( $r$  -0.40,  $P=0.08$ ), and positively correlated with propionate molar percentage ( $r$  0.47,  $P=0.03$ ) at +2.5 h relative to the commencement of the morning feeding (online Supplementary Fig. S3). When the greatest dH<sub>2</sub> data point was removed, only a trend ( $r$  -0.39,  $P=0.09$ , data not shown) to a correlation between dH<sub>2</sub> and acetate molar percentage at +2.5 h was observed.


Elemental Mg supplementation decreased the copy numbers of fungi (-63.6%,  $P=0.006$ ) and the relative abundance of *S. ruminantium* (-39.7%,  $P=0.008$ ), and increased the copy numbers of methanogens (+47.9%,  $P<0.001$ ) and relative abundance of *R. flavefaciens* (+71%,  $P=0.02$ ) (online Supplementary Table S3). Ruminal +2.5 h dH<sub>2</sub> was negatively correlated with copy numbers of fungi ( $r$  -0.53,  $P=0.02$ ) and the relative abundance of *S. Ruminantium* ( $r$  -0.43,  $P=0.05$ ), and positively correlated with copy numbers of methanogens ( $r$  0.54,  $P=0.01$ ) (Fig. 2). When the greatest dH<sub>2</sub> data point was removed, a significant correlation ( $r$  -0.54,  $P=0.01$ , data not shown) between +2.5 h dH<sub>2</sub> and copy numbers of fungi was still observed.

Pyrosequencing of the 16S rRNA gene resulted in 38730 (SD 3761) sequences per sample (range of 30 525–44 554 sequences per sample). Supplementation with elemental Mg increased community Ace ( $P=0.04$ ) and bias-corrected Chao 1 ( $P=0.04$ ) richness, but did not alter bacterial diversity of

**Table 3.** Concentration of rumen metabolites in growing goats fed diets with 1.45% Mg(OH)<sub>2</sub> or 0.6% elemental magnesium after 28 d of adaption ( $n$  10) (Mean values with their standard errors)

Items	Mg(OH) <sub>2</sub>		Elemental Mg		P		
	Mean	SE	Mean	SE	Treatment	Time	Treatment × Time
Mg <sup>2+</sup> (mmol/l)	5.71	0.332	5.29	0.314	0.16	<0.001	0.39
pH	6.66	0.027	6.64	0.034	0.31	<0.001	0.003
Glucose (mmol/l)	4.64	0.389	5.28	0.352	0.18	0.254	0.92
<b>Dissolved gases</b>							
dH <sub>2</sub> (μmol/l)	1.02	0.121	1.99	0.415	0.02	<0.001	0.02
dCH <sub>4</sub> (mmol/l)	1.05	0.045	1.46	0.058	<0.001	<0.001	<0.001
dCO <sub>2</sub> (mmol/l)	57.4	5.82	58.3	4.75	0.82	0.002	0.48
<b>Rumen fermentation</b>							
Ammonia (mmol/l)	7.54	0.829	8.23	0.994	0.17	0.02	0.13
Total VFA (mmol/l)	98.5	2.59	90.0	2.29	<0.001	<0.001	0.85
Acetate:Propionate	3.38	0.060	2.98	0.087	<0.001	0.003	0.97
<b>Individual VFA (mol/100 mol total VFA)</b>							
Acetate	63.1	0.62	60.7	0.77	<0.001	0.19	0.99
Propionate	18.9	0.25	21.1	0.43	<0.001	<0.001	0.81
Butyrate	11.9	0.61	11.8	0.62	0.91	0.69	0.99
Valerate	1.84	0.088	1.92	0.116	0.57	<0.001	0.65
/iso-butyrate	1.45	0.067	1.56	0.078	0.10	<0.001	0.29
/iso-valerate	2.88	0.119	3.00	0.157	0.39	<0.001	0.68
<b>Estimated net H<sub>2</sub> production relative to the amount of total VFA produced (mol/mol)</b>							
R <sub>NH2</sub>	1.29	0.008	1.22	0.012	<0.001	<0.001	0.88

dH<sub>2</sub>, dissolved H<sub>2</sub>; dCH<sub>4</sub>, dissolved methane; dCO<sub>2</sub>, dissolved carbon dioxide; VFA, volatile fatty acids.



**Fig. 1.** Dissolved hydrogen (dH<sub>2</sub>, a) and methane (dCH<sub>4</sub>, b) in rumen contents at 0, 2.5 and 6 h after the commencement of the morning feeding in goats fed diets with 1.45% Mg(OH)<sub>2</sub> (■) or 0.6% elemental magnesium (□) after 28 d of adaption. Values are means (*n* 10), with their standard errors represented by vertical bars. \*\*\* *P* < 0.001.

Shannon and Simpson index (online Supplementary Table S4). The PC1 and PC2 explained 15.2 and 10.1% of variation in bacterial OTU, respectively, and the score plot did not show a distinct clustering of goats supplemented with Mg(OH)<sub>2</sub> or elemental Mg (online Supplementary Fig. S4). Bacteroidetes and Firmicutes were the most abundant ruminal bacterial phyla, and did not differ between treatments (Table 5). Few differences were observed in the abundance of genera with >1% abundance, which are shown in Table 5. Elemental Mg supplementation decreased the abundance of Bacteroidaceae (*P* = 0.03) and increased the abundance of Veillonellaceae (*P* = 0.06) and *Succiniclasticum* (*P* = 0.009) (Table 5).


**Discussion**

H<sub>2</sub> is produced during the reaction of elemental Mg with water: Mg + 2H<sub>2</sub>O → Mg(OH)<sub>2</sub> + H<sub>2</sub>. The Mg(OH)<sub>2</sub> so produced would be further hydrolysed in the rumen environment, in a reaction favoured by low pH: Mg(OH)<sub>2</sub> + 2H<sup>+</sup> → Mg<sup>2+</sup> + 2H<sub>2</sub>O. Ruminal Mg<sup>2+</sup> concentration was thus expected to be pH dependent, which agreed with the negative association observed between Mg<sup>2+</sup> concentration and pH. At all time points, ruminal Mg<sup>2+</sup> concentration was similar for both treatments of elemental Mg and Mg(OH)<sub>2</sub> supplementation, which suggests that rumen fluid passage rates to the lower tract were similar for both treatments. The reaction of elemental Mg with water resulting in H<sub>2</sub> formation might have been short-lived, or perhaps incomplete because of rumen outflow of elemental Mg, because dH<sub>2</sub> increased at +2.5 h, but not at +6 h, relative to the commencement of the morning feeding.

**Table 4.** Estimated Gibbs energy changes (kJ/reaction) of seven reaction pathways in the rumen of goats fed diets 1.45% Mg(OH)<sub>2</sub> or 0.6% elemental magnesium after 28 d of adaption (*n* 10)\* (Mean values with their standard errors)

Items	Mg(OH) <sub>2</sub>		Elemental Mg		<i>P</i>
	Mean	SE	Mean	SE	
<b>0 h relative to the commencement of morning feeding</b>					
Glucose → 2 acetate + 4 H <sub>2</sub>	-337	1.32	-341	2.14	0.10
Glucose → acetate + 1/2 butyrate + 3 H <sub>2</sub>	-331	1.13	-334	1.5	0.11
Glucose → butyrate + 2 H <sub>2</sub>	-324	1.05	-327	0.97	0.15
Glucose → acetate + propionate + H <sub>2</sub>	-407	0.92	-410	1.41	0.02
Glucose → 2/3 acetate + 4/3 propionate	-338	1.00	-339	0.94	0.10
Glucose + 2 H <sub>2</sub> → 2 propionate	-350	1.68	-350	1.25	0.99
CO <sub>2</sub> + 4 H <sub>2</sub> → CH <sub>4</sub>	-30.3	1.49	-27.7	1.79	0.24
<b>+2.5 h relative to the commencement of morning feeding</b>					
Glucose → 2 acetate + 4 H <sub>2</sub>	-324	2.56	-316	3.96	0.03
Glucose → acetate + 1/2 butyrate + 3 H <sub>2</sub>	-321	2.23	-314	3.24	0.03
Glucose → butyrate + 2 H <sub>2</sub>	-319	1.93	-315	2.56	0.04
Glucose → acetate + propionate + H <sub>2</sub>	-397	1.52	-393	2.29	0.07
Glucose → 2/3 acetate + 4/3 propionate	-334	0.67	-333	0.77	0.36
Glucose + 2 H <sub>2</sub> → 2 propionate	-351	1.24	-354	1.81	0.09
CO <sub>2</sub> + 4 H <sub>2</sub> → CH <sub>4</sub>	-39.8	2.14	-46.4	3.89	0.05
<b>+6 h relative to the commencement of morning feeding</b>					
Glucose → 2 acetate + 4 H <sub>2</sub>	-328	1.13	-329	2.25	0.67
Glucose → acetate + 1/2 butyrate + 3 H <sub>2</sub>	-324	1.15	-325	1.93	0.67
Glucose → butyrate + 2 H <sub>2</sub>	-321	1.22	-322	1.67	0.68
Glucose → acetate + propionate + H <sub>2</sub>	-401	0.79	-402	1.18	0.50
Glucose → 2/3 acetate + 4/3 propionate	-335	0.76	-336	0.51	0.52
Glucose + 2 H <sub>2</sub> → 2 propionate	-350	1.23	-350	1.29	0.96
CO <sub>2</sub> + 4 H <sub>2</sub> → CH <sub>4</sub>	-37.0	1.24	-36.0	2.27	0.70

\* Water is omitted for simplicity.



**Fig. 2.** Relationships between 2.5 h post-feeding dissolved hydrogen ( $dH_2$ ) and select microorganisms (a, copy number of fungi; b, copy number of methanogens; c, relative abundance of *Selenomonas ruminantium*), as determined by RT-PCR, in goats ( $n$  10) fed diets with 1.45%  $Mg(OH)_2$  or 0.6% elemental Mg after 28 d of adaption. Each point represents one goat, with a total of 20 data points.  $\Delta$ ,  $Mg(OH)_2$  treatment;  $\circ$ , elemental magnesium treatment.

**Table 5.** Select bacterial phyla and genera, as determined by 16S rRNA pyrosequencing, in the rumen contents of growing goats fed diets with 1.45%  $Mg(OH)_2$  or 0.6% elemental magnesium after 28 d of adaption ( $n$  10)\* (Mean values with their standard errors)

Phyla	Genera	% of sequences				P
		$Mg(OH)_2$		Elemental Mg		
		Mean	SE	Mean	SE	
Bacteroidetes		63.8	2.96	62.1	2.09	0.61
	Bacteroidaceae†	0.871	0.139	0.532	0.109	0.03
Firmicutes		26.2	2.54	28.3	2.49	0.57
	Veillonellaceae‡	0.486	0.142	0.883	0.198	0.06
	<i>Succiniclasticum</i>	0.987	0.217	2.88	0.556	0.009

\* Only the most abundant phyla and the genera that differed ( $P < 0.10$ ) between treatments are shown.

† Unknown genera within family Bacteroidaceae.

‡ Unknown genera within family Veillonellaceae.

It is understood that ruminal  $H_2$  accumulation hampers the oxidation of reduced electron carriers, such as NADH, to form  $H_2$ , leading to impaired glycolysis, decreased microbial ATP generation and fibre degradation in the rumen<sup>(32)</sup>. However, the rumen microbial ecosystem might exhibit the capability to cope with elevated  $H_2$ . Previous *in vivo* studies indicated that  $H_2$  accumulation under methanogenesis inhibition did not affect feed intake, total tract digestibility and meat or milk production in ruminants<sup>(7,33,34)</sup>. In the current study, we also did not find differences in feed intake and digestibility in goats fed diets supplemented with elemental Mg and  $Mg(OH)_2$ , but elevated  $dH_2$  resulting from elemental Mg supplementation decreased total VFA concentration in the rumen, which suggests that fermentation might have been impaired by elevated rumen  $dH_2$ . At first sight, it is difficult to reconcile the lack of effect of elevated  $dH_2$  on digestibility with its negative effects on total VFA concentration. However, both measurements are proxies of digestion and fermentation events. We measured apparent overall tract digestibility, but any changes in microbial biomass in faeces could have masked differences in true overall digestibility. Similarly, we measured VFA concentration in the rumen, but we did not measure actual fluxes of VFA production. A more complete understanding of the effects of

elevated  $dH_2$  would require a more in-depth study of the dynamics of digestion, microbial biomass production and fluxes of VFA production in the rumen.

The rumen VFA profile varies widely depending on the types of carbohydrates fermented. Diets rich in readily degraded starch generally stimulate propionate production, whereas on the other hand cellulose is fermented preferentially to acetate<sup>(35)</sup>. These shifts of fermentation pathways are associated with the concentration of ruminal  $dH_2$ <sup>(5)</sup>. Lower  $dH_2$  facilitates  $H_2$  generation through acetate production, whereas greater  $dH_2$  facilitates the disposition of electrons derived from fermentation into propionate production<sup>(2)</sup>. In this study, we confirmed rumen  $dH_2$  as a direct effector of the shift of glucose fermentation pathways, as elevation of  $dH_2$  caused by elemental Mg supplementation decreased acetate molar percentage and increased propionate molar percentage. This study allowed isolating the effects of elevated  $dH_2$  on fermentation from other effects caused by changes in the diet.

Elevated  $dH_2$  facilitated electron-incorporating glucose fermentation pathways, which agreed with the negative correlation between  $dH_2$  and  $R_{NH_2}$ . The shifts in fermentation pathways caused by elevated  $dH_2$  resulting from elemental Mg

supplementation also agreed with the effects of elemental Mg supplementation on  $\Delta G$  of the main fermentation pathways. The  $\Delta G$  of glucose fermentation to acetate and  $H_2$  was increased, and to propionate was decreased, by elemental Mg supplementation, along with the observed elevated  $dH_2$ . These results were in agreement with our previous study in dairy cows fed four types of carbohydrates<sup>(5)</sup>. The magnitude of the effects of elemental Mg supplementation on  $\Delta G$  of most rumen pathways at +2.5 h relative to the commencement of the morning feeding were rather small, which suggested that stimulation of pathways incorporating  $H_2$  might have taken place through more favourable kinetics of  $H_2$  uptake rather than through thermodynamic changes.

Changes in the composition of the rumen microbial community when  $gH_2$  accumulates as a consequence of methanogenesis inhibition have been demonstrated<sup>(6,7,36)</sup>. In dairy cows fed four types of carbohydrates, greater  $dH_2$  was associated with less acetate and lower number of  $H_2$  producers such as anaerobic fungi<sup>(5)</sup>. In that study, however, the concentration of  $dH_2$  was a consequence of the type of carbohydrate supplemented, and therefore changes in variables other than  $dH_2$  resulting from the dietary changes could have also influenced the abundance of microbial groups. To our knowledge, the present study was the first investigation on how elevated  $dH_2$  resulting from a non-fermentative source affected rumen microbiota, and we found that elevated  $dH_2$  generated by elemental Mg supplementation greatly decreased (–63.6%) the copy numbers of fungi. Decreased fungal numbers did not adversely affect fibre degradation, which might indicate that other fibrolytic organisms, such as *R. flavefaciens*, might have compensated for the negative effects of elemental Mg on fungi. *R. flavefaciens* produces  $H_2$ , but also succinate as a major product<sup>(37,38)</sup>, and it would be possible that *R. Flavefaciens* shifted fermentation towards succinate production under elevated  $H_2$  resulting from elemental Mg supplementation. That said, the relative abundance of *R. flavefaciens* was small, which casts doubt on the influence it might have had on fibre digestion.

The impact of increased  $dH_2$  generated by supplementation with elemental Mg on the rumen bacterial community composition was further analysed using high-throughput sequencing. Supplementation with elemental Mg increased species richness in terms of Ace and Chao 1 diversity, although total copy numbers of bacteria were similar for both treatments. However, the PCoA score plot indicated that ruminal bacterial community was not distinctly different between both treatments. In a previous study, elevated  $H_2$  was associated with increased Bacteroidetes: Firmicutes ratio when methanogenesis was inhibited by chloroform–cyclodextrin<sup>(39)</sup>. However, in the present study, elemental Mg supplementation decreased the abundance of family Bacteroidaceae, but did not alter the abundance of Bacteroidetes and Firmicutes. An interesting result was that genus *Succiniclasticum*, which belongs to family Firmicutes and is involved in converting succinate to propionate<sup>(40)</sup>, was 2-fold more abundant in goats that received elemental Mg supplementation. This result was in agreement with the increased propionate molar percentage in the rumen when  $dH_2$  concentration was increased by elemental Mg supplementation. In contrast, an unexpected observation was that elemental Mg supplementation and elevated  $dH_2$  decreased the

abundance of *S. ruminantium*, an important propionate-producing bacterium in the rumen<sup>(41)</sup>. The important increase in *Succiniclasticum* abundance might have compensated for the decreased abundance of *S. ruminantium* to produce propionate in this study, if in goats supplemented elemental Mg succinate conversion to propionate limited propionate production.

$H_2$  and  $CO_2$  are the main substrates for methanogenesis in the rumen, and the majority of  $H_2$  produced from carbohydrate fermentation is used in  $CH_4$  production<sup>(42)</sup>. In our study, we found that elemental Mg supplementation increased  $dCH_4$  concentration, the copy number of methanogens and  $CH_4$  emissions, indicating that increased  $dH_2$  generated by elemental Mg supplementation promoted  $CH_4$  production in the rumen of goats. Olijhoek *et al.*<sup>(11)</sup> also reported that short-term infusion of  $H_2$  into the rumen increased  $CH_4$  production. Furthermore, we estimated that elemental Mg supplementation decreased  $\Delta G$  of methanogenesis at +2.5 h after the commencement of the morning feeding, indicating that elevated  $dH_2$  would have thermodynamically favoured growth and metabolism of methanogens at that time point. Dissolved  $H_2$  concentration was positively correlated with copy numbers of methanogens, but not with  $CH_4$  emissions. This suggests that ruminal  $dH_2$  might have exceeded the capacity of methanogens to take up  $H_2$  in the treatment with elemental Mg supplementation, so that methanogenic population and activity might have been limiting for  $CH_4$  generation in the rumen.

In summary, goats adapted to elevated ruminal  $dH_2$  generated by elemental Mg supplementation had lower rumen total VFA concentration, copy number of fungi and greater propionate molar percentage and  $CH_4$  emissions when compared with goats fed the control diet supplemented with  $Mg(OH)_2$ . Elevation of ruminal  $dH_2$  concentration seemed to inhibit rumen fermentation and altered fermentation pathways and some microbial groups, without affecting total tract digestibility. Effects of extra  $H_2$  obtained by supplementing elemental Mg were short-lived, because elevated  $dH_2$  had returned to control concentration by +6 h relative to the commencement of the morning feeding. Furthermore, the extent of elevated ruminal  $dH_2$  concentration caused by elemental Mg supplementation varied widely among the ten goats, and the goat with the greatest  $dH_2$  concentration (17 mmol/l) also had the greatest methanogen population and the lowest VFA concentration, acetate molar percentage, acetate:propionate ratio and copy number of fungi. This study provides insights on the effects of elevated  $dH_2$  on fermentation in the rumen and methanogenesis, although high dietary Mg concentration in both treatments might have altered the availability of other minerals and exerted some postabsorptive effects in goats. More studies with sustained and consistent elevated  $dH_2$  are needed to further elucidate how elevated  $dH_2$  concentration affects the rumen ecosystem, fermentation and digestion.

### Acknowledgements

This work was supported by National Natural Science Foundation of China (grant nos 31561143009, 31320103917 and 31472133), National key research and development program of China (grant no. 2016YFD0500504), Youth Innovation Promotion Association

CAS (2016327), China Agriculture Research System (grant no. CARS-37) and Hunan province science and technology plan (grant no. 2015WK3043).

M. W., R. W., H. X. M., D. L. L. and X. M. Z. conducted the research; M. W., R. W. and E. M. U. analysed the data; and M. W., E. M. U., K. A. B and Z. L. T. wrote the paper. M. W. and Z. L. T. designed the research and had primary responsibility for the final content. All authors read and approved the final manuscript.

The authors declare that there are no conflicts of interest.

### Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114517002161>

### References

- Vyas D, McGinn SM, Duval SM, *et al.* (2016) Effects of sustained reduction of enteric methane emissions with dietary supplementation of 3-nitrooxypropanol on growth performance of growing and finishing beef cattle1. *J Anim Sci* **94**, 2024–2034.
- Janssen PH (2010) Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim Feed Sci Technol* **160**, 1–22.
- Joblin KN (1999) Ruminant acetogens and their potential to lower ruminant methane emissions. *Aust J Agric Res* **50**, 1307–1313.
- Ungerfeld EM & Kohn RA (2006) The role of thermodynamics in control of ruminal fermentation. In *Ruminant Physiology Digestion, Metabolism and Impact of Nutrition on Gene Expression, Immunology and Stress*, pp. 55–85 [K Sejrsen, T Hvelpund and MO Nielsen, editors]. Wageningen: Wageningen Academic Publishers.
- Wang M, Wang R, Xie T, *et al.* (2016) Shifts in rumen fermentation and microbiota are associated with dissolved ruminal hydrogen concentrations in lactating dairy cows fed different types of carbohydrates. *J Nutr* **146**, 1714–1721.
- Martinez-Fernandez G, Denman SE, Yang C, *et al.* (2016) Methane inhibition alters the microbial community, hydrogen flow, and fermentation response in the rumen of cattle. *Front Microbiol* **7**, 1122.
- Mitsumori M, Shinkai T, Takenaka A, *et al.* (2012) Responses in digestion, rumen fermentation and microbial populations to inhibition of methane formation by a halogenated methane analogue. *Br J Nutr* **108**, 482–491.
- Patra AK & Yu Z (2013) Effects of gas composition in headspace and bicarbonate concentrations in media on gas and methane production, degradability, and rumen fermentation using *in vitro* gas production techniques. *J Dairy Sci* **96**, 4592–4600.
- Klop G, Dijkstra J, Dieho K, *et al.* (2017) Enteric methane production in lactating dairy cows with continuous feeding of essential oils or rotational feeding of essential oils and lauric acid. *J Dairy Sci* **100**, 3563–3575.
- Qiao JY, Tan ZL, Guan LL, *et al.* (2015) Effects of hydrogen in headspace and bicarbonate in media on rumen fermentation, methane production and methanogenic population using *in vitro* gas production techniques. *Anim Feed Sci Technol* **206**, 19–28.
- Olijhoek DW, Hellwing ALF, Weisbjerg MR, *et al.* (2016) Effect of short-term infusion of hydrogen on enteric gas production and rumen environment in dairy cows. *Anim Prod Sci* **56**, 466–471.
- Broudicou LP, Offner A & Sauvant D (2014) Effects of inoculum source, pH, redox potential and headspace di-hydrogen on rumen *in vitro* fermentation yields. *Animal* **8**, 931–937.
- Wang M, Sun XZ, Janssen PH, *et al.* (2014) Responses of methane production and fermentation pathways to the increased dissolved hydrogen concentration generated by eight substrates in *in vitro* ruminal cultures. *Anim Feed Sci Technol* **194**, 1–11.
- Wang M, Ungerfeld EM, Wang R, *et al.* (2016) Supersaturation of dissolved hydrogen and methane in rumen of Tibetan sheep. *Front Microbiol* **7**, 850.
- Ng F, Kittelmann S, Patchett ML, *et al.* (2016) An adhesin from hydrogen-utilizing rumen methanogen *Methanobrevibacter ruminantium* M1 binds a broad range of hydrogen-producing microorganisms. *Environ Microbiol* **18**, 3010–3021.
- Wang M, Wang R, Janssen PH, *et al.* (2016) Sampling procedure for the measurement of dissolved hydrogen and volatile fatty acids in the rumen of dairy cows. *J Anim Sci* **94**, 1159–1169.
- Wang M, Wang R, Sun X, *et al.* (2015) A mathematical model to describe the diurnal pattern of enteric methane emissions from non-lactating dairy cows post-feeding. *Anim Nutr* **1**, 329–338.
- McGinn SM, Beauchemin KA, Coates T, *et al.* (2004) Methane emissions from beef cattle: effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *J Anim Sci* **82**, 3346–3356.
- Association of Official Analytical Chemists (editor) (1995) *Official Methods of Analysis*, 16th ed. Arlington, VA: AOAC.
- Van Soest PJ, Robertson JB & Lewis BA (1991) Symposium: carbohydrate methodology, metabolism and nutritional implications in dairy cattle. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* **74**, 3583–3597.
- Kartchner RJ & Theurer B (1981) Comparison of hydrolysis methods used in feed, digesta, and fecal starch analysis. *J Agri Food Chem* **29**, 8–11.
- Weatherburn MW (1967) Phenol-hypochlorite reaction for determination of ammonia. *Anal Chem* **39**, 971–974.
- Nelson N (1944) A photometric adaptation of the Somogyi method for the determination of glucose. *J Biol Chem* **153**, 375–380.
- Hille KT, Hetz SK, Rosendahl J, *et al.* (2016) Determination of Henry's constant, the dissociation constant, and the buffer capacity of the bicarbonate system in ruminal fluid. *J Dairy Sci* **99**, 369–385.
- Carroll JJ, Slupsky JD & Mather AE (1991) The solubility of carbon-dioxide in water at low-pressure. *J Phys Chem Ref Data* **20**, 1201–1209.
- Guyader J, Eugène M, Doreau M, *et al.* (2017) Tea saponin reduced methanogenesis *in vitro* but increased methane yield in lactating dairy cows. *J Dairy Sci* **100**, 1845–1855.
- Jiao J, Lu Q, Tan Z, *et al.* (2014) *In vitro* evaluation of effects of gut region and fiber structure on the intestinal dominant bacterial diversity and functional bacterial species. *Anaerobe* **28**, 168–177.
- Stevenson D & Weimer P (2007) Dominance of *Prevotella* and low abundance of classical ruminal bacterial species in the bovine rumen revealed by relative quantification real-time PCR. *Appl Microbiol Biotechnol* **75**, 165–174.
- Hackmann TJ & Firkins JL (2015) Electron transport phosphorylation in rumen butyrovibrios: unprecedented ATP yield for glucose fermentation to butyrate. *Front Microbiol* **6**, 622.

30. Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461.
31. DeSantis TZ, Hugenholtz P, Larsen N, *et al.* (2006) Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* **72**, 5069–5072.
32. Ungerfeld EM (2015) Shifts in metabolic hydrogen sinks in the methanogenesis-inhibited ruminal fermentation: a meta-analysis. *Front Microbiol* **6**, 37.
33. Hristov AN, Oh J, Giallongo F, *et al.* (2015) An inhibitor persistently decreased enteric methane emission from dairy cows with no negative effect on milk production. *PNAS* **112**, 10663–10668.
34. Romero-Perez A, Okine EK, McGinn SM, *et al.* (2015) Sustained reduction in methane production from long-term addition of 3-nitrooxypropanol to a beef cattle diet. *J Anim Sci* **93**, 1780–1791.
35. Bannink A, Kogut J, Dijkstra J, *et al.* (2006) Estimation of the stoichiometry of volatile fatty acid production in the rumen of lactating cows. *J Theor Biol* **238**, 36–51.
36. Shinkai T, Enishi O, Mitsumori M, *et al.* (2012) Mitigation of methane production from cattle by feeding cashew nut shell liquid. *J Dairy Sci* **95**, 5308–5316.
37. Stewart CS, Flint HJ & Bryant MP (1997) The rumen bacteria. In *The Rumen Microbial Ecosystem*, pp. 10–72 [PN Hobson and CS Stewart, editors]. London: Blackie Academic & Professional.
38. Philippeau C, Lettat A, Martin C, *et al.* (2017) Effects of bacterial direct-fed microbials on ruminal characteristics, methane emission, and milk fatty acid composition in cows fed high- or low-starch diets. *J Dairy Sci* **100**, 2637–2650.
39. Fonty G, Joblin K, Chavarot M, *et al.* (2007) Establishment and development of ruminal hydrogenotrophs in methanogen-free lambs. *Appl Environ Microbiol* **73**, 6391–6403.
40. Scheifinger CC, Linehan B & Wolin MJ (1975) H<sub>2</sub> production by *Selenomonas ruminantium* in the absence and presence of methanogenic bacteria. *Appl Microbiol* **29**, 480–483.
41. Sawanon S, Koike S & Kobayashi Y (2011) Evidence for the possible involvement of *Selenomonas ruminantium* in rumen fiber digestion. *FEMS Microbiol Lett* **325**, 170–179.
42. McAllister TA & Newbold CJ (2008) Redirecting rumen fermentation to reduce methanogenesis. *Aust J Exp Agric* **48**, 7–13.