

CORRIGENDUM

Thin Film Nanoelectronic Probe for Protein Detection – **CORRIGENDUM**

Rahim Esfandyarpour, Mehdi Javanmard, Zahra Koochak, Hesaam Esfandyarpour,
James S. Harris, and Ronald W. Davis

doi: 10.1557/opl.2013.660, Published by Materials Research Society, 5 June 2013.

The article's reference list has been updated with the proper citation format.

Reference

Rahim Esfandyarpour, Mehdi Javanmard, Zahra Koochak, Hesaam Esfandyarpour,
James S. Harris and Ronald W. Davis (2013). Thin Film Nanoelectronic Probe for
Protein Detection. MRS Proceedings, 1572, mrss13-1572-ss01-10
doi:10.1557/opl.2013.660.

REFERENCES

1. Dunphy, W. G.; Kumagai, A., The cdc25 protein contains an intrinsic phosphatase activity. *Cell* 1991, 67 (1), 189-196.
2. Stillman, B., Smart machines at the DNA replication fork. *Cell* 1994, 78 (5), 725-728.
3. Stevens, M. M.; George, J. H., Exploring and engineering the cell surface interface. *Science* 2005, 310 (5751), 1135-1138.
4. Astumian, R. D., Adiabatic pumping mechanism for ion motive ATPases. *Physical review letters* 2003, 91 (11), 118102.
5. Esfandyarpour, R.; Esfandyarpour, H.; Javanmard, M.; Harris, J. S.; Davis, R. W., Microneedle Biosensor: A Method for Direct Label-free Real Time Protein Detection. *Sensors and Actuators B: Chemical* 2012.
6. Esfandyarpour, R.; Esfandyarpour, H.; Javanmard, M.; Harris, J. S.; Davis, R. W. In *Electrical Detection of Protein Biomarkers Using Nanoneedle Biosensors*, MRS Proceedings, Cambridge Univ Press: 2012.
7. Koritsas, V. M.; Atkinson, H. J., An assay for detecting nanogram levels of proteolytic enzymes. *Analytical biochemistry* 1995, 227 (1), 22-26.
8. van Oss, C. J.; Giese, R. F.; Bronson, P. M.; Docoslis, A.; Edwards, P.; Ruyechan, W. T., Macroscopic-scale surface properties of streptavidin and their influence on aspecific interactions between streptavidin and dissolved biopolymers. *Colloids and Surfaces B: Biointerfaces* 2003, 30 (1), 25-36.
9. Butler, J.; Ni, L.; Brown, W.; Joshi, K.; Chang, J.; Rosenberg, B.; Voss Jr, E., The immunochemistry of sandwich ELISAs—VI. Greater than 90% of monoclonal and 75% of polyclonal anti-fluorescyl capture antibodies (Cabs) are denatured by passive adsorption. *Molecular immunology* 1993, 30 (13), 1165-1175.
10. Nordlund, H., *Avidin engineering: modification of function, oligomerization, stability and structure topology*. Jyväskylän yliopisto: 2003.
11. Salama, A. D.; Dougan, T.; Levy, J. B.; Cook, H. T.; Morgan, S. H.; Naudeer, S.; Maidment, G.; George, A. J.; Evans, D.; Lightstone, L., Goodpasture's disease in the absence of circulating anti-glomerular basement membrane antibodies as detected by standard techniques. *American journal of kidney diseases* 2002, 39 (6), 1162-1167.
12. Pitera, J. W.; Falta, M.; van Gunsteren, W. F., Dielectric properties of proteins from simulation: the effects of solvent, ligands, pH, and temperature. *Biophysical journal* 2001, 80 (6), 2546-2555.
13. Gray, H. B.; Winkler, J. R., Electron tunneling through proteins. *Quarterly reviews of biophysics* 2003, 36 (3), 341-372.
14. Larsson, S., Electron transfer in proteins. *J. Chem. Soc., Faraday Trans. 2* 1983, 79 (9), 1375-1388.