

THE ÉTALE GROUPOID OF AN INVERSE SEMIGROUP AS A GROUPOID OF FILTERS

M. V. LAWSON[✉], S. W. MARGOLIS and B. STEINBERG

(Received 27 June 2011; accepted 9 April 2012; first published online 15 May 2013)

Communicated by M. Jackson

This paper is dedicated to the memory of our friend and colleague Steve Haataja.

Abstract

Paterson showed how to construct an étale groupoid from an inverse semigroup using ideas from functional analysis. This construction was later simplified by Lenz. We show that Lenz's construction can itself be further simplified by using filters: the topological groupoid associated with an inverse semigroup is precisely a groupoid of filters. In addition, idempotent filters are closed inverse subsemigroups and so determine transitive representations by means of partial bijections. This connection between filters and representations by partial bijections is exploited to show how linear representations of inverse semigroups can be constructed from the groups occurring in the associated topological groupoid.

2010 Mathematics subject classification: primary 20M18; secondary 20M30, 18B40, 46L05.

Keywords and phrases: Étale groupoids, inverse semigroups, filters.

1. Introduction and motivation

In his influential book, Renault [20] showed how to construct C^* -algebras from locally compact topological groupoids. This can be seen as a far-reaching generalization of both commutative C^* -algebras and finite-dimensional C^* -algebras. From this perspective, topological groupoids can be viewed as ‘noncommutative topological spaces’. Renault also showed that, in addition to groupoids and C^* -algebras, a third class of structures naturally intervenes: inverse semigroups. Local bisections of topological groupoids form inverse semigroups and, conversely, inverse semigroups can be used to construct topological groupoids.

The relationship between inverse semigroups and topological groupoids can be seen as a generalization of that between (pre)sheaves of groups and their corresponding display spaces, since an inverse semigroup with central idempotents is a presheaf of groups over its semilattice of idempotents. This relationship has been investigated by

a number of authors, notably Paterson [18], Kellendonk [4–7] and Resende [21]. Our paper is related to Paterson’s work but mediated through a more recent redaction due to Daniel Lenz [15].

We prove two main results. First, we show that Lenz’s construction of the topological groupoid can be interpreted entirely in terms of down-directed cosets on inverse semigroups—these are precisely the filters in an inverse semigroup. Such filters arise naturally from those transitive actions which we term ‘universal’. Second, we show how representations of an inverse semigroup can be constructed from the groups occurring in the associated topological groupoid. This is related to Steinberg’s results on constructing finite-dimensional representations of inverse semigroups using groupoid techniques described in [28]. The first result proved in this paper has already been developed further in [11, 12].

Lenz [15] was the main spur that led us to write this paper; but in the course of doing so, we realized that the first four chapters of Ruyle’s unpublished thesis [22] could be viewed as a major contribution to the aims of this paper in the case of free inverse monoids. Ruyle’s work has proved indispensable for our Section 2. In addition, Leech [14], with its emphasis on the order-theoretic structure of inverse semigroups, can be seen with mathematical hindsight to be a precursor of our approach. Last, but not least, Boris Schein in a number of seminars talked about ways of constructing *infinitesimal elements* of an inverse semigroup: the maximal filters of an inverse semigroup can be regarded as just that [24, 25].

For general inverse semigroup theory we refer the reader to [9]. However, we note the following. The product in a semigroup will usually be denoted by concatenation but sometimes we shall use \cdot for emphasis; we shall also use it to denote actions. In an inverse semigroup S we define

$$\mathbf{d}(s) = s^{-1}s \quad \text{and} \quad \mathbf{r}(s) = ss^{-1}.$$

Green’s relation \mathcal{H} can be defined in terms of this notation as follows: $s\mathcal{H}t$ if and only if $\mathbf{d}(s) = \mathbf{d}(t)$ and $\mathbf{r}(s) = \mathbf{r}(t)$. If e is an idempotent in a semigroup S then G_e will denote the \mathcal{H} -class in S containing e ; this is a maximal subgroup. The natural partial order will be the only partial order considered when we deal with inverse semigroups. If $X \subseteq S$ then $E(X)$ denotes the set of idempotents in X . An inverse subsemigroup of S is said to be *wide* if it contains all the idempotents of S . A *primitive idempotent* e in an inverse semigroup S with zero is one with the property that if $f \leq e$ then either $f = e$ or $f = 0$. Let S be an inverse semigroup. The *minimum group congruence* σ on S is defined by $a \sigma b$ if and only if $c \leq a, b$ for some $c \in S$. This congruence has the property that S/σ is a group, and if ρ is any congruence on S for which S/ρ is a group, then $\sigma \subseteq \rho$. We denote by $\sigma^{\mathfrak{h}}$ the associated natural homomorphism $S \rightarrow S/\sigma$. See [9] for more information on this important congruence.

After an early version of this paper was written, we discovered that Jonathon Funk and Pieter Hofstra had independently arrived at what we call universal actions, and which they call torsors [1]. They show that these correspond exactly to the points of the classifying topos of the inverse semigroup. Further connections between our work

and their work will be explored in a forthcoming paper by Funk, Hofstra and the third author. In particular, we connect the filter construction of Paterson's groupoid with the soberification of the inductive groupoid of the inverse semigroup and the soberification of the inverse semigroup. We also show that actions of the inverse semigroup on sober spaces correspond to actions of the soberification of the inductive groupoid on sober spaces.

2. The structure of transitive actions

In this section, we begin by reviewing the general theory of representations of inverse semigroups by partial permutations. [19, Ch. IV, Section 4] contains an exposition of this elementary theory and we refer the reader there for any proofs we omit. We also incorporate some results by Ruyle from [22] which can be viewed as anticipating some of the ideas in this paper. We then introduce the concept of universal transitive actions which provides the connection with the work of Lenz to be explained in Section 3.

2.1. The classical theory. A *representation* of an inverse semigroup by means of partial bijections (or partial permutations) is a homomorphism $\theta: S \rightarrow I(X)$ to the symmetric inverse monoid on a set X . A representation of an inverse semigroup in this sense leads to a corresponding notion of an action of the inverse semigroup S on the set X : the associated action is defined by $s \cdot x = \theta(s)(x)$, if x belongs to the set-theoretic domain of $\theta(s)$. The action is therefore a partial function from $S \times X$ to X mapping (s, x) to $s \cdot x$ when $s \cdot x$ exists satisfying the following two axioms.

(A1) If $e \cdot x$ exists, where e is an idempotent, then $e \cdot x = x$.

(A2) $(st) \cdot x$ exists if and only if $s \cdot (t \cdot x)$ exists, in which case they are equal.

It is easy to check that representations and actions are different ways of describing the same thing. For convenience, we shall use the words 'action' and 'representation' interchangeably: if we say that the inverse semigroup S acts on a set X then this will imply the existence of an appropriate homomorphism from S to $I(X)$. If S acts on X we shall often refer to X as a *space* or as an S -*space* and its elements as *points*. A subset $Y \subseteq X$ closed under the action is called a *subspace*. Disjoint unions of actions are again actions. An action is said to be *effective* if for each $x \in X$ there is $s \in S$ such that $s \cdot x$ exists. We shall assume that all our actions are effective. An effective action of an inverse semigroup S on the set X induces an equivalence relation \sim on the set X when we define $x \sim y$ if and only if $s \cdot x = y$ for some $s \in S$. The action is said to be *transitive* if \sim is $X \times X$. Just as in the theory of permutation representations of groups, every representation of an inverse semigroup is a disjoint union of transitive representations. Thus the transitive representations of inverse semigroups are of especial significance.

Let X and Y be S -spaces. A *morphism* from X to Y is a function $\alpha: X \rightarrow Y$ such that $s \cdot x$ exists implies that $s \cdot \alpha(x)$ exists and $\alpha(s \cdot x) = s \cdot \alpha(x)$. A *strong morphism* from X to Y is a function $\alpha: X \rightarrow Y$ such that $s \cdot x$ exists if and only if $s \cdot \alpha(x)$ exists, and if $s \cdot x$ exists then $\alpha(s \cdot x) = s \cdot \alpha(x)$. Bijective strong morphisms are called *equivalences*. The proofs of the following two lemmas are straightforward.

LEMMA 2.1. *The class of S -spaces, for a fixed inverse semigroup S , together with their morphisms (respectively, strong morphisms) forms a category.*

LEMMA 2.2. *Let S be an inverse semigroup acting on X , Y and Z .*

- (1) *The image of a strong morphism $\alpha: X \rightarrow Y$ is a subspace of Y .*
- (2) *If X and Y are transitive S -spaces and $\alpha: X \rightarrow Y$ is a strong morphism then α is surjective.*

If we fix an inverse semigroup S there are a number of categories of actions associated with it: actions and morphisms, actions and strong morphisms, transitive actions and morphisms, and transitive actions and strong morphisms. As we indicated above, these two categories of transitive actions will be of central importance.

A *congruence* on X is an equivalence relation \sim on the set X such that if $x \sim y$ and if $s \cdot x$ exists and $s \cdot y$ exists then $s \cdot x \sim s \cdot y$. A *strong congruence* on X is an equivalence relation \approx on the set X such that if $x \approx y$ and $s \in S$ we have that $s \cdot x$ exists if and only if $s \cdot y$ exists, and if the actions are defined then $s \cdot x \approx s \cdot y$.

Strong morphisms and strong congruences are united by a classical first isomorphism theorem. Recall that the *kernel* of a function is the equivalence relation induced on its domain. The proofs of the following are routine.

PROPOSITION 2.3. *Let S be an inverse semigroup and let X and Y be S -spaces.*

- (1) *Let $\alpha: X \rightarrow Y$ be a strong morphism. Then the kernel of α is a strong congruence.*
- (2) *Let \sim be a strong congruence on X . Denote the \sim -class containing the element x by $[x]$. Define $s \cdot [x] = [s \cdot x]$ if $s \cdot x$ exists. Then this defines an action S on the set of \sim -congruence classes X/\sim and the natural map $v: X \rightarrow X/\sim$ is a strong morphism.*
- (3) *Let $\alpha: X \rightarrow Y$ be a strong morphism, let its kernel be \sim and let $v: X \rightarrow X/\sim$ be the associated natural map. Then there is a unique injective strong morphism β from X/\sim to Y such that $\beta v = \alpha$.*

The above result tells us that the category of transitive representations of a fixed inverse semigroup with strong morphisms between them has a particularly nice structure.

We may analyze transitive actions of inverse semigroups in a way generalizing the relationship between transitive group actions and subgroups. To describe this relationship we need some definitions. If $A \subseteq S$ is a subset then define

$$A^\uparrow = \{s \in S : a \leq s \text{ for some } a \in A\}.$$

If $A = A^\uparrow$ then A is said to be *closed (upwards)*.

Let X be an S -space. Fix a point $x \in X$, and consider the set S_x consisting of all $s \in S$ such that $s \cdot x = x$. We call S_x the *stabilizer* of the point x .

We do not assume in this paper that homomorphisms of inverse semigroups with zero preserve the zero. If $\theta: S \rightarrow I(X)$ is a representation that does preserve zero then the zero of S is mapped to the empty function of $I(X)$. Clearly, the empty function

cannot belong to any stabilizer. We say that a closed inverse subsemigroup is *proper* if it does not contain a zero. In the theory we summarize below, proper closed inverse subsemigroups arise from actions where the zero acts as the empty partial function.

Now let $y \in X$ be any point. By transitivity, there is an element $s \in S$ such that $s \cdot x = y$. Observe that because $s \cdot x$ is defined so too is $s^{-1}s \cdot x$ and that $s^{-1}s \in S_x$. The set of all elements of S which map x to y is $(sS_x)^\uparrow$.

Let H be a closed inverse subsemigroup of S . Define a *left coset* of H to be a set of the form $(sH)^\uparrow$ where $s^{-1}s \in H$. We give the proof of the following for completeness.

LEMMA 2.4. *Let S be an inverse semigroup.*

- (1) *Two cosets $(sH)^\uparrow$ and $(tH)^\uparrow$ are equal if and only if $s^{-1}t \in H$.*
- (2) *If $(sH)^\uparrow \cap (tH)^\uparrow \neq \emptyset$ then $(sH)^\uparrow = (tH)^\uparrow$.*

PROOF. (1) Suppose that $(sH)^\uparrow = (tH)^\uparrow$. Then $t \in (sH)^\uparrow$ and so $sh \leq t$ for some $h \in H$. Thus $s^{-1}sh \leq s^{-1}t$. But $s^{-1}sh \in H$ and H is closed and so $s^{-1}t \in H$. Conversely, suppose that $s^{-1}t \in H$. Then $s^{-1}t = h$ for some $h \in H$ and so $sh = ss^{-1}t \leq t$. It follows that $tH \subseteq sH$ and so $(tH)^\uparrow \subseteq (sH)^\uparrow$. The reverse inclusion follows from the fact that $t^{-1}s \in H$ since H is closed under inverses.

(2) Suppose that $a \in (sH)^\uparrow \cap (tH)^\uparrow$. Then $sh_1 \leq a$ and $th_2 \leq a$ for some $h_1, h_2 \in H$. Thus $s^{-1}sh_1 \leq s^{-1}a$ and $t^{-1}th_2 \leq t^{-1}a$. Hence $s^{-1}a, t^{-1}a \in H$. It follows that $s^{-1}aa^{-1}t \in H$, but $s^{-1}aa^{-1}t \leq s^{-1}t$. This gives the result by (1) above. □

We denote by S/H the set of all left cosets of H in S . The inverse semigroup S acts on the set S/H when we define

$$a \cdot (sH)^\uparrow = (asH)^\uparrow \text{ whenever } \mathbf{d}(as) \in H.$$

This defines a transitive action. The following is [19, Lemma IV.4.9] and [3, Proposition 5.8.5].

THEOREM 2.5. *Let S act transitively on the set X . Then the action is equivalent to the action of S on the set S/S_x where x is any point of X .*

The following is [19, Proposition IV.4.13].

PROPOSITION 2.6. *If H and K are any closed inverse subsemigroups of S then they determine equivalent actions if and only if there exists $s \in S$ such that*

$$sHs^{-1} \subseteq K \quad \text{and} \quad s^{-1}Ks \subseteq H.$$

The above relationship between closed inverse subsemigroups is called *conjugacy* and defines an equivalence relation on the set of closed inverse subsemigroups. The proof of the following is given for completeness.

LEMMA 2.7. *H and K are conjugate if and only if*

$$(sHs^{-1})^\uparrow = K \quad \text{and} \quad (s^{-1}Ks)^\uparrow = H.$$

PROOF. Let H and K be conjugate. Let $e \in H$ be any idempotent. Then $ses^{-1} \in K$. But $ses^{-1} \leq ss^{-1}$ and so $ss^{-1} \in K$. Similarly $s^{-1}s \in H$. We have that $sHs^{-1} \subseteq K$ and so $(sHs^{-1})^\uparrow \subseteq K$. Let $k \in K$. Then $s^{-1}ks \in H$ and $s(s^{-1}ks)s^{-1} \in sHs^{-1}$ and $s(s^{-1}ks)s^{-1} \leq k$. Thus $(sHs^{-1})^\uparrow = K$, as required. The converse is immediate. \square

Thus to study the transitive actions of an inverse semigroups S it is enough to study the closed inverse subsemigroups of S up to conjugacy.

The following result is motivated by Lemma 2.16 of Ruyle’s thesis [22] and brings morphisms and strong morphisms back into the picture.

THEOREM 2.8. *Let S be an inverse semigroup acting transitively on the sets X and Y , and let $x \in X$ and $y \in Y$. Let S_x and S_y be the stabilizers in S of x and y respectively.*

- (1) *There is a (unique) morphism $\alpha: X \rightarrow Y$ such that $\alpha(x) = y$ if and only if $S_x \subseteq S_y$.*
- (2) *There is a (unique) strong morphism $\alpha: X \rightarrow Y$ such that $\alpha(x) = y$ if and only if $S_x \subseteq S_y$ and $E(S_x) = E(S_y)$.*

PROOF. (1) We begin by proving uniqueness. Let $\alpha, \beta: X \rightarrow Y$ be morphisms such that $\alpha(x) = \beta(x) = y$. Let $x' \in X$ be arbitrary. By transitivity there exists $a \in S$ such that $x' = a \cdot x$. By the definition of morphisms, we have that $a \cdot \alpha(x)$ exists and $a \cdot \beta(x)$ exists and that

$$\alpha(x') = \alpha(a \cdot x) = a \cdot \alpha(x)$$

and

$$\beta(x') = \beta(a \cdot x) = a \cdot \beta(x).$$

But by assumption $\alpha(x) = \beta(x) = y$ and so $\alpha(x') = \beta(x')$. It follows that $\alpha = \beta$.

Let $\alpha: X \rightarrow Y$ be a morphism such that $\alpha(x) = y$. Let $s \in S_x$. Then $s \cdot x$ exists and $s \cdot x = x$. By the definition of morphism, it follows that $s \cdot \alpha(x)$ exists and that $\alpha(s \cdot x) = s \cdot \alpha(x)$. But $s \cdot x = x$ and so $\alpha(x) = s \cdot \alpha(x)$. Hence $s \cdot y = y$. We have therefore proved that $s \in S_y$, and so $S_x \subseteq S_y$.

Suppose now that $S_x \subseteq S_y$. We have to define a morphism $\alpha: X \rightarrow Y$ such that $\alpha(x) = y$. We start by defining $\alpha(x) = y$. Let $x' \in X$ be any point in X . Then $x' = a \cdot x$ for some $a \in S$. We need to show that $a \cdot y$ exists. Since $a \cdot x$ exists we know that $a^{-1}a \cdot x$ exists and this is equal to x . It follows that $a^{-1}a \in S_x$ and so $a^{-1}a \in S_y$, by assumption. Thus $a^{-1}a \cdot y$ exists and is equal to y . But from the existence of $a^{-1}a \cdot y$ we can deduce the existence of $a \cdot y$. We would therefore like to define $\alpha(x') = a \cdot y$. We have to check that this is well defined. Suppose that $x' = a \cdot x = b \cdot x$. Then $b^{-1}a \cdot x = x$ and so $b^{-1}a \in S_x$. By assumption, $b^{-1}a \in S_y$ and so $b^{-1}a \cdot y = y$. Thus $bb^{-1}a \cdot y = b \cdot y$ and $bb^{-1}a \cdot y = bb^{-1} \cdot (a \cdot y) = a \cdot y$. Thus $a \cdot y = b \cdot y$. It follows that α is a well-defined function mapping x to y . It remains to show that α is a morphism. Suppose that $s \cdot x'$ is defined. By assumption, there exists $a \in S$ such that $x' = a \cdot x$. By definition $\alpha(x') = a \cdot y$. We have that $s \cdot x' = s \cdot (a \cdot x) = sa \cdot x$. By definition, $\alpha(s \cdot x') = sa \cdot y$. But $sa \cdot y = s \cdot (a \cdot y) = s \cdot \alpha(x')$. Hence $\alpha(s \cdot x') = s \cdot \alpha(x')$, as required.

(2) We begin by proving uniqueness. Let $\alpha, \beta: X \rightarrow Y$ be strong morphisms such that $\alpha(x) = \beta(x) = y$. Let $x' \in X$ be arbitrary. By transitivity there exists $a \in S$ such

that $x' = a \cdot x$. By the definition of strong morphisms we have that $a \cdot \alpha(x)$ exists and $a \cdot \beta(x)$ exists and that

$$\alpha(x') = \alpha(a \cdot x) = a \cdot \alpha(x)$$

and

$$\beta(x') = \beta(a \cdot x) = a \cdot \beta(x).$$

But by assumption, $\alpha(x) = \beta(x) = y$ and so $\alpha(x') = \beta(x')$. It follows that $\alpha = \beta$.

Next we prove existence. Suppose that $S_x \subseteq S_y$ and $E(S_x) = E(S_y)$. We have to define a strong morphism $\alpha: X \rightarrow Y$ such that $\alpha(x) = y$. We start by defining $\alpha(x) = y$. Let $x' \in X$ be any point in X . Then $x' = a \cdot x$ for some $a \in S$. We need to show that $a \cdot y$ exists. Since $a \cdot x$ exists we know that $a^{-1}a \cdot x$ exists and this is equal to x . It follows that $a^{-1}a \in S_x$ and so $a^{-1}a \in S_y$, by assumption. Thus $a^{-1}a \cdot y$ exists and is equal to y . But from the existence of $a^{-1}a \cdot y$ we can deduce the existence of $a \cdot y$. We therefore define $\alpha(x') = a \cdot y$. We have to check that this is well defined. Suppose that $x' = a \cdot x = b \cdot x$. Then $b^{-1}a \cdot x = x$ and so $b^{-1}a \in S_x$. By assumption, $b^{-1}a \in S_y$ and so $b^{-1}a \cdot y = y$. Thus $bb^{-1}a \cdot y = b \cdot y$ and $bb^{-1}a \cdot y = bb^{-1} \cdot (a \cdot y) = a \cdot y$. Thus $a \cdot y = b \cdot y$. It follows that α is a well-defined function mapping x to y .

It remains to show that α is a strong morphism. Suppose that $s \cdot x'$ is defined. By assumption, there exists $a \in S$ such that $x' = a \cdot x$. By definition, $\alpha(x') = a \cdot y$. We have that $s \cdot x' = s \cdot (a \cdot x) = sa \cdot x$. By definition $\alpha(s \cdot x') = sa \cdot y$. But $sa \cdot y = s \cdot (a \cdot y) = s \cdot \alpha(x')$. Hence $\alpha(s \cdot x') = s \cdot \alpha(x')$.

Now suppose that $\alpha(x') = y'$ and $s \cdot y'$ exists. We shall prove that $s \cdot x'$ exists. Observe that $s^{-1}s \cdot y'$ exists and that it is enough to prove that $s^{-1}s \cdot x'$ exists. Let $x' = u \cdot x$, which exists since we are assuming that our action is transitive. Then, by what we proved above, $y' = u \cdot y$. Observe that $u^{-1}(s^{-1}s)u \cdot y = y$ and so $u^{-1}(s^{-1}s)u \in E(S_y)$. It follows by our assumption that $u^{-1}(s^{-1}s)u \in E(S_x)$ and so $u^{-1}(s^{-1}s)u \cdot x = x$. It readily follows that $s^{-1}s \cdot x'$ exists, and so $s \cdot x'$ exists, as required.

We now prove the converse. Let $\alpha: X \rightarrow Y$ be a strong morphism such that $\alpha(x) = y$. Let $s \in S_x$. Then $s \cdot x$ exists and $s \cdot x = x$. By the definition of strong morphism, it follows that $s \cdot \alpha(x)$ exists and that $\alpha(s \cdot x) = s \cdot \alpha(x)$. But $s \cdot x = x$ and so $\alpha(x) = s \cdot \alpha(x)$. Hence $s \cdot y = y$. We have therefore proved that $s \in S_y$, and so $S_x \subseteq S_y$. Let $e \in E(S_y)$. Then $e \cdot \alpha(x)$ exists. But α is a strong morphism and so $e \cdot x$ exists. Clearly $e \in E(S_x)$. It follows that $E(S_x) = E(S_y)$. □

The following result is adapted from Lemma 1.9 of Ruyle [22] and will be useful to us later.

LEMMA 2.9. *Let F be a closed inverse subsemigroup of the semilattice of idempotents of the inverse subsemigroup S . Define*

$$\overline{F} = \{s \in S : s^{-1}Fs \subseteq F, sFs^{-1} \subseteq F\}.$$

Then \overline{F} is a closed inverse subsemigroup of S whose semilattice of idempotents is F . Furthermore, if T is any closed subsemigroup of S with semilattice of idempotents F then $T \subseteq \overline{F}$.

PROOF. Clearly the set \overline{F} is closed under inverses. Let $s, t \in \overline{F}$. We calculate

$$(st)^{-1}F(st) = t^{-1}(s^{-1}Fs)t \subseteq t^{-1}Ft \subseteq F$$

and

$$(st)F(st)^{-1} = s(tFt^{-1})s^{-1} \subseteq sFs^{-1} \subseteq F.$$

Thus $st \in \overline{F}$. It follows that \overline{F} is an inverse subsemigroup of S .

Let $e \in \overline{F}$ and $f \in F$. Then by assumption $ef \in F$. But $ef \leq e$ and F is a closed inverse subsemigroup of the semilattice of idempotents and so $e \in F$. Thus $E(\overline{F}) = F$.

Let $s \leq t$ where $s \in \overline{F}$. Then $s = ss^{-1}t = ft$. Let $e \in F$. Then

$$s^{-1}es = t^{-1}feft = t^{-1}eft \leq t^{-1}et.$$

Now $s^{-1}es, t^{-1}et$ are idempotents and $s^{-1}es \in F$ thus $t^{-1}et \in F$, because F is a closed inverse subsemigroup of the semilattice of idempotents. Similarly, $tet^{-1} \in F$. It follows that $t \in \overline{F}$ and so \overline{F} is a closed inverse subsemigroup of S .

Finally, let T be a closed inverse subsemigroup of S such that $E(T) = F$. Let $t \in T$. Then for each $e \in F$ we have that $t^{-1}et, tet^{-1} \in F$. Thus $T \subseteq \overline{F}$. □

A closed inverse subsemigroup T of S will be said to be *fully closed* if $T = \overline{E(T)}$. Closed inverse subsemigroups of the semilattice of idempotents of an inverse semigroup are called *filters in $E(S)$* . Observe the emphasis on the word ‘in’. A filter in $E(S)$ is said to be *principal* if it is of the form e^\uparrow . We denote by $\mathcal{F}_{E(S)}$ the set of all closed inverse subsemigroups of $E(S)$ and call it the *filter space of the semilattice of idempotents of S* . This filter space is a poset when we define $F \leq F'$ if and only if $F' \subseteq F$ so that, in particular, $e^\uparrow \leq f^\uparrow$ if and only if $e \leq f$.

Let F be a filter in $E(S)$. Then F^\uparrow is a closed inverse subsemigroup containing F and clearly the smallest such inverse subsemigroup. On the other hand, by Lemma 2.9, \overline{F} is the largest closed inverse subsemigroup with semilattice of idempotents F . We have therefore proved the following lemma.

LEMMA 2.10. *The semilattice of idempotents of any closed inverse subsemigroup H of an inverse semigroup S is a filter F in $E(S)$ and $F^\uparrow \subseteq H \subseteq \overline{F}$. Thus F^\uparrow is the smallest closed inverse subsemigroup with semilattice of idempotents F and \overline{F} is the largest.*

PROPOSITION 2.11. *Let S be an inverse semigroup and let $G = S/\sigma$. Then there is an inclusion-preserving bijection between the wide closed inverse subsemigroups of S and the subgroups of G .*

PROOF. Let T be a wide inverse subsemigroup of S . Then the image of T in G is a subgroup since inverse subsemigroups map to inverse subsemigroups under homomorphisms. Suppose that T and T' , where also $E(S) \subseteq T' \subseteq S$, have the same image in G . Let $t \in T$. Then $\sigma^{\mathfrak{h}}(t) = \sigma^{\mathfrak{h}}(t')$ for some $t' \in T'$. Thus $a \leq t, t'$ from the definition of σ . But both T and T' are order ideals of S and so $a \in T \cap T'$. Thus $a \leq t$ and $a \in T'$ and T' is closed thus $t \in T'$. We have shown that $T \subseteq T'$. The reverse inclusion follows by symmetry. If H is a subgroup of G then the full inverse image

of H under σ^{\sharp} is a wide inverse subsemigroup of S . This defines an order-preserving map going in the opposite direction. It is now clear that the result holds. \square

The following is a special case of [22, Lemma 2.17]. We include it for interest since we shall not use it explicitly.

LEMMA 2.12. *Let F be a filter in $E(S)$ in the inverse semigroup S .*

- (1) *The intersection of any family of closed inverse subsemigroups with common semilattice of idempotents F is again a closed inverse subsemigroup with semilattice of idempotents F .*
- (2) *Given any family of closed inverse subsemigroups with common semilattice of idempotents F there is a smallest closed inverse subsemigroup with semilattice F which contains them all.*

2.2. Universal and fundamental transitive actions. We shall now define two special classes of transitive actions that play a decisive role in this paper. Let S be an inverse semigroup and let H be a closed inverse subsemigroup of S . By Lemma 2.10,

$$E(H)^{\uparrow} \subseteq H \subseteq \overline{E(H)}$$

where $E(H)$ is a filter in $E(S)$. We shall use this observation as the basis of two definitions, the first of which is by far the most important. We shall say that a transitive S -space X is *universal* if the stabilizer of a point of X is the closure F^{\uparrow} for some filter F of $E(S)$, and *fundamental* if the stabilizer of a point of X is \overline{F} for some filter F in $E(S)$. Both definitions are independent of the point chosen.

LEMMA 2.13. *With the above definitions, the following hold.*

- (1) *A strong morphism between universal transitive actions is an equivalence.*
- (2) *Any strong morphism with domain a fundamental transitive action and codomain a transitive action is an equivalence.*

PROOF. (1) Let X and Y be universal transitive spaces. Let $\alpha: X \rightarrow Y$ be a strong morphism. Choose $x \in X$. Then $S_x \subseteq S_{\alpha(x)}$ and $E(S_x) = E(S_{\alpha(x)})$. But the actions are universal and so all stabilizers are the full closures of their semilattices of idempotents. Thus $S_x = S_{\alpha(x)}$ and so α is an equivalence by the second part of Theorem 2.8.

(2) Let X and Y be transitive spaces where X is fundamental and let $\alpha: X \rightarrow Y$ be a strong morphism. Choose $x \in X$ and let $y = \alpha(x)$. Then $S_x \subseteq S_y$ and $E(S_x) = E(S_y)$ by the second part of Theorem 2.8. But S_x is fundamental and so $S_x = S_y$. We may deduce from the second part of Theorem 2.8 that there is a unique strong morphism from Y to X mapping y to x . It follows that α is an equivalence. \square

If $\alpha: X \rightarrow Y$ is a strong morphism between two transitive S -spaces, we shall say that Y is *strongly covered* by X . The importance of universal actions arises from the following result.

PROPOSITION 2.14. *Let S be an inverse semigroup.*

- (1) *Each transitive action of S is strongly covered by a universal one.*
- (2) *Each transitive action of S strongly covers a fundamental one.*

PROOF. (1) Let Y be an arbitrary transitive S -space. Choose a point $y \in Y$. Let $F = E(S_y)$ and put $H = F^\uparrow$. Then $E(H) = E(S_y)$ and $H \subseteq S_y$. Put $X = S/H$ and choose the point x in X to be the coset H . Then there is a unique strong morphism $\alpha: X \rightarrow Y$ such that $\alpha(x) = y$ by the second part of Theorem 2.8 which is surjective by the second part of Lemma 2.2 and X is a universal transitive space by construction.

(2) Let Y be an arbitrary transitive S -space. Choose a point $y \in Y$. Let $F = E(S_y)$ and put $H = \overline{F}$. Thus, by Lemma 2.10, $S_y \subseteq H$ and $E(S_y) = E(H)$. Put $X = S/H$ and choose the point x in X to be the coset H . Then there is a unique strong morphism $\alpha: Y \rightarrow X$ such that $\alpha(y) = x$ by the second part of Theorem 2.8 which is surjective by the second part of Lemma 2.2 and X is a fundamental transitive space by construction. □

THEOREM 2.15. *Let X be a universal, transitive S -space and let x be a point of X . Put $S_x = F^\uparrow$, where F is a filter in $E(S)$ and $G_F = \overline{F}/\sigma$. Then there is an order-preserving bijection between the set of strong congruences on X and the set of subgroups of G_F .*

PROOF. Put $G = G_F$. By Proposition 2.11, there is an order-preserving bijection between the closed inverse subsemigroups H such that $F^\uparrow \subseteq H \subseteq \overline{F}$ and the subgroups of G . Thus we need to show that there is a bijection between the set of strong congruences on X and the set of closed wide inverse subsemigroups of \overline{F} . Observe that we use the fact that strong morphisms between transitive spaces are surjective by the second part of Lemma 2.2.

Let \sim be a strong congruence defined on X . Then by Proposition 2.3 it determines a strong morphism $\nu: X \rightarrow X/\sim$. For x given in the statement of the theorem, we have that the stabilizer of $[x]$, the \sim -class containing x , is a closed inverse subsemigroup H_x such that $F^\uparrow \subseteq H_x \subseteq \overline{F}$ by the second part of Theorem 2.8. We have thus defined a function from strong congruences on X to the set of closed wide inverse subsemigroups of \overline{F} .

Suppose that \sim_1 and \sim_2 are two strong congruences on X that map to the same closed wide inverse subsemigroup. Denote the \sim_i equivalence class containing x by $[x]_i$ and let $\nu_i: X \rightarrow X/\sim_i$ be the natural map. Let $x \in X$. Then the stabilizer of $[x]_1$ and the stabilizer of $[x]_2$ are the same, namely H . Suppose that $x \sim_1 y$. Thus $[x]_1 = [y]_1$. Since X is an universal transitive S -space there exists $b \in B$ such that $b \cdot x = y$. It follows that $b \cdot [x]_1 = [y]_1 = [x]_1$ and so $b \in H$. By assumption, $b \cdot [x]_2 = [x]_2$. But \sim_2 is a strong congruence and so $y = b \cdot x \sim_2 x$ and so $x \sim_2 y$. A symmetrical argument shows that \sim_1 and \sim_2 are equal. Thus the correspondence we have defined is injective. We now show that it is surjective.

Let $F^\uparrow \subseteq H \subseteq \overline{F}$ be such a closed wide inverse subsemigroup. Then $Y = H/S$ is a transitive S -space. Choose the point $y = H \in Y$. Then by the second part of Theorem 2.8 there is a unique strong morphism $\alpha_H: X \rightarrow Y$ such that $\alpha(x) = y$.

The kernel of α_H , which we denote by \sim_H , is a strong congruence defined on X by Proposition 2.3, and the kernel of α_H maps to H . \square

Observe that the above theorem requires a chosen point in X .

2.3. A topological interpretation. Let S be an inverse semigroup and X an S -space. Define an S -labeled graph $G(X)$ whose vertices are X and whose edges go from x to sx , where $x \in X$, $s \in S$ and sx is defined, with label s on this edge in this case. There is an obvious involution on the graph by inversion, so this is a graph *in the sense of Serre*. Observe that the directed graph $G(X)$ is connected if and only if X is transitive. From now on we shall deal only with transitive actions and so our graphs will be connected.

The *star* of a vertex x in $G(X)$ is the set of all edges that start at x . Now let G and H be arbitrary graphs. A morphism f from G to H is called an *immersion* if it induces an injection from the star set of x to that of $f(x)$ for each vertex x of G . The morphism f is called a *cover* if it induces a bijection between such star sets. The following is the key link between the algebraic and the topological interpretations of inverse semigroup actions.

LEMMA 2.16. *Let S be an inverse semigroup and let X and Y be transitive S -spaces. There is a morphism from X to Y if and only if there is a label-preserving immersion from $G(X)$ to $G(Y)$, and there is a strong morphism from X to Y if and only if there is a label-preserving cover from $G(X)$ to $G(Y)$.*

PROOF. Let $\alpha: X \rightarrow Y$ be a morphism of transitive S -spaces. Consider the directed edge $x \xrightarrow{s} y$ in the graph $G(X)$. Then $s \cdot x = y$. Since α is a morphism, $\alpha(s \cdot x) = s \cdot \alpha(x) = \alpha(y)$. We may therefore define $f: G(X) \rightarrow G(Y)$ by mapping the edge $x \xrightarrow{s} y$ to the edge $\alpha(x) \xrightarrow{s} \alpha(y)$. It is immediate that this is an immersion. The fact that immersions arise from morphisms is now straightforward to prove. Finally, suppose that α is a strong morphism. Let $\alpha(x) \xrightarrow{s} \alpha(y)$ be an edge. This means that $s \cdot \alpha(x) = \alpha(y)$. But α is a strong morphism and so $s \cdot x$ exists and $\alpha(s \cdot x) = s \cdot \alpha(x)$. It follows that the graph map is a cover. \square

For a more complete account of the connection between immersions, inverse monoids and inverse categories, see [16, 27].

3. The étale groupoid associated with an inverse semigroup

In Section 2, we investigated the relationship between transitive actions of an inverse semigroup and closed inverse subsemigroups. We found that the universal transitive actions played a special role. We shall show in this section how these universal transitive actions, via their stabilizers, lead to the inverse semigroup introduced by Lenz and thence to Paterson's étale groupoid.

3.1. The inverse semigroup of cosets $\mathcal{K}(S)$. We begin by reviewing a construction studied by a number of authors [8, 9, 14, 26]. A subset $A \subseteq S$ of an inverse semigroup is called an *atlas* if $A = AA^{-1}A$. A closed atlas is precisely a coset of a closed inverse

subsemigroup of S [8]. We shall therefore refer to a closed atlas as a *coset*. Observe that the intersection of cosets, if nonempty, is a coset. The set of cosets of S is denoted by $\mathcal{K}(S)$. There is a product on $\mathcal{K}(S)$, denoted by \otimes , and defined as follows: if $A, B \in \mathcal{K}(S)$ then $A \otimes B$ is the intersection of all cosets of S that contain the set AB . More explicitly if $X = (aH)^\uparrow$, where $a^{-1}a \in H$, and $Y = (bK)^\uparrow$, where $b^{-1}b \in K$, then $X \otimes Y = (ab\langle b^{-1}Hb, K \rangle)^\uparrow$ where $\langle C, D \rangle$ is the inverse subsemigroup of S generated by $C \cup D$. In fact, $\mathcal{K}(S)$ is an inverse semigroup called the *(full) coset semigroup of S* . Note that its natural partial order is reverse inclusion. Thus S is the zero element of $\mathcal{K}(S)$. The idempotents of $\mathcal{K}(S)$ are just the closed inverse subsemigroups of S .

There is an embedding $\iota: S \rightarrow \mathcal{K}(S)$ that maps s to s^\uparrow . Observe now that if $A \in \mathcal{K}(S)$ then for each $s \in A$ we have that $s^\uparrow \subseteq A$ and so $A \leq s^\uparrow$. It follows readily from this that A is in fact the meet of the set $\{s^\uparrow: s \in A\}$. More generally, every nonempty subset of $\mathcal{K}(S)$ has a meet and so the inverse semigroup $\mathcal{K}(S)$ is *meet complete*. The map $\iota: S \rightarrow \mathcal{K}(S)$ is universal for maps to meet complete inverse semigroups. Thus the inverse semigroup $\mathcal{K}(S)$ is the *meet completion* of the inverse semigroup S [14]. It is worth noting that the category of meet complete inverse semigroups and their morphisms is not a full subcategory of the category of inverse semigroups and their homomorphisms and so the meet completion of $\mathcal{K}(S)$ is $\mathcal{K}(\mathcal{K}(S))$ and not just $\mathcal{K}(S)$.

At this point, we want to highlight a class of transitive actions that will play an important role both here and in Section 4. Let T be an inverse semigroup and let e be any idempotent in T . We denote by L_e the \mathcal{L} -class containing e . The set L_e therefore consists of all elements $t \in T$ such that $\mathbf{d}(t) = e$. Define a partial function from $T \times L_e$ to L_e by $a \cdot x$ exists if and only if $\mathbf{d}(ax) = e$. This defines a transitive action of T on L_e called the *(left) Schützenberger action determined by the idempotent e* . This is the transitive action determined by the closed inverse subsemigroup e^\uparrow .

The structure of $\mathcal{K}(S)$ is inextricably linked to the structure of transitive actions of S . The following proposition was first stated in [8].

PROPOSITION 3.1. *Let S be an inverse semigroup. Every transitive representation of S is the restriction of a Schützenberger representation of $\mathcal{K}(S)$.*

PROOF. Let H be a closed inverse subsemigroup of S . In the inverse semigroup $\mathcal{K}(S)$, the \mathcal{L} -class L_H of the idempotent H consists of all $A \in \mathcal{K}(S)$ such that $A^{-1} \otimes A = H$. Let $a \in A$. Then $A = (aH)^\uparrow$. It follows that L_H consists of precisely the left cosets of H in S . Let $A \in L_H$ and consider the product $s^\uparrow \otimes A$. Then this again belongs to L_H precisely when $(sa)^{-1}sa \in H$ and is equal to $(saH)^\uparrow$. It follows that via the map ι the inverse semigroup acts on L_H precisely as it acts on S/H . □

If H and K are two idempotents of $\mathcal{K}(S)$ then they are \mathcal{D} -related if and only if there exists $A \in \mathcal{K}(S)$ such that $A^{-1} \otimes A = H$ and $A \otimes A^{-1} = K$ if and only if H and K are conjugate. Thus the \mathcal{D} -classes of $\mathcal{K}(S)$ are in bijective correspondence with the conjugacy classes of closed inverse subsemigroups.

We may, in some sense, ‘globalize’ the connection between $\mathcal{K}(S)$ and transitive actions of S . Denote by $\mathbf{O}(S)$ the category whose objects are the *right S -spaces*

H/S and whose arrows are the (right) morphisms. We now recall the following construction [10]. Let S be an inverse semigroup. We can construct from S a right cancellative category, denoted $\mathbf{R}(S)$, whose elements are pairs $(s, e) \in S \times E(S)$ such that $\mathbf{d}(s) \leq e$. We regard (s, e) as an arrow from e to $\mathbf{r}(s)$ and define a product by $(s, e)(t, f) = (st, e)$.

The following proposition generalizes [10, Example 2.2.3].

PROPOSITION 3.2. *The category $\mathbf{O}(S)$ is isomorphic to the category $\mathbf{R}(\mathcal{K}(S))$.*

PROOF. We observe first that a morphism with a transitive space as its domain is determined by its value on any element of that domain. Let $\phi: U/S \rightarrow V/S$ be a morphism. Then ϕ is determined by the value taken by $\phi(U) = (Va)^\uparrow$. Now the stabilizer S_U of U is U itself and the stabilizer $S_{(Va)^\uparrow}$ is $(a^{-1}Va)^\uparrow$. Thus by Theorem 2.8, $U \subseteq (a^{-1}Va)^\uparrow$. Conversely, if we are given that $U \subseteq (a^{-1}Va)^\uparrow$ then we can define a morphism from U/S to V/S by $U \mapsto (Va)^\uparrow$. There is therefore a bijection between morphisms from U/S to V/S and inclusions $U \subseteq (a^{-1}Va)^\uparrow$. We shall encode the morphism ϕ by the triple $(V, (Va)^\uparrow, U)$. Let $\psi: V/S \rightarrow W/S$ be a morphism encoded by the triple $(W, (Wb)^\uparrow, V)$. The triple encoding $\psi\phi$ is of the form $(W, (Wc)^\uparrow, U)$ where $\psi\phi(U) = (Wc)^\uparrow$. Thus $(W, (Wb)^\uparrow, V)(V, (Va)^\uparrow, U) = (W, (Wba)^\uparrow, U)$. The product $(Wb)^\uparrow \otimes (Va)^\uparrow$ in $\mathcal{K}(S)$ is precisely $(Wba)^\uparrow$. We now recall that the natural partial order in $\mathcal{K}(S)$ is reverse inclusion. It follows that the triple $(V, (Va)^\uparrow, U)$ can be identified with the pair $((Va)^\uparrow, U)$ where $\mathbf{d}((Va)^\uparrow) \leq U$. We regard $((Va)^\uparrow, U)$ as an arrow with domain U and codomain V . The result now follows. \square

3.2. The inverse semigroup of filters $\mathcal{L}(S)$. We shall now describe an inverse subsemigroup of $\mathcal{K}(S)$. A subset $A \subseteq S$ of an inverse semigroup S is said to be (*down*) *directed* if it is nonempty and, for each $a, b \in A$, there exists $c \in A$ such that $c \leq a, b$. Closed directed sets in a poset are called *filters*. When this definition is applied to semilattices we recover the definition given earlier.

LEMMA 3.3. *The closed directed subsets are precisely the directed cosets.*

PROOF. A directed coset is certainly a closed directed subset. Let A be a closed directed subset. We prove that it is an atlas. Clearly $A \subseteq AA^{-1}A$. Thus we need only check that $AA^{-1}A \subseteq A$. Let $a, b, c \in A$. Then since A is directed there is $d \in A$ such that $d \leq a, b, c$. Thus $d = dd^{-1}d \leq ab^{-1}c$ and so $ab^{-1}c \in A$ since A is also closed. \square

LEMMA 3.4. *A closed inverse subsemigroup T of an inverse semigroup S is directed if and only if there is a filter $F \subseteq E(S)$ such that $T = F^\uparrow$.*

PROOF. Suppose that $T = F^\uparrow$. Let $a, b \in T$. Then $e \leq a$ and $f \leq b$ for some $e, f \in F$. But F is a filter in the semilattice of idempotents and so closed under multiplication. Thus $ef \in F$. But then $ef \leq a, b$ and so T is directed.

Let T be a closed directed inverse subsemigroup. Put $F = E(S)$. Let $e, f \in F$. Now T is directed and so there is $i \in T$ such that $i \leq e, f$. Thus i is an idempotent. But $i \leq ef \leq e, f$ and so, since F is closed, we have that $ef \in F$. It follows that F is a filter

in $E(S)$. Clearly $F^\uparrow \subseteq T$. Let $t \in T$. Then $t^{-1}t \in T$ since T is an inverse subsemigroup. But T is directed so there exists $j \leq t, t^{-1}t$. But then j is an idempotent and so $j \leq t$ gives that $t \in F^\uparrow$. Hence $T \subseteq F^\uparrow$. Thus $T = F^\uparrow$, as required. \square

LEMMA 3.5. *If A and B are both directed cosets then $(AB)^\uparrow$ is the smallest directed coset containing AB ; it is also the smallest coset containing AB .*

PROOF. The set $(AB)^\uparrow$ is closed so we need only show that it is directed. Let $ab, a'b' \in AB$. Then there exist $c \leq a, a'$ where $c \in A$ and $d \leq b, b'$ where $d \in B$. It follows that $cd \in AB$ and $cd \leq ab, a'b'$. Thus the set is directed.

Now let X be any coset containing AB . Then X is closed and so $(AB)^\uparrow \subseteq X$. \square

The subset of $\mathcal{K}(S)$ consisting of directed cosets is denoted by $\mathcal{L}(S)$.

PROPOSITION 3.6. *Let S be an inverse semigroup.*

- (1) $\mathcal{L}(S)$ is an inverse subsemigroup of $\mathcal{K}(S)$.
- (2) The directed cosets of S are precisely the cosets of the closed directed inverse subsemigroups of S .
- (3) Each element of $\mathcal{K}(S)$ is the meet of a subset of $\mathcal{L}(S)$ contained in an \mathcal{H} -class of $\mathcal{L}(S)$.

PROOF. (1) If $A, B \in \mathcal{K}(S)$ then their product is the intersection of all cosets containing AB . But if $A, B \in \mathcal{L}(S)$ then by Lemma 3.5 this intersection will also belong to $\mathcal{L}(S)$. Closure under inverses is immediate. Thus $\mathcal{L}(S)$ is an inverse subsemigroup of $\mathcal{K}(S)$.

(2) If $A \in \mathcal{K}(S)$ then $A = (aH)^\uparrow = (a)^\uparrow \otimes H$ where $H = A^{-1} \otimes A$ and $a \in A$. Thus A is directed if and only if H is directed.

(3) Let $A \in \mathcal{K}(S)$ be a coset. Define a relation \sim on the set A by $a \sim b$ if and only if there exists $c \in A$ such that $c \leq a, b$. We show that \sim is an equivalence relation on A . Clearly \sim is reflexive and symmetric. It only remains to prove that it is transitive. Let $a \sim b$ and $b \sim c$. Then there exist $x \leq a, b$ and $y \leq b, c$ where $x, y \in A$. In particular, $x, y \leq b$. Thus $z = xy^{-1}y = yx^{-1}x$ is the meet of x and y . Since A is a coset $xy^{-1}y, yx^{-1}x \in A$. It follows that $z \leq a, c$. Denote the blocks of the partition induced by \sim on A by A_i where $i \in I$. Each block is directed by construction and easily seen to be closed. It follows that each block is a directed coset and so $A_i \in \mathcal{L}(S)$. We have therefore proved that $A = \bigwedge_{i \in I} A_i$.

It remains to show that $A_i \mathcal{H} A_j$. To do this it is enough to compute $A_i^{-1} \otimes A_j$ and $A_i \otimes A_j^{-1}$ and observe that these idempotents do not depend on the suffix i . We may write $A = (aH)^\uparrow$ for some closed inverse subsemigroup H of S and element a such that $\mathbf{d}(a) \in H$. Put $F = E(H)$ the semilattice of idempotents of H . Put $K = F^\uparrow$ and $L = (aKa^{-1})^\uparrow$, both closed directed inverse subsemigroups of S and so elements of $\mathcal{L}(S)$. We prove that $K = A_i^{-1} \otimes A_i$ and $L = A_i \otimes A_i^{-1}$. From $A \leq A_i$ we have that $H = A^{-1} \otimes A \leq A_i^{-1} \otimes A_i$ and $(aHa^{-1})^\uparrow \leq A_i \otimes A_i^{-1}$. By construction, $H \leq K$ and K is in fact the smallest idempotent of $\mathcal{L}(S)$ above H . It follows that $K \leq A_i^{-1} \otimes A_i$ and similarly $L \leq A_i \otimes A_i^{-1}$. It remains to show that equality holds in each case, which means checking that $K \subseteq A_i^{-1} \otimes A_i$ and $L \subseteq A_i \otimes A_i^{-1}$.

Let $k \in K$ and $a_i \in A_i$. Now $k \in K \subseteq H$ and $a_i \in A_i \subseteq A$. Thus $a_i k \in A$. But $a_i k \leq a_i$. Now if $a_i k \in A_j$ then by closure $a_i \in A_j$ and so we must have that $a_i k \in A_i$. Thus $ka_i^{-1}a_i \in A_i^{-1} \otimes A_i$ and so by closure $k \in A_i^{-1} \otimes A_i$, as required.

Let $l \in L$. Let $a_i \in A_i$. Then $A = (a_i H)^\uparrow$. Thus $L = (a_i K a_i^{-1})^\uparrow$. It follows that $a_i^{-1} l a_i \in K$ and so $a_i a_i^{-1} l \in a_i K a_i^{-1}$ giving $l \in A_i \otimes A_i^{-1}$.

An alternative way of proving this result is to observe that K is a closed inverse subsemigroup of H and so H can be written as a disjoint union of some of the left cosets of K . We can then use this decomposition to write A itself as a disjoint union of left cosets of K . □

We say that an inverse semigroup S is *meet complete* if every nonempty subset of S has a meet. Meet completions of inverse semigroups are discussed at the end of [8, 9, Section 1.4] and most importantly in [14]. The meet completion of an inverse semigroup S is in fact $\mathcal{K}(S)$ [14].

The inverse semigroup S is said to have all *directed meets* if it has meets of all nonempty directed subsets. The result below shows that $\mathcal{L}(S)$ is the *directed meet completion* of S in the same way that $\mathcal{K}(S)$ is the meet completion.

PROPOSITION 3.7. *Let S be an inverse semigroup. Then $\mathcal{L}(S)$ is the directed meet completion of S .*

PROOF. We have the embedding $\iota: S \rightarrow \mathcal{L}(S)$ and once again each $A \in \mathcal{L}(S)$ is the join of all the s^\uparrow where $s \in A$. This time the set over which we are calculating the meet is directed. Let $\mathcal{A} = \{A_i : i \in I\}$ be a directed subset of $\mathcal{K}(S)$. Thus for each pair of cosets A_i and A_j there is a coset A_k such that $A_k \leq A_i, A_j$. Put $A = \bigcup_{i \in I} A_i$. It is clearly a closed subset. If $a, b \in A$ then $a \in A_i$ and $b \in A_j$ for some i and j . By assumption $A_i, A_j \subseteq A_k$ for some k . Thus $a, b \in A_k$. But A_k is a directed subset and so there exists $c \in A_k$ such that $c \leq a, b$. It follows that A is a closed and directed subset and so is a directed coset by Lemma 3.3. It is now immediate that A is the meet of the set \mathcal{A} . Let $\theta: S \rightarrow T$ be a homomorphism to an inverse semigroup T which has all meets of directed subsets. Define $\psi: \mathcal{K}(S) \rightarrow T$ by $\psi(A) = \bigwedge \theta(A)$. Then ψ is a homomorphism and the unique one such that $\psi \iota = \theta$. □

In [15], Lenz constructs an inverse semigroup $\mathcal{O}(S)$ from an inverse semigroup S , which is the basis for his étale groupoid associated with S . The key result for our paper is the following theorem.

THEOREM 3.8. *The inverse semigroup $\mathcal{L}(S)$ is isomorphic to Lenz’s semigroup $\mathcal{O}(S)$.*

PROOF. Let $\mathcal{F} = \mathcal{F}(S)$ denote the set of directed subsets of S . For $A, B \in \mathcal{F}$ define $A < B$ if and only if for each $b \in B$ there exists $a \in A$ such that $a \leq b$. This is a preorder. The associated equivalence relation is given by $A \sim B$ if and only if $A < B$ and $B < A$. We now make two key observations. (1) $A \sim A^\uparrow$. It is easy to check that A^\uparrow is directed. By definition $A < A^\uparrow$, whereas $A^\uparrow < A$ is immediate. (2) $A^\uparrow \sim B^\uparrow$ if and only if $A^\uparrow = B^\uparrow$. There is only one direction that needs proving. Suppose that $A^\uparrow \sim B^\uparrow$. Let $a \in A^\uparrow$. Then $B^\uparrow < A^\uparrow$ and so there is $b \in B$ such that $b \leq a$. But then $a \in B^\uparrow$. Thus $A^\uparrow \subseteq B^\uparrow$.

The reverse inclusion is proved similarly. By (1) and (2), it follows that $A \sim B$ if and only if $A^\uparrow = B^\uparrow$. As a set, $\mathcal{O}(S) = \mathcal{F}(S)/\sim$. We have therefore set up a bijection between $\mathcal{O}(S)$ and $\mathcal{L}(S)$. Lemma 3.5 tells us that the multiplication defined in [15] in $\mathcal{O}(S)$ ensures that this bijection is an isomorphism. \square

Denote by $\mathbf{U}(S)$ the category whose objects are the *right* S -spaces H/S where H is directed and whose arrows are the (right) morphisms. We have the following analog of Proposition 3.2.

PROPOSITION 3.9. *The category $\mathbf{U}(S)$ is isomorphic to the category $\mathbf{R}(\mathcal{L}(S))$.*

3.3. Paterson's étale groupoid. Theorem 3.8 brings us to the beginning of Section 4 of Lenz's paper [15] where he describes Paterson's étale groupoid. If T is an inverse semigroup, then it becomes a groupoid when we define a partial binary operation \cdot , called the *restricted product*, by $s \cdot t$ exists if and only if $\mathbf{d}(s) = \mathbf{r}(t)$ in which case $s \cdot t = st$. Paterson's groupoid is precisely $(\mathcal{L}(S), \cdot)$ equipped with a suitable topology. The isomorphism functor defined by Lenz from $\mathcal{L}(S)$ to Paterson's groupoid can be very easily described in terms of the ideas introduced in our paper. Let $A \in \mathcal{L}(S)$. Define $P = (AA^{-1})^\uparrow$. Then for any $a \in A$ we have that $A = (Pa)^\uparrow$. Thus we may regard A as a *right coset* of the closed, directed inverse subsemigroup P . By the dual of Lemma 2.4(1), we have that $(Pa)^\uparrow = (Pb)^\uparrow$, where $aa^{-1}, bb^{-1} \in P$, if and only if $ab^{-1} \in P$ if and only if $pa = pb$ for some $p \in P$, where we use the fact that every element of P is above an idempotent. The ordered pair (P, a) where $\mathbf{r}(a) \in P$ determines the right coset $(Pa)^\uparrow$, and another such pair (P, b) determines the same right coset if and only if $pa = pb$ for some $p \in P$. This leads to an equivalence relation and we denote the equivalence class containing (P, a) by $[P, a]$. The isomorphism functor between the Lenz groupoid $\mathcal{L}(S)$ and Paterson's groupoid is therefore defined by $A \mapsto [(AA^{-1})^\uparrow, a]$ where $a \in A$. We see that Paterson has to work with equivalence classes because of the nonuniqueness of coset-representatives, and Lenz has to work with equivalence classes because he works with generating sets of filters rather than with the filters themselves. In our approach, the use of equivalence classes in both cases is avoided.

Recall from Section 2.2, that a transitive S -space X is universal if the stabilizer H of a point of X is F^\uparrow where F is a filter in $E(S)$. In other words, by Lemma 3.4 the closed inverse subsemigroup H is directed. It follows that the universal transitive actions of S are determined by the directed filters that are also inverse subsemigroups. We shall now describe how the structure of the groupoid $(\mathcal{L}(S), \cdot)$ reflects the properties of transitive actions of S . In what follows, we can just as easily work in the inverse semigroup as in the groupoid.

PROPOSITION 3.10. *Let S be an inverse semigroup.*

- (1) *The connected components of the groupoid $\mathcal{L}(S)$ are in bijective correspondence with the equivalence classes of universal transitive actions of S .*
- (2) *Let H be an identity in $\mathcal{L}(S)$. Then the local group G_H at H is isomorphic to the group $\overline{E(H)}/\sigma$.*

PROOF. (1) The identities of $\mathcal{L}(S)$ are the closed directed inverse subsemigroups of S . Two such identities belong to the same connected component if and only if they are conjugate. The result now follows by Proposition 2.6.

(2) Put $F = E(H)$ so that $H = \overline{F}$. Let A be in the local group determined by H . Then $H = (A^{-1}A)^\uparrow = (AA^{-1})^\uparrow$. Define $\theta: G_H \rightarrow \overline{E(H)}/\sigma$ by $\theta(A) = \sigma(a)$ where $a \in A$.

We show first that this map is well defined. Let $f \in F$ and let $a \in A$. Then $a^{-1}fa \in A^{-1}E(A)A \subseteq HA = A$ and so $a^{-1}fa \in F$ and $afa^{-1} \in AE(A)A^{-1} \subseteq AH = A$ and so $afa^{-1} \in F$. Thus $A \subseteq \overline{F}$. Next suppose that $a, b \in A$. Then there is an element $c \in A$ such that $c \leq a, b$. Thus $\sigma(a) = \sigma(b)$. It follows that θ is well defined.

We now show that θ defines a bijection. Suppose that $\theta(A) = \theta(B)$. Then $a\sigma b$ where $a \in A$ and $b \in B$. Thus there exists $c \in \overline{F}$ such that $c \leq a, b$. It follows that $c = ac^{-1}c = bc^{-1}c$ and so $a^{-1}ac^{-1}c \leq a^{-1}b$. But $a^{-1}ac^{-1}c \in F$ and so $A = B$. Thus θ is injective. Let $a \in \overline{F}$. Then $a^{-1}a, \in \overline{F}$ and so $a^{-1}a \in F$. Thus $A = (aH)^\uparrow$ is a well-defined coset and then $(A^{-1}A)^\uparrow = H = (AA^{-1})^\uparrow$. It follows that $A \in G_H$ and $\theta(A) = \sigma(a)$. Thus θ is surjective.

Finally, we show that θ defines a homomorphism. Let $A, B \in G_H$ and $a \in A$ and $b \in B$. By Lemma 3.5, $A \otimes B = (AB)^\uparrow$ and contains ab . Thus $\theta(A)\theta(B) = \sigma(a)\sigma(b) = \sigma(ab) = \theta(A \otimes B)$. □

We now have the following theorem.

THEOREM 3.11. *Let S be an inverse semigroup. Then $\mathcal{L}(S)$ explicitly encodes universal transitive actions of S via its Schützenberger actions, and implicitly encodes all transitive actions via its local groups.*

PROOF. An idempotent of $\mathcal{L}(S)$ is just an inverse subsemigroup H of S that is also a filter. Denote by \mathcal{L}_H the \mathcal{L} -class of H in the inverse semigroup $\mathcal{L}(S)$. The elements of \mathcal{L}_H are just the left cosets of H in S . The inverse semigroup $\mathcal{L}(S)$ acts on the set \mathcal{L}_H , a Schützenberger action, and so too does S via the map ι of Proposition 3.7. This latter action is equivalent to the action of S on S/H . We have therefore shown that $\mathcal{L}(S)$ encodes universal transitive actions of S via its Schützenberger actions.

By the first part of Proposition 2.14 each transitive action of S on a set Y is strongly covered by a universal one X . Let H be a stabilizer of this universal action of S on X . Then the strong covering is determined by a strong congruence which by Theorem 2.15 is determined by a subgroup of the \mathcal{H} -class in $\mathcal{L}(S)$ containing the idempotent H ; in other words, by a subgroup of the local group determined by the idempotent H . □

Finally, the topology on the groupoid $\mathcal{L}(S)$ is defined in terms of the embedding $S \rightarrow \mathcal{L}(S)$ as follows. Let $s \in S$. Define

$$U_s = \{A \in \mathcal{L}(S) : s \in A\}$$

and for $s_1, \dots, s_n \leq s$ define

$$U_{s; s_1, \dots, s_n} = U_s \cap U_{s_1}^c \cap \dots \cap U_{s_n}^c.$$

Then the sets $U_{s; s_1, \dots, s_n}$ form a basis for a topology.

4. Matrix representations of inverse semigroups

We deduce here results of the third author on the finite-dimensional irreducible representations of inverse semigroups [28]. There an approach based on groupoid algebras was used, whereas here we use results of Green [2, Ch. 6] and the universal property of $\mathcal{L}(S)$.

4.1. Green’s theorem and primitive idempotents. The following theorem summarizes the contents of [2, Ch. 6]. Let A be a ring. A module is assumed to be a left A -module unless otherwise stated. We also consider only *unitary* A -modules, that is, A -modules M such that $AM = M$ (where AM means the submodule generated by elements am with $a \in A$ and $m \in M$). If A has a unit, then this is the same as saying that the unit acts as the identity on M . In particular, a *simple* A -module is an A -module M such that $AM \neq 0$ and there are no nonzero proper submodules of M . If e is an idempotent of A and M is an A -module, then eM is an eAe -module. The functor $M \mapsto eM$ is called *restriction* and we sometimes denote it by $\text{Res}_e(M)$. It is well known and easy to check that $eM \cong \text{Hom}_A(Ae, M)$, where the latter has a left eAe -action induced by the right action of eAe on Ae . For an eAe -module N , define

$$\text{Ind}_e(N) = Ae \otimes_{eAe} N.$$

The usual hom-tensor adjunction implies that Ind_e is the left adjoint of Res_e . Moreover, $\text{Res}_e \text{Ind}_e$ is isomorphic to the identity functor on the category eAe -modules. Indeed, $eaen \otimes n \mapsto eaen$ is an isomorphism with inverse $n \mapsto e \otimes n$. These isomorphisms are natural in N .

THEOREM 4.1 (Green). *Let A be a ring and $e \in A$ an idempotent.*

(1) *If N is a simple eAe -module, then the induced module*

$$\text{Ind}_e(N) = Ae \otimes_{eAe} N$$

has a unique maximal submodule $R(N)$, which can be described as the largest submodule of $\text{Ind}_e(N)$ annihilated by e . Moreover, the simple module $\tilde{N} = \text{Ind}_e(N)/R(N)$ satisfies $\tilde{N} \cong e\tilde{N}$.

- (2) *If M is a simple A -module with $eM \neq 0$, then eM is a simple eAe -module and $M \cong e\overline{M}$.*

Let S be an inverse semigroup and suppose that e is a minimum idempotent of S . Then $eSe = G_e$, the maximal subgroup of S at e , and is also the maximal group image of S . Moreover, $Se = G_e = eS$ and the action of S on the left of Se factors through the maximal group image homomorphism. Let k be a commutative ring with unit. Then $ekSe \cong kG_e$ and so Green’s theorem shows that simple kS -modules M with $eM \neq 0$ are in bijection with simple kG_e -modules via induction and restriction. Moreover, since $kSe = kG_e$, we have that $\text{Ind}_e(N) = N$ with the action of S induced by the maximal group image homomorphism. Thus $\text{Ind}_e(N)$ already is a simple kS -module. Let us consider the analogous situation for primitive idempotents.

Let e be a primitive idempotent of an inverse semigroup with 0. Observe that in this case $eSe = G_e \cup \{0\}$ since $e, 0$ are the only idempotents of eSe and so if $s \neq 0$, then $ss^{-1} = e = s^{-1}s$. Thus if k_0S is the contracted semigroup algebra of S (meaning the quotient of kS by the ideal of scalar multiples of the zero of S), then $ek_0Se \cong kG_e$ and so, again by Green’s theorem, we have a bijection between simple k_0S -modules M with $eM \neq 0$ and kG_e -modules via induction. We aim to show now that if N is a simple kG_e -module, then $\text{Ind}_e(N)$ is already a simple k_0S -module. Let L_e be the \mathcal{L} -class of e . Then since e is primitive, it follows that $L_e = Se \setminus \{0\}$ and so $k_0Se = kL_e$ where S acts on the left of kL_e via linearly extending the left Schützenberger representation. The group G_e acts freely on the right of L_e with orbits the \mathcal{H} -classes contained in L_e . Thus $k_0Se = kL_e$ is free as a right $ek_0Se = kG_e$ -module. Let T be a transversal to the \mathcal{H} -classes of L_e and let N be a kG_e -module. Then as a k -module, $\text{Ind}_e(N) = \bigoplus_{t \in T} t \otimes_k N$. A fact we shall use is that any element of L_e is primitive and so if $t_1 \neq t_2 \in T$, then $t_1t_1^{-1} \neq t_2t_2^{-1}$ and hence $t_1t_1^{-1}t_2 = 0$.

LEMMA 4.2. *If N is a nonzero kG_e -module, then no nontrivial submodule of $\text{Ind}_e(N)$ is annihilated by e .*

PROOF. Let M be a nonzero submodule of $\text{Ind}_e(N)$. Notice that M is annihilated by e if and only if it is annihilated by the ideal generated by e . So let $m = \sum_{t \in T} t \otimes n_t$ (with only finitely many terms nonzero) be a nonzero element of M . Then there exists $t \in T$ with $n_t \neq 0$. By the observation just before the proof $tt^{-1}m = t \otimes n_t \neq 0$ and so tt^{-1} does not annihilate m . But $e = t^{-1}t$ generates the same ideal as tt^{-1} and so M is not annihilated by e . □

As a corollary, we obtain from Green’s Theorem 4.1 that if N is a simple kG_e -module, then $\text{Ind}_e(N)$ is a simple k_0S -module.

COROLLARY 4.3. *Let S be an inverse semigroup, $e \in E(S)$ a primitive idempotent and k a commutative ring with unit. If N is a simple kG_e -module, then $\text{Ind}_e(N)$ is a simple kS -module.*

If k is a field, then from $\text{Ind}_e(N) = \bigoplus_{t \in T} t \otimes_k N$, we see that $\text{Ind}_e(N)$ is finite-dimensional if and only if T is finite and N is finite-dimensional.

4.2. The main result. Suppose now that S is any inverse semigroup and $e \in E(S)$. Let $I_e = SeS \setminus J_e$ be the ideal of elements strictly \mathcal{J} -below e . If N is a kG_e -module, then let

$$\text{Ind}_e(N) = k_0[S/I_e]e \otimes_{kG_e} N = (kS/kI_e)e \otimes_{kG_e} N.$$

Equivalently, if L_e is the \mathcal{L} -class of e , then kL_e is a free right kG_e -module with basis the set of \mathcal{H} -classes of L_e and also it is a left kS -module by means of the action of S on the left of L_e by partial bijections via the Schützenberger representation. Then $\text{Ind}_e(N) = kL_e \otimes_{kG_e} N$. Suppose now that the \mathcal{D} -class of e contains only finitely many idempotents; in this case we say that e has *finite index* in S . Under the hypothesis that e has finite index it is well known that if $f \in E(S)$ with $f < e$, then $SfS \neq SeS$ and so $f \in I_e$. Thus e is primitive in S/I_e and so Corollary 4.3 shows that $\text{Ind}_e(N)$ is simple for any simple kG_e -module in this setting.

We are now ready to construct the finite-dimensional irreducible representations of an inverse semigroup over a field. This was first carried out by Munn [17], whereas the construction presented here first appeared in [28] where it was deduced as a special case of a result on étale groupoids. Our approach here uses the inverse semigroup $\mathcal{L}(S)$. Fix a field k . First we construct a collection of simple kS -modules.

PROPOSITION 4.4. *Let $e \in E(\mathcal{L}(S))$ have finite index and let N be a simple kG_e -module. Then $\text{Ind}_e(N)$ is a simple kS -module. Moreover, $\text{Ind}_e(N)$ is finite-dimensional if and only if N is.*

PROOF. The above discussion shows that $\text{Ind}_e(N)$ is simple as a $k\mathcal{L}(S)$ -module so we just need to show that any S -invariant subspace is $\mathcal{L}(S)$ -invariant. In fact, we show that each element of $\mathcal{L}(S)$ acts the same on $\text{Ind}_e(N)$ as some element of S . It will then follow that any S -invariant subspace is $\mathcal{L}(S)$ -invariant and so $\text{Ind}_e(N)$ is a simple kS -module.

Let T be a transversal for the orbits of G_e on L_e . Then T is finite since these orbits are in bijection with \mathcal{R} -classes of D_e , which in turn are in bijection with idempotents of D_e . Let $A \in \mathcal{L}(S)$ and write $A = \bigwedge_{d \in D} s_d$ with $s \in S$ and D a directed set. We claim that if $t \otimes n$ is an elementary tensor with $t \in T$, then there exists $d_t \in D$ depending only on t (and not n) such that $A(t \otimes n) = s_{d_t}(t \otimes n)$ for all $d \geq d_t$. By [9, Section 1.4, Proposition 19], $At = \bigwedge_{d \in D} (s_d t)$. Since the \mathcal{D} -class of e has only finitely many idempotents, it follows by [9, Theorem 3.2.16] that distinct elements of \mathcal{D} are not comparable in the natural partial order. Since the set $\{s_d t \mid d \in D\}$ is directed, either $s_d t \leq_{\mathcal{L}} e$ for all sufficiently large elements of D or $s_d t$ is an element ℓ of L_e independent of d . In the first case $At \leq_{\mathcal{L}} e$ and in the second case $At = \ell$. Thus in the first case, $A(t \otimes n) = 0 = s_d(t \otimes n)$ for d large enough, whereas in the second case $A(t \otimes n) = \ell \otimes n = s_d(t \otimes n)$ for all $d \in D$. We conclude that d_t exists.

Since T is finite, we can find $d_0 \in D$ with $d_0 \geq d_t$ for all $t \in R$. Then A and s_{d_0} agree on all elements of the form $t \otimes n$ with $t \in T$ and $n \in N$. But such elements span $\text{Ind}_e(N)$ and so we conclude that A and s_{d_0} agree on $\text{Ind}_e(N)$.

The final statement follows from the previous discussion. □

Note that application of the restriction functor and the fact that $\text{Res}_e \text{Ind}_e$ is isomorphic to the identity shows that $\text{Ind}_e(N) \cong \text{Ind}_e(M)$ implies that $N \cong M$. Also, if e, f are two finite-index idempotents of $\mathcal{L}(S)$ and $e \not\leq_{\mathcal{J}} f$, then f annihilates $\text{Ind}_e(N)$ for any kG_e -module and hence all elements of f , viewed as a filter, annihilate $\text{Ind}_e(N)$. On the other hand, no element of the filter e annihilates $\text{Ind}_e(N)$. It follows that if e, f are finite-index idempotents that are not \mathcal{D} -equivalent, then the modules of the form $\text{Ind}_e(N)$ and $\text{Ind}_f(M)$ are never isomorphic. Clearly, \mathcal{D} -equivalent idempotents give isomorphic collections of simple modules. Thus, for each \mathcal{D} -class with finitely many idempotents, we get a distinct set of simple kS -modules (up to isomorphism).

The following fact is well known and easy to prove.

PROPOSITION 4.5. *Let k be a field and V an n -dimensional k -vector space. Then any semilattice in $\text{End}_k(V)$ has size at most 2^n .*

PROOF. Any idempotent matrix is diagonalizable and so any semilattice of matrices is simultaneously diagonalizable. But the multiplicative monoid of k^n has 2^n idempotents. □

We can now complete the description of the finite-dimensional irreducible representations of an inverse semigroup. In the statement of the theorem below, it is worth recalling that $e = H$ is a finite-index, closed directed subsemigroup of S and G_e is the group $\overline{E(H)}/\sigma$ described in Theorem 2.15.

THEOREM 4.6. *Let k be a field and S an inverse semigroup. Then the finite-dimensional simple kS -modules are precisely those of the form $\text{Ind}_e(N)$ where e is a finite-index idempotent of $\mathcal{L}(S)$ and N is a finite-dimensional simple kG_e -module.*

PROOF. It remains to show that every simple kS -module M is of this form. Let $\theta: S \rightarrow \text{End}_k(V)$ be the corresponding irreducible representation. Then $T = \theta(S)$ is an inverse semigroup with finitely many idempotents and so trivially directed meet complete. Thus θ extends to a homomorphism $\bar{\theta}: \mathcal{L}(S) \rightarrow \text{End}_k(V)$ by the universal property. Trivially $\bar{\theta}$ must be irreducible as well. Let f be a minimal nonzero idempotent of $T = \theta(S) = \bar{\theta}(\mathcal{L}(S))$. Then $\bar{\theta}^{-1}(f)$ is directed and so has a minimum element e .

Suppose that $e' \in \mathcal{D}e$. Suppose that $e'' < e'$. We claim that $\bar{\theta}(e'') = 0$. Indeed, choose $A \in \mathcal{L}(S)$ such that $A^{-1}A = e$ and $AA^{-1} = e'$. Then $A^{-1}e''A < A^{-1}e'A = e$ and so $\bar{\theta}(A^{-1}e''A) = 0$. Thus $\bar{\theta}(e'') = \bar{\theta}(AA^{-1}e''AA^{-1}) = 0$. We conclude that $\bar{\theta}$ is injective on the idempotents of D_e . Otherwise, we can find $e_1, e_2 \in D_e$ with $\bar{\theta}(e_1) = \bar{\theta}(e_2)$. Then $e_1e_2 \leq e_1, e_2$ and $\bar{\theta}(e_1) = \bar{\theta}(e_1e_2) = \bar{\theta}(e_2)$. Thus $e_1 = e_1e_2 = e_2$ by the above claim. We conclude that e has finite index since T has finitely many idempotents.

By choice of e , it now follows that $\bar{\theta}$ factors through S/I_e and hence is a $k_0[S/I_e]$ -module. Moreover, e is primitive in S/I_e . (If $I_e = \emptyset$, then we interpret $k_0[S/I_e]$ as kS and e is the minimum idempotent.) Since $eM = fM \neq 0$ by choice of f , it follows by Green's theorem that $N = eM$ is a simple $ek_0[S/I_e]e = kG_e$ -module, necessarily finite-dimensional. The identity map $N \rightarrow eM$ corresponds under the adjunction to

a nonzero homomorphism $\psi: \text{Ind}_e(N) \rightarrow M$. But we already know that $\text{Ind}_e(N)$ is simple by Proposition 4.4. Schur's lemma then yields that ψ is an isomorphism. This completes the proof. \square

References

- [1] J. Funk and P. Hofstra, 'Topos theoretic aspects of semigroup actions', *Theory Appl. Categ.* **24** (2010), 117–147.
- [2] J. A. Green, *Polynomial Representations of GL_n* , Lecture Notes in Mathematics, 830 (Springer, Berlin, 1980).
- [3] J. M. Howie, *Fundamentals of Semigroup Theory* (Clarendon Press, Oxford, 1995).
- [4] J. Kellendonk, 'The local structure of tilings and their integer group of coinvariants', *Commun. Math. Phys.* **187** (1997), 115–157.
- [5] J. Kellendonk, 'Topological equivalence of tilings', *J. Math. Phys.* **38** (1997), 1823–1842.
- [6] J. Kellendonk and M. V. Lawson, 'Tiling semigroups', *J. Algebra* **224** (2000), 140–150.
- [7] J. Kellendonk and M. V. Lawson, 'Universal groups for point-sets and tilings', *J. Algebra* **276** (2004), 462–492.
- [8] M. V. Lawson, 'Coverings and embeddings of inverse semigroups', *Proc. Edinb. Math. Soc.* **36** (1993), 399–419.
- [9] M. V. Lawson, *Inverse Semigroups* (World Scientific, Singapore, 1998).
- [10] M. V. Lawson, 'Ordered groupoids and left cancellative categories', *Semigroup Forum* **68** (2004), 458–476.
- [11] M. V. Lawson, 'A noncommutative generalization of Stone duality', *J. Austral. Math. Soc.* **88** (2010), 385–404.
- [12] M. V. Lawson and D. H. Lenz, 'Pseudogroups and their étale groupoids', arXiv:1107.5511v2.
- [13] M. V. Lawson and B. Steinberg, 'Ordered groupoids and étendues', *Cah. Topol. Geom. Differ. Categ.* **45** (2004), 82–108.
- [14] J. Leech, 'Inverse monoids with a natural semilattice ordering', *Proc. Lond. Math. Soc.* **70**(3) (1995), 146–182.
- [15] D. H. Lenz, 'On an order-based construction of a topological groupoid from an inverse semigroup', *Proc. Edinb. Math. Soc.* **51** (2008), 387–406.
- [16] S. W. Margolis and J. C. Meakin, 'Free inverse monoids and graph immersions', *Internat. J. Algebra Comput.* **3** (1993), 79–99.
- [17] W. D. Munn, 'Matrix representations of inverse semigroups', *Proc. Lond. Math. Soc.* **14**(3) (1964), 165–181.
- [18] A. L. T. Paterson, *Groupoids, Inverse Semigroups, and their Operator Algebras* (Birkhäuser, Boston, 1999).
- [19] M. Petrich, *Inverse Semigroups* (John Wiley & Sons, New York, 1984).
- [20] J. Renault, *A Groupoid Approach to C^* -algebras* (Springer, Berlin, 1980).
- [21] P. Resende, 'Étale groupoids and their quantales', *Adv. Math.* **208** (2007), 147–209.
- [22] R. L. Ruyle, 'Pseudovarieties of inverse monoids', PhD Thesis, University of Nebraska-Lincoln, 1997.
- [23] B. M. Schein, 'Representations of generalized groups', *Izv. Vyssh. Uchebn. Zaved. Mat.* **3** (1962), 164–176 (in Russian).
- [24] B. M. Schein, 'Noble inverse semigroups with bisimple core', *Semigroup Forum* **36** (1987), 175–178.
- [25] B. M. Schein, 'Infinitesimal elements and transitive representations of inverse semigroups', in: *Proc. Int. Symp. on the Semigroup Theory and its Related Fields*, Kyoto, 1990 (Shimane University, Matsue, 1990), 197–203.
- [26] B. M. Schein, 'Cosets in groups and semigroups', in: *Semigroups with Applications* (eds. J. M. Howie, W. D. Munn and H. J. Weinert) (World Scientific, Singapore, 1992), 205–221.

- [27] B. Steinberg, 'A topological approach to inverse and regular semigroups', *Pacific J. Math.* **208** (2003), 367–396.
- [28] B. Steinberg, 'A groupoid approach to discrete inverse semigroup algebras', *Adv. Math.* **223** (2010), 689–727.

M. V. LAWSON, Department of Mathematics and
Maxwell Institute for Mathematical Sciences, Heriot-Watt University,
Riccarton, Edinburgh EH14 4AS, UK
e-mail: markl@ma.hw.ac.uk

S. W. MARGOLIS, Department of Mathematics, Bar-Ilan University,
52900 Ramat Gan, Israel
e-mail: margolis@math.biu.ac.il

B. STEINBERG, Department of Mathematics, The City College of New York,
NAC 8/133, Convent Ave. at 138th Street, New York, NY 10031, USA
e-mail: bsteinberg@ccny.cuny.edu