

- Miami, 1994. *MMWR* 1994;43:773-775.
50. Centers for Disease Control and Prevention. Human rabies—West Virginia, 1994. *MMWR* 1995;44:86-87, 93.
 51. Centers for Disease Control and Prevention. Human rabies—Alabama, Tennessee, and Texas, 1994. *MMWR* 1995;44:269-272.
 52. Centers for Disease Control and Prevention. Human rabies—Washington, 1995. *MMWR* 1995;44:625-627.
 53. Centers for Disease Control and Prevention. Human rabies—Connecticut, 1995. *MMWR* 1996;45:207-209.
 54. Centers for Disease Control and Prevention. Human rabies—California, 1995. *MMWR* 1996;45:353-356.
 55. Moellering RC Jr. The Garrod lecture: the *Enterococcus*: a classic example of the impact of antimicrobial resistance on therapeutic options. *J Antimicrob Chemother* 1991;28:1-12.
 56. Frieden TR, Munsiff SS, Low DE, et al. Emergence of vancomycin-resistant enterococci in New York City. *Lancet* 1993;342:76-79.
 57. Cohen ML. Epidemiology of drug resistance: implications for a post-antimicrobial era. *Science* 1992;257:1050-1055.
 58. Hospital Infection Control Practices Advisory Committee. Recommendations for preventing the spread of vancomycin resistance. *MMWR* 1995;44(RR-12):1-13.
 59. Jernigan DB, Hofmann J, Cetron MS, et al. Outbreak of Legionnaires' disease among cruise ship passengers exposed to a contaminated whirlpool spa. *Lancet* 1996;347:494-499.
 60. Duchin JS, Koster F, Peters CJ, et al. Hantavirus pulmonary syndrome: clinical description of seventeen patients with a newly recognized disease. *N Engl J Med* 1994;330:949-955.
 61. Bell BP, Goldoft M, Griffin PM, et al. A multistate outbreak of *Escherichia coli* O157:H7-associated bloody diarrhea and hemolytic uremic syndrome from hamburgers: the Washington experience. *JAMA* 1994;272:1349-1353.
 62. Boyce TG, Pemberton AG, Wells JG, Griffin PM. Screening for *Escherichia coli* O157:H7—a nationwide survey of clinical laboratories. *J Clin Microbiol* 1995;33:3275-3277.
 63. Tenover FC, Tokars J, Swenson J, Paul S, Spitalny K, Jarvis W. Ability of clinical laboratories to detect antimicrobial agent-resistant enterococci. *J Clin Microbiol* 1993;31:1695-1699.

Outbreaks of *Cyclospora* Infections in US and Canada

**Gina Pugliese, RN, MS;
Martin S. Favero, PhD**
Medical News Editors

The CDC recently reported a number of outbreaks of *Cyclospora cayetanensis* infection in May and June 1996 in at least 10 states in the United States and Ontario, Canada. *Cyclospora cayetanensis* is a parasite that causes a diarrheal illness. Only three outbreaks have been reported in the United States since it was first diagnosed in humans in 1977.

Several hundred laboratory-confirmed cases have been reported to the CDC, mostly involving immunocompetent adults. Preliminary evidence suggests that consumption of raspberries and mixtures of berries and other fresh fruits may be associated with *Cyclospora* infection. The CDC, the FDA, and health officials in the

United States and Canada are collaborating to determine the extent and cause of the outbreaks, the sources of contamination, and whether transmission is ongoing. Additional efforts include testing samples of produce for *Cyclospora*.

Although *Cyclospora* is transmitted by the fecal-oral route, direct person-to-person transmission is unlikely, because excreted oocysts require days to weeks under favorable conditions to become infectious (ie, sporulate). The incubation period is 1 week, and illness may be protracted (from days to weeks), with frequent, watery stools and other gastrointestinal symptoms; symptoms may remit and relapse. Oocysts can be identified in the stool by examination of wet mounts under phase microscopy, use of modified acid-fast stains, or demonstration of autofluorescence with ultra-

violet epifluorescence microscopy. However, these procedures are not routine for most clinical laboratories and therefore confirmation by an experienced reference laboratory is recommended. Infection can be treated with trimethoprim-sulfamethoxazole.

The specific sources or modes of transmission have not been identified, and the complex distribution routes and handling of seasonal produce complicates tracing. As always, raw produce should be washed thoroughly before eating. Cases should be reported to local and state health departments.

FROM: Centers for Disease Control and Prevention. Outbreaks of *Cyclospora cayetanensis* infection—United States, 1996. *MMWR* 1996;45(25):549-551.