

A NOTE ON FREE ACTIONS OF GROUPS ON PRODUCTS OF SPHERES

JANG HYUN JO[✉] and JONG BUM LEE

(Received 14 September 2012; accepted 21 January 2013; first published online 8 March 2013)

Abstract

It has been conjectured that if $G = (\mathbb{Z}_p)^r$ acts freely on a finite CW -complex X which is homotopy equivalent to a product of spheres $S^{n_1} \times S^{n_2} \times \cdots \times S^{n_k}$, then $r \leq k$. We address this question with the relaxation that X is finite-dimensional, and show that, to answer the question, it suffices to consider the case where the dimensions of the spheres are greater than or equal to 2.

2010 *Mathematics subject classification*: primary 57S25; secondary 22E25.

Keywords and phrases: CW -complex, free action, free rank, solvmanifolds, strongly polycyclic groups.

1. Introduction

The purpose of this note is to give a contribution to the following long-standing conjecture in algebraic topology, which remains unanswered despite considerable effort by many mathematicians.

CONJECTURE 1.1. If $G = (\mathbb{Z}_p)^r$ acts freely on a finite CW -complex which is homotopy equivalent to a product of spheres $S^{n_1} \times S^{n_2} \times \cdots \times S^{n_k}$, then $r \leq k$.

In fact, the converse of Conjecture 1.1 is also open, that is, it is not known that any finite group G with $\text{rk } G = r$ can act freely on a finite (or finite-dimensional) CW -complex which is homotopy equivalent to a product of spheres $S^{n_1} \times S^{n_2} \times \cdots \times S^{n_r}$, where $\text{rk } G$ is the maximum of p -ranks $\text{rk}_p G$ as p runs over the primes dividing $|G|$. The case of a single sphere for Conjecture 1.1 and its converse were solved by Smith [13], Milnor [11], Swan [14] and Madsen *et al.* [10]. Moreover, a generalisation of this result has been established by Adem and Smith [2] and Mislin and Talelli [12]. The affirmative answer to Conjecture 1.1 for the case of free $(\mathbb{Z}/p)^r$ -actions on $(S^n)^k$ with trivial action on homology was first given by Carlsson [5]

The first author has been supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012R1A1A2006395). The second author has been supported by the Mid-career Researcher Program through an NRF grant funded by the MEST (No. 2010-0008640) and by a Sogang University Research Grant for 2010.

© 2013 Australian Mathematical Publishing Association Inc. 0004-9727/2013 \$16.00

using a spectral sequence argument. Browder [4] and Benson and Carlson [3] gave other beautiful proofs of this result via different approaches. Adem and Browder [1] settled the homologically nontrivial case except in the cases $p = 2$ and $n = 1, 3$ or 7 . Subsequently, Yalçın [15] solved the case for $p = 2$ and $n = 1$, leaving the other two cases open. The nonequidimensional case is known for the product of two spheres [8]. Recently, Hanke [7] gave the most general result using the subject of tame homotopy theory for Conjecture 1.1.

To settle the case for $p = 2$ and $n = 1$, Yalçın used the following algebraic result to prove the case of $p = 2$ and $n = 1$ for Conjecture 1.1.

THEOREM 1.2 [15, Theorem 3.2]. *Let G be a finite group, and let $0 \rightarrow \mathbb{Z}^k \rightarrow \Gamma \rightarrow G \rightarrow 1$ be an extension of G . If Γ is torsion-free, then $\text{rk}(G) \leq k$.*

Recently, we generalised Yalçın's result from free abelian groups to strongly polycyclic groups.

THEOREM 1.3 [9, Theorem 3.2]. *Let Γ be a strongly polycyclic group and G be a finite group. Let $1 \rightarrow \Gamma \rightarrow \Pi \rightarrow G \rightarrow 1$ be an extension. Then $\text{rk}_p(G) - \text{rk}_p(\Pi) \leq h(\Gamma)$ for all primes p with $p \mid |G|$, where $h(\Gamma)$ is the Hirsch rank of Γ .*

Note that Yalçın's topological application hinges on the fact that the k -torus is a finite dimensional $K(\mathbb{Z}^k, 1)$ -space. In particular, $\pi_1(G \setminus X)$ is torsion-free, where X is a free G -complex with the homotopy type of the k -torus. Thus Theorem 1.3 sheds some light on the following conjecture which is relaxed with the condition of finite-dimensionality.

CONJECTURE 1.4. If $G = (\mathbb{Z}_p)^r$ acts freely on a finite-dimensional CW -complex which is homotopy equivalent to a product of spheres $S^{n_1} \times S^{n_2} \times \cdots \times S^{n_k}$, then $r \leq k$.

The purpose of this note is to show that, to prove Conjecture 1.4, it suffices to consider the case where the dimensions of the spheres are greater than or equal to 2. We will consider this in a more general setting in Theorem 2.2.

CONJECTURE 1.5. If $G = (\mathbb{Z}_p)^r$ acts freely on a finite-dimensional CW -complex which is homotopy equivalent to a product of spheres $S^{n_1} \times S^{n_2} \times \cdots \times S^{n_k}$ and all $n_i \geq 2$, then $r \leq k$.

2. Main results

Recall that for a finite-dimensional CW -complex X and p a prime, the free p -rank of X , denoted by $\text{frk}_p(X)$, is the largest r such that $(\mathbb{Z}_p)^r$ acts freely on X [16, Definition 1.1]. Similarly, we can give the following definition to handle Conjectures 1.4 and 1.5.

DEFINITION 2.1. Let X be a finite-dimensional CW -complex and p a prime. The homotopy free p -rank of X , denoted by $\text{hfrk}_p(X)$, is defined as the largest r such that $(\mathbb{Z}_p)^r$ acts freely on Y , where Y is a finite-dimensional CW -complex homotopy equivalent to X .

By definition, $\text{frk}_p(X) \leq \text{hfrk}_p(X)$ for any finite-dimensional CW-complex X . It is well known that the following statements hold:

- (1) $\text{frk}_p(S^n) = \text{hfrk}_p(S^n) = 1$ for n odd;
- (2) $\text{frk}_p(S^n \times S^m) \leq \text{hfrk}_p(S^n \times S^m) \leq 2$ [8].

Note also that if Conjecture 1.4 is true, then $\text{hfrk}_p(S^{n_1} \times S^{n_2} \times \dots \times S^{n_k}) \leq k$. From [16] it is known that $\text{hfrk}_p(\prod^k L_p^{2n-1}) = k$, where L_p^{2n-1} is the lens space. We do not know of an example of a finite-dimensional CW-complex satisfying the inequality $\text{frk}_p(X) < \text{hfrk}_p(X)$. It is conceivable that $\text{frk}_p(X) = \text{hfrk}_p(X)$.

THEOREM 2.2. *Let M be a compact solvmanifold and L a finite-dimensional CW-complex with a finite fundamental group $\pi_1(L)$. Then*

$$\text{hfrk}_p(M \times L) \leq \dim M + \text{hfrk}_p(L).$$

PROOF. Suppose that $G = (\mathbb{Z}_p)^r$ acts freely on a finite-dimensional CW-complex X which is homotopy equivalent to the product $M \times L$. Let $F = \pi_1(L)$, $\Gamma = \pi_1(M)$ and $\pi = \pi_1(G \setminus X)$. Then the free G -action on X induces a short exact sequence

$$1 \longrightarrow \Gamma \times F \longrightarrow \pi \longrightarrow G \longrightarrow 1.$$

Since F is a characteristic subgroup of $\Gamma \times F$, the above short exact sequence yields the following commutative diagram of short exact sequences:

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & \\
 & & \uparrow & & \uparrow & & \\
 1 & \longrightarrow & \Gamma & \longrightarrow & \pi/F & \longrightarrow & G \longrightarrow 1 \\
 & & \uparrow & & \uparrow & & \uparrow = \\
 1 & \longrightarrow & \Gamma \times F & \longrightarrow & \pi & \longrightarrow & G \longrightarrow 1 \\
 & & \uparrow & & \uparrow & & \\
 & & F & \xrightarrow{=} & F & & \\
 & & \uparrow & & \uparrow & & \\
 & & 1 & & 1 & &
 \end{array}$$

Note that M is a $K(\Gamma, 1)$ -space whose dimension is the Hirsch length $h(\Gamma)$ of Γ and Γ is a strongly polycyclic group [6, 9]. Note also that the top horizontal exact sequence

$$1 \longrightarrow \Gamma \longrightarrow \pi/F \longrightarrow G \longrightarrow 1$$

is associated to the following maps:

$$\begin{array}{c} \bar{X} \simeq \widetilde{M} \times L \simeq L \\ \Gamma \downarrow \\ X \simeq M \times L \\ G \downarrow \\ G \setminus X \end{array}$$

By Theorem 1.3, we have $\text{rk}_p(G) \leq \text{rk}_p(\pi/F) + h(\Gamma) = \text{rk}_p(\pi/F) + \dim M$. Note that every finite subgroup H of π/F is isomorphic to a subgroup of G . Since π/F acts freely on \bar{X} , the subgroup H acts freely on \bar{X} , which is homotopy equivalent to L . Hence $\text{rk}_p(\pi/F) \leq \text{hfrk}_p(L)$ and therefore $\text{rk}_p(G) \leq \dim(M) + \text{hfrk}_p(L)$. \square

COROLLARY 2.3. *Conjecture 1.5 implies Conjecture 1.4.*

PROOF. Let $G = (\mathbb{Z}_p)^r$. Suppose that G acts freely on a finite dimensional CW-complex X which is homotopy equivalent to $S^{n_1} \times S^{n_2} \times \dots \times S^{n_k}$. Without loss of generality, we may assume that $n_1 = 1$ and $n_2, \dots, n_k \geq 2$. Write $M = S^1$ and $L = S^{n_2} \times \dots \times S^{n_k}$. Conjecture 1.5 implies that $\text{hfrk}_p(L) \leq k - 1$. By Theorem 2.2, $\text{rk}_p(G) \leq 1 + k - 1 = k$. \square

COROLLARY 2.4. *If $G = (\mathbb{Z}_p)^r$ acts freely on a finite-dimensional CW-complex X which is homotopy equivalent to a product $(S^1)^\ell \times Y$, where Y is one of the following:*

- (1) $S^n \times S^m$ where $n, m \geq 2$;
- (2) $\prod^k L_p^{2n-1}(a_1, \dots, a_n)$, where $L_p^{2n-1}(a_1, \dots, a_n)$ is the standard lens space with $(a_i, p) = 1$ for all i ;

then $r \leq \ell + 2$ for case (1) and $r \leq \ell + k$ for case (2).

PROOF. This follows from combining the results of [8, 16] with Theorem 2.2. \square

References

- [1] A. Adem and W. Browder, ‘The free rank of symmetry of $(S^n)^k$ ’, *Invent. Math.* **92** (1988), 431–440.
- [2] A. Adem and J. H. Smith, ‘Periodic complexes and group actions’, *Ann. of Math.* (2) **154** (2001), 407–435.
- [3] D. J. Benson and J. F. Carlson, ‘Complexity and multiple complexes’, *Math. Z.* **195** (1987), 221–238.
- [4] W. Browder, ‘Cohomology and group actions’, *Invent. Math.* **71** (1983), 599–608.
- [5] G. Carlsson, ‘On the rank of abelian groups acting freely on $(S^n)^k$ ’, *Invent. Math.* **69** (1982), 393–400.
- [6] F. T. Farrell and L. E. Jones, *Classical Aspherical Manifolds*, CBMS Regional Conference Series in Mathematics, 75 (American Mathematical Society, Providence, RI, 1990).
- [7] B. Hanke, ‘The stable free rank of symmetry of products of spheres’, *Invent. Math.* **178** (2009), 265–298.

- [8] A. Heller, 'A note on spaces with operators', *Illinois J. Math.* **3** (1959), 98–100.
- [9] J. H. Jo and J. B. Lee, 'Group extensions and free actions by finite groups on solvmanifolds', *Math. Nachr.* **283** (2010), 1054–1059.
- [10] I. Madsen, C. B. Thomas and C. T. C. Wall, 'The topological spherical space form problem. II. Existence of free actions', *Topology* **15** (1976), 375–382.
- [11] J. Milnor, 'Groups which act on S^n without fixed points', *Amer. J. Math.* **79** (1957), 623–630.
- [12] G. Mislin and O. Talelli, 'On groups which act freely and properly on finite dimensional homotopy spheres', in: *Computational and Geometric Aspects of Modern Algebra (Edinburgh, 1998)*, London Mathematical Society Lecture Note Series, 275 (Cambridge University Press, Cambridge, 2000), 208–228.
- [13] P. A. Smith, 'Permutable periodic transformations', *Proc. Nat. Acad. Sci.* **30** (1944), 105–108.
- [14] R. G. Swan, 'Periodic resolutions for finite groups', *Ann. of Math. (2)* **72** (1960), 267–291.
- [15] E. Yalçın, 'Group actions and group extensions', *Trans. Amer. Math. Soc.* **352** (2000), 2689–2700.
- [16] E. Yalçın, 'Free actions of p -groups on products of lens spaces', *Proc. Amer. Math. Soc.* **129** (2001), 887–898.

JANG HYUN JO, Department of Mathematics, Sogang University,
Seoul 121-742, Korea
e-mail: jhjo@sogang.ac.kr

JONG BUM LEE, Department of Mathematics, Sogang University,
Seoul 121-742, Korea
e-mail: jlee@sogang.ac.kr