

continued for 5 days after antimicrobial discontinuation. Focus groups and interviews were conducted to identify barriers, and the implementation strategy was adapted to address the key identified barriers. The implementation strategy included clinical decision support involving a linked flag on antibiotic ordering and a 1-click order entry within the electronic medical record (EMR), provider and patient education (written/videos/in-person), and local site champions. Protocol adherence was measured by tracking the number of patients on therapeutic antimicrobials that received BioK+ based on the bedside nursing EMR medication administration records. Adherence rates were sorted by hospital and unit in 48- and 72-hour intervals with recording of percentile distribution of time (days) to receipt of the first antimicrobial. **Results:** In total, 340 education sessions with >1,800 key stakeholders occurred before and during implementation across the 4 involved hospitals. The overall adherence of probiotic ordering for wards with antimicrobial orders was 78% and 80% at 48 and 72 hours, respectively over 72 patient months. Individual hospital adherence rates varied between 77% and 80% at 48 hours and between 79% and 83% at 72 hours. Of 246,144 scheduled probiotic orders, 94% were administered at the bedside within a median of 0.61 days (75th percentile, 0.88), 0.47 days (75th percentile, 0.86), 0.71 days (75th percentile, 0.92) and 0.67 days (75th percentile, 0.93), respectively, at the 4 sites after receipt of first antimicrobial. The key themes from the focus groups emphasized the usefulness of the linked flag alert for probiotics on antibiotic ordering, the ease of the EMR 1-click order entry, and the importance of the education sessions. **Conclusions:** Electronic clinical decision support, education, and local champion support achieved a high implementation rate consistent across all sites. Use of a 1-click order entry in the EMR was considered a key component of the success of the implementation and should be considered for any implementation strategy for a stewardship initiative. Achieving high prescribing adherence allows more precision in evaluating the effectiveness of the probiotic strategy.

Funding: Partnerships for Research and Innovation in the Health System, Alberta Innovates/Health Solutions **Funding:** Award

Disclosures: None

Doi:[10.1017/ice.2020.849](https://doi.org/10.1017/ice.2020.849)

Presentation Type:

Poster Presentation

Implementing a Centralized Surveillance and Validation Program for Infection Prevention

Lauren Le Goff, The Children's Hospital of Philadelphia; Sarah Smathers, The Children's Hospital of Philadelphia; Lauren Satchell, Children's Hospital of Philadelphia; Lori Handy, Children's Hospital of Philadelphia; Julia Sammons, The Children's Hospital of Philadelphia

Background: Mandatory reporting of all healthcare-associated infections (HAIs) leads to substantial surveillance volume for infection prevention and control (IPC) programs. Prior to 2019, 6 infection preventionists were performing system-wide surveillance for all infection types using NHSN definitions at a large quaternary-care center in Pennsylvania. Limited surveillance validation was performed. With the continued expansion of the health system, increased demands for IPC expertise, and a growing team, the need for streamlined surveillance, and a validation program were identified. **Methods:** A surveillance training program for novice team members was developed and implemented. Infection prevention associates (IPAs), whose primary role was data management, began training. The new program included NHSN training videos, direct observation of surveillance with infection

preventionists and practice case studies. Following training, IPAs performed surveillance for experienced infection preventionists covering high-risk inpatient units. To ensure high reliability, surveillance validation was initiated. Each month, ~10% of investigated infections were randomly pulled from the electronic surveillance system and divided among experienced infection preventionists. These validators performed unbiased reviews of the charts based on limited data, including patient demographics and culture results. Validation documentation included noting whether an infection was reportable to NHSN and a rationale. Data on whether or not each patient had a complex medical history and time spent validating each case were collected. Compliance of validator documentation aligning with original documentation was tracked. Discrepancies were discussed as a team and were adjudicated as needed. IPAs tracked hours spent on surveillance to capture effort transitioned from infection preventionists. **Results:** Between March and July 2019, an average of 223 (range, 178–261) potential infections were reviewed per month. From March through June 2019, 61 infections were selected for validation, with 98% compliance with original documentation. One minor discrepancy was attributed to interpretation of documentation in the medical record. Medical complexity accounted for 78% of reviews and validation time spent averaged 12 minutes per infection (range, 3–28 minutes). Self-reported effort directed from infection preventionists to 2 IPAs for surveillance was ~20 hours per week. An additional IPA was hired to perform surveillance in addition to other job responsibilities. **Conclusions:** Centralized surveillance programs can promote high reliability and cost-efficient IPC staffing for large healthcare systems, especially those with mandatory reporting requirements or medically complex patient populations. Improving surveillance skills among associate staff can increase experienced infection preventionist bandwidth for project management, staff supervision, and other leadership responsibilities. Lastly, validation programs are crucial to ensuring quality assurance of data reporting to both internal and external stakeholders.

Funding: None

Disclosures: None

Doi:[10.1017/ice.2020.850](https://doi.org/10.1017/ice.2020.850)

Presentation Type:

Poster Presentation

Implementing a Massive Personal Protective Equipment Education—A Multidisciplinary Team Approach

Mayar Al Mohajer, Baylor College of Medicine; Megan Fischer, Baylor St. Luke's Medical Center; Melissa Rouse, Baylor St. Luke's Medical Center; Takei Pipkins, Baylor St. Luke's Medical Center; John Byrne, Baylor St. Luke's Medical Center

Background: Personal protective equipment (PPE) is defined by the Occupational Safety and Health Administration as specialized clothing or equipment worn by an employee for protection against infectious materials. They include gloves, gowns, masks, respirators, googles and face shields. The CDC has issued guidelines on appropriateness of when, what, and how to use PPE. Despite these guidelines, compliance with PPE remains challenging. **Methods:** We implemented a massive hospital-wide rapid education program on PPE donning and doffing of all employees and staff. This program included an online video, return demonstration and just-in-time training. To develop the program, we recorded PPE training video, reviewed PPE validation checklist, developed new isolation precaution signage with quick response (QR) code to video, developed a nutrition tray removal video and a equipment cleaning video, developed family and visitor guidelines for isolation precautions, and created an audit