

## O279

**Factors affecting cognitive remediation outcome in schizophrenia: The role of treatment resistance**

M. Spangaro<sup>1\*</sup>, M. Bosia<sup>2</sup>, F. Martini<sup>1</sup>, M. Bechi<sup>1</sup>, M. Buonocore<sup>1</sup> and R. Cavallaro<sup>2</sup>

<sup>1</sup>Clinical Neurosciences, IRCCS San Raffaele Scientific Institute, milano, Italy and <sup>2</sup>Faculty Of Medicine, Vita-Salute San Raffaele University, Milan, Italy

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.446

**Introduction:** Treatment-resistant schizophrenia (TRS) represents a major clinical issue, characterized by worse psychopathological outcome, a more disrupted neurobiological substrate and higher health-care costs. Cognitive impairment is a core feature of schizophrenia, strongly associated with patients' functional outcome. Different studies showed that TRS patients exhibit poorer neurocognitive performance, particularly on verbal domains. To date Cognitive Remediation Therapy (CRT) represents the best available tool for treating cognitive deficits in schizophrenia. However, CRT outcomes are highly heterogeneous and significant treatment predictors are still lacking.

**Objectives:** To investigate possible differences of CRT outcome among patients with schizophrenia, stratified according to antipsychotic response (TRSs vs. first-line responders - FLRs).

**Methods:** 150 patients with schizophrenia, (95 FLRs, 55 TRSs) were assessed for neurocognition with BACS and WCST at baseline and after CRT. General Linear Models (GLMs) were performed to investigate possible differences between groups on basal cognition and CRT outcome (Cohen's d Effect Size).

**Results:** At baseline, GLMs showed significant differences in Verbal Memory ( $F=4,66$ ;  $p=0,03$ ) and WCST-executive functions ( $F=5,59$ ;  $p=0,02$ ), both worse in TRS group. Effect Sizes of CRT outcome resulted significantly different in domains of Verbal Memory ( $F=4,68$ ;  $p=0,03$ ) and WCST-executive functions ( $F=4,62$ ;  $p=0,03$ ), with greater improvements among TRS patients.

**Conclusions:** This is the first study to indicate treatment-resistance as a possible predictor of CRT outcome in schizophrenia. Moreover, we observed that CRT resulted able to fill the cognitive gap between treatment groups. Thus, these results further highlight the importance of early cognitive interventions in order to reduce the neuropsychological and functional burden associated with the disease, especially for TRS patients.

**Disclosure:** No significant relationships.

**Keywords:** treatment resistance; cognitive remediation; schizophrénia

## O276

**Influenza and schizophrenia: How can we shed a light in the new virus from an old association?**

D. Magalhães<sup>1\*</sup>, F. Ferreira<sup>1</sup>, T. Ferreira<sup>1</sup>, I. Figueiredo<sup>2</sup>, F. Martinho<sup>1</sup>, R. Felício<sup>2</sup> and N. Santos<sup>1</sup>

<sup>1</sup>Psychiatry, Hospital Professor Doutor Fernando Fonseca, Lisboa (Amadora), Portugal and <sup>2</sup>Mental Health Department, Hospital Professor Doutor Fernando Fonseca, Lisboa (Amadora), Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.447

**Introduction:** COVID-19 raises serious concerns regarding its unknown consequences for health, including psychiatric long term

outcomes. Historically, influenza virus has been responsible for pandemics associated with schizophrenia. Epidemiological studies showed increased risk for schizophrenia in children of mothers exposed to the 1957 influenza A2 pandemic. Controversy remains concerning the mechanisms of pathogenesis underlying this risk.

**Objectives:** We aim to review the evidence for the association between influenza infection and schizophrenia risk, the possible pathogenic mechanisms underlying and correlate these findings with the schizophrenia hypothesis of neurodevelopment.

**Methods:** We reviewed literature regarding evidence from epidemiological, translational animal models and serological studies using medline database.

**Results:** The biological mechanisms likely to be relevant account to the effects of infection-induced maternal immune activation, microglial activation, infection-induced neuronal autoimmunity, molecular mimicry of the influenza virus, neuronal surface auto-antibodies and psychosis with potential infectious antecedents. Influenza infection may fit into the theory of the neurodevelopment of schizophrenia as a factor that alters the normal maturation processes of the brain (possible second or third hit).

**Conclusions:** Influenza infection has multiple pathogenic pathways in both pre and post natal processes that might increase the risk of schizophrenia or psychosis. The existing evidence regarding the relationship between influenza virus and psychosis might help us draw similar long-term concerns of COVID-19.

**Disclosure:** No significant relationships.

**Keywords:** schizophrénia; influenza; viral; infection

## O277

**Negative symptoms in first episode schizophrenia: Results from the “parma early psychosis” program**

L. Pelizza<sup>1\*</sup>, E. Leuci<sup>1</sup>, D. Maestri<sup>1</sup>, E. Quattrone<sup>1</sup>, G. Paulillo<sup>1</sup>, P. Pellegrini<sup>1</sup> and S. Azzali<sup>2</sup>

<sup>1</sup>Department Of Mental Health, Azienda USL di Parma, Parma, Italy and <sup>2</sup>Department Of Mental Health, Azienda USL-IRCCS di Reggio Emilia, Reggio Emilia, Italy

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.448

**Introduction:** Identifying distinct dimensions of negative symptoms in First Episode Schizophrenia (FES) might result in a better understanding and treatment of this invalidating symptomatology.

**Objectives:** Aim of this study was to examine negative symptom structure in FES patients using the Positive and Negative Syndrome Scale (PANSS).

**Methods:** All 147 participants, aged 12–35 years, completed the PANSS and the Global Assessment of Functioning (GAF) scale. A principal component analysis with varimax rotation was performed to investigate PANSS negative symptom structure in the FES total sample.

**Results:** A 2-factor model (i.e. “Expressive Deficits” and “Asociality” dimensions) was identified. Only “Expressive Deficits” domain had a significant negative correlation with baseline GAF score.

**Conclusions:** This bipartite solution seems to be adequate to describe the phenomenological variety of negative symptoms experienced by FES individuals at the point of entry in early intervention services.

**Disclosure:** No significant relationships.

**Keywords:** psychopathology; negative symptoms; schizophrénia; early psychosis

## Sexual medicine and mental health

## O279

## The prevalence of anxiety and depression in transgender people living in Russia

E. Chumakov<sup>1,2\*</sup>, Y. Ashenbrenner<sup>1</sup>, N. Petrova<sup>2</sup>, M. Zastrozhin<sup>3,4</sup>, L. Azarova<sup>1</sup> and O. Limankin<sup>1,5,6</sup>

<sup>1</sup>Day Inpatient Department, St-Petersburg Psychiatric Hospital No 1 named after P.P. Kaschenko, Saint-Petersburg, Russian Federation; <sup>2</sup>Department Of Psychiatry And Addiction, Saint-Petersburg University, Saint-Petersburg, Russian Federation; <sup>3</sup>Department Of Addictology, Russian Medical Academy Of Continuous Professional Education, Moscow, Russian Federation; <sup>4</sup>Laboratory Of Genetics And Basic Research, Moscow Research & Practical Centre on Addictions of the Moscow Department of Healthcare, Moscow, Russian Federation; <sup>5</sup>Department Of Psychotherapy, Medical Psychology And Sexology, North-Western State Medical University named after I.I. Mechnikov, Saint-Petersburg, Russian Federation and <sup>6</sup>Department Of Social Psychiatry And Psychology, St-Petersburg Institute of Postgraduate Improvement of Physicians-experts of the Ministry of Labour and Social Protection, Saint-Petersburg, Russian Federation

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.449

**Introduction:** The prevalence rates of mental health issues, particularly anxiety and depression, is high among transgender people. However, the incidence of anxiety and depression in transgender people living in Russia is unclear until now.

**Objectives:** To examine the frequency of anxiety and depression in transgender people living in Russia.

**Methods:** The Hospital Anxiety and Depression Scale (HADS) was used for online screening for symptoms of anxiety and depression in transgender people living in Russia throughout November 2019. 588 transgender adults living in all Federal Districts of Russia (mean age 24.0±6.7) were included in the final analysis. 69.6% (n=409) of the survey participants indicated the direction of transition as transmasculine (TM), 23.1% (n=136) – as transfeminine (TW), and 7.3% (n=43) – as other (TO).

**Results:** It was found that 45.1% (n=265) and 24.0% (n=141) of transgender people had clinically significant levels of anxiety and depression, respectively (HADS score of 11 or higher). The rates of anxiety (TM=10.21±4.68; TW=8.72±3.91; TO=10.72±4.43) and depression (TM=7.53±4.09; TW=7.40±4.19; TO=7.74±4.33) did not have statistically significant differences within the direction of transition. The anxiety and depression mean scores in all subgroups were statistically significantly higher than in the general Russian population (p<0.001; one sample t-test).

**Conclusions:** Our findings suggest a high prevalence of depression and anxiety disorders in the transgender population as compared to the cisgender population in Russia. The identified frequency of anxiety and depression in transgender people in Russia is worrying and requires immediate action to improve the availability and quality of medical and psychological care for this group of people.

**Disclosure:** No significant relationships.

**Keywords:** Depression; Transgender; Anxiety

## O280

## Sexual fantasies, subjective satisfaction and quality of sexual life in patients of sexual dysfunction: A comparative study

N. Ohri<sup>1\*</sup>, A. Dubey<sup>2</sup>, G. Vankar<sup>3</sup>, P. Rathi<sup>2</sup> and A. Gill<sup>1</sup>

<sup>1</sup>Psychiatry, New Life Hospital, Varanasi, India; <sup>2</sup>Psychiatry, Sri Aurobindo Medical College and PGI, Indore, India and <sup>3</sup>Psychiatry, Parul Institute of Medical Sciences, Vadodara, India

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.450

**Introduction:** Exploring the ways in which sexual fantasies may affect sexual experience and satisfaction is of relevance in the clinical setting involving sexual dysfunction.

**Objectives:** To observe how the sexual fantasy scores differ in their relationship with sexual satisfaction, experience and quality between sexual dysfunction cases and normal controls.

**Methods:** Scales included: Wilson's sex fantasy questionnaire (WSFQ), Arizona Sexual Experience Scale (ASEX), Sexual Quality of Life Questionnaire (SQoL), and a subjective sexual satisfaction meter. Differences in responses of both groups on WSFQ (item-wise and domain-wise) were analysed using T-tests. Two-way ANOVA was applied to see how other scales affected sexual fantasy.

**Results:** Cases scored significantly higher on ASEX scale, and low on satisfaction, SQoL and WSFQ

	Cases N=100	Controls N=100	t-test
Satisfaction Mean(sd)	4.27(1.85)	7.82(1.31)	t=3.052;df=198, p=0.0026
Asex	17.52(4.73)	8.28(1.34)	t=15.24;df=198, p<0.0001
SQoL	29.41(12.12)	49.5(6.67)	t=14.52;df=198, p<0.0001
WSFQ	26.80(17.61)	30.59(15.32)	t=1.62,df=98,p=0.106

Majority of WSFQ responses, both in cases and controls, fell in the intimate and impersonal domains. Sexual fantasy scores and sexual satisfaction had a strong positive and significant correlation in controls but no linear correlation in the case-subjects. sexual fantasy scores contributed to 5.7% of difference in the scores of SQoL between groups. Major variance in scores of satisfaction in our subjects depended on presence or absence of sexual dysfunction (46.5%) but sexual fantasies also contributed to 8.8% of the variance.

**Conclusions:** The study showed that fantasies contribute to positive sexual outcomes only in absence of sexual dysfunction. ANOVA analysis revealed that in case-subjects sexual satisfaction briefly increases initially with increase in fantasy scores but starts to decline as fantasies increase.

**Disclosure:** No significant relationships.

**Keywords:** Sexual Dysfunction; sexual fantasy; Sexual Quality of Life; Sexual experience