MICROSCOPY TODAY




MARCH 1994

ISSUE #94-2

Aurora... The Next Step In SPM Technology

Sanning Near-Field Optical Microscopy, SNOM, is the latest advance in the evolution of Scanning Probe Microscopy. And Aurora,[™] from TopoMetrix, is the first production SNOM system available to researchers.

SNOM is the SPM technique that breaks the diffraction barrier to spatial resolution. It expands SPM's nanometer-scale imaging



SNOM image (above) and topographic image (right) of eye tissue were acquired simultaneously by Aurora.

capability by adding an optical probe which yields information that complements data from other SPM techniques. It combines all the advantages of optical microscopy, including contrast mechanisms and spectroscopic methods, with the high resolution of SPM. Resolution that's better than the best conventional or confocal microscope by an order-ofmagnitude or more.

Aurora is the newest member of TopoMetrix' TMX 2000 family of SPMs. It features proven piezoelectric scanners, optics, control electronics and software.



Aurora lets you collect data in either reflection or transmission mode while you are simultaneously gathering topographic data.

New applications for Aurora are being uncovered every day. Besides being the natural extension to every existing optical-microscopy application, it can be used for optical lithography, highdensity data storage, polarization imaging of magneto optics and polymers, and to characterize light-wave devices like semiconductor lasers and single-mode optical fibers. And that's just the beginning.

To learn more about Aurora, call us today at **1-800-765-5067**. We'll get product literature and sample images on their way to you right away.



TopoMetrix, 5403 Betsy Ross Drive, Santa Clara, California, 95054 Tel 408-982-9700, Fax 408-982-9751

New Light On An Old Problem

Jean-Paul Revel, CALTECH

Like many other people I find myself not having the time to do all the things that I would like to do in a day. I have always wished I could emulate the young lady in Prof. Arthur Bullers (1874-1944) limerick which was published in the December 19, 1933 issue of that reputed journal, Punch:

"There was a young lady named Bright Whose speed was faster than light. She set out one day, In a relative way, And returned home the previous night."

Well, up to now I have not succeeded in my attempts to gain time in this way, but there is hope, there is hope as I will try to make clear in a minute.

First of course, as we all know, the speed of light is finite. This was first demonstrated by Roemer in 1676. He measured the interval between successive eclipses of the intermost satellite of Jupiter (that would be lo, with its sulfur spewing volcanoes), at a time when Jupiter was close to Earth. From his data he calculated when the 100th eclipse should occur, and (I surmise) invited his friends to come and observe. Well the eclipse did not materialize when he had expected it! Here they all were, waiting for the little moon to cooperate and it did not! And it was not by seconds that the event was amiss but by a whopping 15 minutes. With all his friends gone home shaking their heads, with doubts about his ability to predict anything, Roemer started thinking about this and realized that, because of where they were in their orbits, the distance between Earth and Jupiter was increasing and that his embarrassment could be explained by the fact that light from the satellite would therefore take longer to arrive to his telescope. He calculated that his observations could be accounted for if the speed of light was 48,303 leagues/second. As we know, this number, which corresponds in the units we prefer today to 214,000 km/sec, was in fact of the correct order of magnitude. With improved methods for measuring the speed of light we now use the figure of 299,792,75 km/sec. Since the speed of light is finite there is always a possibility that one could better it, and thereby regain time?

Now what gave me hope that this might in fact already have been achieved, is a paper by A.M Steinberg, P.G. Kwiat and R.Y. Chiao of the Department of Physics at the University of California at Berkeley, on "Measurement of the Single-Photon Tunneling Time", published in Physical Review Letters 1993, 71 pp. 708-711. They say "Using a two-photon interferometer, we have measured the time delay for a photon to tunnel across a barrier consisting of a 1.1 mµ thick ID photonic band-gap material. The peak of the photon wave packet appears on the far side of the barrier 1.47 -0.21 fs earlier (their italics!) than it would if it were to travel at the vacuum speed of light c ... " Now all I have to do, thinks I, is to somehow take advantage of these photons and I'll be in the company of Ms. Bright whenever I need time to finish all the things I want to do. Too bad that the next sentence of the abstract quoted above, goes on to say "Although the apparent tunneling velocity (1.7 +/- 0.2) cis superluminal", imagine! almost twice the speed of light! "this is not a genuine signal velocity and Einstein causality is not violated ... "What a let down! As explained by Rolf Landauer¹ there are other instances in which there appears to be superluminal velocity, which does not really mean that the "accepted light velocity barrier" has "crumbled". Imagine for example watching the light fromga lighthouse beacon impinging on a distant wall. The spot it forms may "move faster that the velocity of the light, but this is not a violation of relativity; one spot is not the source of the next". It is easy to understand this image, but the restor the story leaves me somewhat puzzled. I feel a bit like Chaim Weizmann, the President of Israel, who said after a transatlantic crossing in the company bf Albert Einstein: "he explained his theory to me every day, and on arrival I was fully convinced that he understood it." The neat thing about this is that I can keep on dreaming with even higher hopes that I had before about catching up, someday, somehow.

Reference

Landauer, R. 1993. Light Faster than Light? Nature, 365, pp 692-3

**************** **INTERNATIONAL READERS:**

While we wish to extend our overseas readership, we logically want to mail the newsletter only to those who wish to receive copies. A "?" following your name on the address of this copy indicates that, as of 14 March, we have not received an indication of that interest. While a first copy for some, this will be the last copy for all who have not requested a subscription.

To request a no cost subscription, you have only to send your 🙀 mailing label back to us or, if you wish, advise your interest by Fax. And, while you are at it, you might inquire if others in your 🜟 organization would like a copy - and advise.

And, as we agree that this newsletter could be much more t interesting, we would appreciate any and all contributions. - - - Don Grimes, Editor

Front Cover Image

A large variety of shapes of fullerenes have been reported in the literature. Regardless of their shape, all these particles have a hollow core surrounded by graphite layers of 0.34 nm spacing. Particles in the form of squares, polygons and tubes up to several microns in length have been observed. The hollow core is a sphere, or a cylindrical cavity in the case of tubes, of about 2 nm diameter. The heart shaped example shown here is guite different in the size and shape of the core, but it is still surrounded by graphite 0.34 nm layers. Notably, it was observed on February 14, indicating date rather than time dependence in the growth process of fullerenes. Happy belated Valentine!

> Augusto A. Morrone University of Florida

MICROSCOPY TODAY

A monthly newsletter dedicated to the unique interests in microscopy of, and mailed at no cost to,

some 14,500 microscopists worldwide.

PO Box 620122, Middleton, WI 53562 - Tel.: (608)836-1970 - Fax: 608)836-1969 - eMail: 74250.331@COMPUSERVE.COM Copyright in 1994 by Today Enterprises, All rights reserved

Don Grimes, Editor