

with 20 mg of intravenous diazepam. Because catatonia was required for these symptoms, there was no NMS; rather, there was a combination of definite catatonia and neuroleptic effects. In order to demonstrate NMS the authors would have had to observe drug-induced symptoms in the absence of catatonia, which they did not; their assertion of separate episodes of both catatonia and NMS is not only unproven, it is untested. The care of catatonic patients has always been risky, and the clinical state of this patient may have been as unstable as NMS, but that does not make it NMS.

GREENFIELD, D., CONRAD, C., KINCARE, P., *et al* (1987) Treatment of catatonia with low-dose lorazepam. *American Journal of Psychiatry*, **144**, 1224–1225.

POPE, H. G., KECK, P. E., JR. & MCELROY, S. L. (1986) Frequency and presentation of neuroleptic malignant syndrome in a large psychiatric hospital. *American Journal of Psychiatry*, **143**, 1227–1233.

SHALEV, A., HERMESH, H. & MUNITZ, H. (1989) Mortality from neuroleptic malignant syndrome. *Journal of Clinical Psychiatry*, **50**, 18–25.

CONRAD MELTON SWARTZ

*Department of Psychiatric Medicine  
East Carolina University  
Greenville  
North Carolina  
USA*

#### 'Mabi bark tea' consumption and psychosis?

SIR: Hassiotis *et al* (*Journal*, September 1992, **161**, 404–407) suggest that the use of mabi bark may be the precipitating cause of a psychotic illness in a 23-year-old West Indian woman. They cite the temporal link between ingestion of the mabi bark drink and the onset of her psychosis, the lack of a previous personal or family history of mental illness, an identical twin not developing a similar illness, and a biochemical basis that mabi bark causes central dopamine release.

We find the hypothesis untenable. Firstly, our observations in Trinidad and Tobago are not consistent with this. Mabi bark drink (mauby) is very widely used by the Trinidad and Tobago population and has not been found to contribute to the onset of psychotic illness. In a recent analysis of 634 schizophrenic patients, the period prevalence rate was found to be 5.0/1000 population with the disorder being three times more common in African Trinidadians than Indian Trinidadians. The use of mauby was investigated and was not found to be a contributory factor in a single case. Mauby has been drunk here for over a century and continues to be a very popular drink. Their patient's consumption of mauby was described as high but two-thirds of a pint

daily for a week can hardly be described as excessive. In addition, for many years it has been served as a cold drink on the psychiatric wards and to date, no reports of its association with psychosis have been made.

In considering the above, it is clear that the association between mabi bark and this girl's psychotic illness is a spurious one and more than likely to be a chance association. There are too many uncontrolled confounding variables. Why implicate mauby? Why not coffee or tea?

HARI D. MAHARAJH  
GERARD A. HUTCHINSON

*St Ann's Hospital  
Port of Spain  
Trinidad  
West Indies*

#### Simultaneous kidney disease and manic-depressive psychosis

SIR: We would like to report the first recorded case of simultaneous transmission of autosomal dominant polycystic kidney disease (ADPKD) and manic-depressive psychosis (MDP).

*Case reports.* Mrs F (aged 51) had her first episode of brief mental disorder at the age of 25 after the birth of her first child. After the birth of her second child five years later she had a further depressive illness which was treated with medication and electroconvulsive therapy (ECT). At the same time she was diagnosed as suffering from chronic renal failure due to ADPKD. In 1976 she had a very prolonged episode of treatment-resistant depressive illness. She was very depressed in mood, psychotic, and almost stuporose. However, after a long period of illness she recovered completely. She has had several further episodes of a prolonged psychotic depressive illness, particularly in 1989 when she was admitted to hospital for over six months – receiving antidepressants, *L*-tryptophan, lithium, and 24 applications of ECT. She has had lithium augmentation for treatment of her depressive illness since 1978. Since 1985 she has been on peritoneal dialysis. Her maternal grandmother had some kind of kidney disease but died in her 70s of an unknown cause. Her husband is well. They have two children.

Her elder son has had an episode of severe depression of similar magnitude to that experienced by his mother. He first presented at the age of 25 with a six-week history of lethargy and depressed mood. He was treated with fluvoxamine without response. He then became suicidal, taking several overdoses and cutting his wrists. On the day of admission he had been found by his flatmate trying to electrocute himself using wires from an electric lamp. He improved slowly with antidepressants and was transferred to Day Care, without regaining his premorbid state. He relapsed, was readmitted and responded very well to ECT.