

review; we suspect that this is partly explained by underdocumentation of criteria such as stool frequency. In healthcare settings where appropriateness of HO-CDI testing is not optimal, mandatory ID attending physician approval may provide an option beyond clinical decision-support tools.

**Funding:** None

**Disclosures:** None

Doi:10.1017/ice.2020.636

### Presentation Type:

Poster Presentation

### Appropriateness of Initiating Antibiotics for Urinary Tract Infection Among Nursing Home Residents

Taniece R. Eure MPH, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention; Nicola D. Thompson MS, PhD, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention; Austin Penna MPH, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention; Wendy M. Bamberg MD, Medical Epidemiology Consulting, Colorado Department of Public Health and Environment; Grant Barney BS, New York Emerging Infections Program; Devra Barter MS, Colorado Department of Public Health and Environment; Paula Clogher MPH, Connecticut Emerging Infections Program and the Yale School of Public Health; Malini DeSilva MD, MPH, Minnesota Department of Health; Ghinwa Dumyati MD, New York Emerging Infections Program, University of Rochester Medical Center; Erin Epton MD, California Department of Health; Christina B. Felsen MPH, New York Emerging Infections Program, University of Rochester Medical Center; Linda Frank RN, BSN, PHN, California Emerging Infections Program; Deborah Godine RN, California Emerging Infections Program; Lourdes Irizarry MD, New Mexico Department of Health; Helen Johnston MPH, Colorado Department of Public Health and Environment; Marion A. Kainer MBBS, MPH, Tennessee Department of Health; Linda Li MPH, Maryland Department of Health; Ruth Lynfield MD, Minnesota Department of Health; JP Mahoehney RN, MPH, Minnesota Department of Health; Joelle Nadle MPH, California Emerging Infections Program; Valerie L. S. Ocampo MPH, BSN, RN, Oregon Health Authority; Susan M. Ray MD, Georgia Emerging Infections Program, Emory University; Monika E. Samper RN, Oregon Health Authority; Sarah Shrum Davis MPH, New Mexico Department of Health; Marla Sievers MPH, New Mexico Department of Health; Krithika Srinivasan MD, MPH, Connecticut Emerging Infections Program and the Yale School of Public Health; Lucy E. Wilson MD, ScM, Maryland Department of Health; Alexia Y. Zhang MPH, Oregon Health Authority; Shelley S. Magill MD, PhD, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention; Nimalie D. Stone MD, Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention

**Background:** Antibiotics are among the most commonly prescribed drugs in nursing homes; urinary tract infections (UTIs) are a frequent indication. Although there is no gold standard for the diagnosis of UTIs, various criteria have been developed to inform and standardize nursing home prescribing decisions, with the goal of reducing unnecessary antibiotic prescribing. Using different published criteria designed to guide decisions on initiating treatment of UTIs (ie, symptomatic, catheter-associated, and uncomplicated cystitis), our objective was to assess the


appropriateness of antibiotic prescribing among NH residents. **Methods:** In 2017, the CDC Emerging Infections Program (EIP) performed a prevalence survey of healthcare-associated infections and antibiotic use in 161 nursing homes from 10 states: California, Colorado, Connecticut, Georgia, Maryland, Minnesota, New Mexico, New York, Oregon, and Tennessee. EIP staff reviewed resident medical records to collect demographic and clinical information, infection signs, symptoms, and diagnostic testing documented on the day an antibiotic was initiated and 6 days prior. We applied 4 criteria to determine whether initiation of treatment for UTI was supported: (1) the Loeb minimum clinical criteria (Loeb); (2) the Suspected UTI Situation, Background, Assessment, and Recommendation tool (UTI SBAR tool); (3) adaptation of Infectious Diseases Society of America UTI treatment guidelines for nursing home residents (Crnich & Drinka); and (4) diagnostic criteria for uncomplicated cystitis (cystitis consensus) (Fig. 1). We calculated the percentage of residents for whom initiating UTI treatment was appropriate by these criteria. **Results:** Of 248 residents for whom UTI treatment was initiated in the nursing home, the median age was 79 years [IQR, 19], 63% were female, and 35% were admitted for postacute care. There was substantial variability in the percentage of residents with antibiotic initiation classified as appropriate by each of the criteria, ranging from 8% for the cystitis consensus, to 27% for Loeb, to 33% for the UTI SBAR tool, to 51% for Crnich and Drinka (Fig. 2). **Conclusions:** Appropriate initiation of UTI treatment among nursing home residents remained low regardless of criteria used. At best only half of antibiotic treatment met published prescribing

Figure 1. Summary of criteria used to assess appropriateness of antibiotic treatment initiation for urinary tract infections.

Loeb Criteria (ICHE 2001)	UTI SBAR Tool (AHRQ 2012)	Crnich & Drinka (Annals of LTC 2014)	Cystitis Consensus (JAMA 2018)
<b>Purpose</b>			
To establish minimum criteria for initiating antibiotics	To promote better communication of information needed for antibiotic decision making	To establish minimum criteria for initiating antibiotics, based on published guidelines	To identify criteria for uncomplicated cystitis and discourage prescribing for asymptomatic bacteriuria
<b>Signs of Systemic Infection</b>			
<ul style="list-style-type: none"> <li>Fever</li> <li>Rigors</li> <li>Mental status change (delirium)</li> </ul>	<ul style="list-style-type: none"> <li>Fever</li> <li>Rigors</li> <li>Mental status change (delirium)</li> <li>Hypotension</li> </ul>	<ul style="list-style-type: none"> <li>Fever</li> <li>Rigors</li> <li>Mental status change (delirium)</li> <li>Unstable Vital Signs               <ul style="list-style-type: none"> <li>Decreased oxygenation</li> <li>Respiratory rate <math>\geq</math>25 breaths per minute</li> <li>Hypotension</li> </ul> </li> </ul>	None
<b>Localizing Urinary Signs and Symptoms</b>			
<ul style="list-style-type: none"> <li>Dysuria</li> <li>Urgency</li> <li>Frequency</li> <li>Hematuria</li> <li>Suprapubic pain</li> <li>Incontinence</li> <li>Costovertebral angle tenderness</li> </ul>	<ul style="list-style-type: none"> <li>Dysuria</li> <li>Urgency</li> <li>Frequency</li> <li>Hematuria</li> <li>Suprapubic pain</li> <li>Incontinence</li> <li>Costovertebral angle tenderness</li> </ul>	<ul style="list-style-type: none"> <li>Dysuria</li> <li>Urgency</li> <li>Frequency</li> <li>Hematuria</li> <li>Suprapubic pain</li> <li>Incontinence</li> <li>Costovertebral angle tenderness</li> <li>Scrotal / Prostate Tenderness</li> <li>Purulent urethral discharge</li> </ul>	<ul style="list-style-type: none"> <li>Dysuria</li> <li>Urgency</li> <li>Frequency</li> <li>Hematuria</li> <li>Suprapubic pain</li> </ul>
<b>Diagnostic Testing</b>			
None	None	Urine Culture Urinalysis	None

Fig. 1.

Figure 2. Appropriateness of Antibiotic Initiation in Nursing Home (NH) Residents for NH initiated, active treatment of UTI (n=248)



**Note:** Please refer to Figure 1 – the summary of criteria used to assess appropriateness of antibiotic treatment initiated for urinary tract infection, for signs of systemic infection and localizing urinary signs and symptoms for each of the criteria listed above.

<sup>1</sup> Denominator for Cystitis Consensus is less than the total due to the exclusion of possible complicated urinary tract infections. These were residents with an indwelling catheter and/or signs and symptoms indicative of pyelonephritis, renal abscess, and prostatitis.

Fig. 2.

criteria. Although insufficient documentation of infection signs, symptoms and testing may have contributed to the low percentages observed, adequate documentation in the medical record to support prescribing should be standard practice, as outlined in the CDC Core Elements of Antibiotic Stewardship for nursing homes. Standardized UTI prescribing criteria should be incorporated into nursing home stewardship activities to improve the assessment and documentation of symptomatic UTI and to reduce inappropriate antibiotic use.

**Funding:** None

**Disclosures:** None

Doi:10.1017/ice.2020.637

**Presentation Type:**

Poster Presentation

**Area-Based Socioeconomic Status Measures and Incidence of Community-Associated ESBL-Producing Enterobacteriaceae, 2017**

Hannah E. Reses, Centers for Disease Control and Prevention; Cedric Brown, CDC Erin C. Phipps Kristina G. Flores; Ghinwa Dumyati, University of Rochester; Rebecca Tsay; Marion Kainer, Western Health; Daniel Muleta, Tennessee Department of Health; Nadezhda Duffy, Centers for Disease Control and Prevention; Isaac See, Centers for Disease Control and Prevention