The Journal of General Microbiology

Volume 52, Part 2, July 1968

- J. ŠMARDA AND U. TAUBENECK Situation of colicin receptors in surface layers of bacterial cells
- J. D. A. MILLER, J. E. HUGHES, G. F. SAUNDERS AND L. L. CAMPBELL Physiological and biochemical characteristics of some strains of sulphate-reducing bacteria
- P. DRAPER AND P. D'ARCY HART The composition of normal and elongated Mycobacterium lepraemurium
- C. RODRIGUEZ-BARRUECO The occurrence of the root-nodule endophytes of Alnus glutinosa and Myrica gale in soils
- T. BAUCHOP AND E. A. DAWES The metabolism of formic acid by Sarcina ventriculi
- J. C. ANAND AND A. D. BROWN Growth rate patterns of the so-called osmophilic and non-osmophilic yeasts in solutions of polyethylene glycol
- I..G. MURRAY Some aspects of the biochemical differentiation of pathogenic fungi: a review
- R. J. ROWBURY The inhibitory action of α -methylmethionine on Escherichia coli
- A. STENDERUP AND A. LETH BAK Deoxyribonucleic acid base composition of some species within the genus Candida
- s. E. JUHASZ Growth retardation, colonial changes and nutritional deficiency in *Mycobacterium phlei* due to lysogeny
- J. R. WARR A mutant of Chlamydomonas reinhardii with abnormal cell division
- A. L. SCHADE, N. H. MYERS AND R. W. REINHART Carbohydrate metabolism and production of diffusible active substances by *Staphylococcus aureus* grown in serum at iron levels in excess of siderophilin iron-saturation and below
- R. F. HILL Modification of lethality and mutagenesis by growth inhibition of ultraviolet-irradiated *Escherichia coli* strain B/R
- H. J. WILLETTS Stromatal rind formation in the brown rot fungi
- w. J. RYAN An X-factor requiring Haemophilus species
- R. J. W. BYRDE AND A. H. FIELDING Pectin methyl-trans-eliminase as the maceration factor of Sclerotinia fructigena and its significance in brown rot of apple
- E. R. LEADBETTER AND S. C. HOLT The fine structure of Bacillus fastidiosus
- J. L. MEERS AND D. W. TEMPEST The influence of extracellular products on the behaviour of mixed microbial populations in magnesium-limited chemostat cultures

45s. net (\$7.50). Annual subscription £25 net (\$87.50) for fifteen issues.

CAMBRIDGE UNIVERSITY PRESS

Advertising rates and data may be obtained from

Mr D. J. French
Journal Sales Promotion
Cambridge University Press
Bentley House
200 Euston Road
London, N.W.1, England

Miss Gretta Goldenman 32 East 57th Street New York, N.Y.10022 U.S.A.

or

MICHROME STAINS

Methyl Green Aquamount Celestin Blue Celloidin *Chromosome Red Fluoroil Fluormount Thionine
May-Grünwald
Giemsa Stain
Janus Green B
Laktoseal
Lipid Crimson
Lacmoid

Leishman Nile Blue Optoil Sudan Black *Trifalgic Acid Wright Stain Pyronine Y etc.

Detection of

PROTEIN-MALNUTRITION

&

State of Health in Man and Animals

can be accomplished most readily and easily in the field or in the laboratory by use of B. T. Squires' modified *FALG STAIN.

*RHODANILE BLUE: an entirely new kind of basic polychrome dye with many uses.

*Information leaflets and catalogue on request

Also dyes for scientific and technological purposes

EDWARD GURR, LTD.

42 Upper Richmond Road West LONDON, S.W.14

Cables: Micromlabs, London, S.W. 14 Telephones: 01-876 8228/9

Publications issued by

COMMONWEALTH INSTITUTE OF ENTOMOLOGY 56 Queen's Gate, London, S.W. 7

BULLETIN OF ENTOMOLOGICAL RESEARCH

Published quarterly and containing original articles on economic entomology. *Post free*. Annual subscription (payable in advance) 150s. od. Prices of back parts and volumes on application.

REVIEW OF APPLIED ENTOMOLOGY

Consisting of abstracts or reviews of current literature on economic entomology throughout the world. Published monthly in two series:

Series "A" dealing with insect and other arthropod pests of cultivated plants, forest trees and stored products of animal and vegetable origin.

Series "B" dealing with insects, ticks, etc. conveying disease or otherwise injurious to man and animals. Post free.

Series "A" Series "B"
Annual subscription (payable in advance) 180s. od. 80s. od.
Prices of back parts and volumes on application.

ZOOLOGICAL RECORD, part INSECTA

Published annually and containing as complete a record as possible of the literature of the previous year, chiefly from the systematic standpoint.

Annual subscription 130s. od. Post free.

Prices of back volumes on application.

NOTES FOR CONTRIBUTORS

CONTRIBUTIONS will be welcomed from scientists of all nationalities, but must be written in English. Papers should be sent to Dr. H. D. Crofton, Department of Zoology, University of Bristol, Bristol 2.

GENERAL. Submission of a paper will be taken to imply that it is unpublished and is not being considered for publication elsewhere. Papers published in *Parasitology* may not be reprinted or published in translation without permission from the Editorial Board. Papers should be as concise as clarity permits, and illustrations kept as few as possible. Attention to the following instructions will assist rapid publication:

TYPESCRIPTS. Papers should be typewritten with double spacing, on one side only of quarto paper, leaving a margin of about 4 cm. on the left-hand side of each sheet. Sheets should be numbered consecutively. Heavily corrected typescripts may have to be returned to the author for retyping. Papers should state the name of the author and the address to which proofs are to be sent. The address of the laboratory at which the work was carried out will be printed with the authors' names at the head of the paper. A short title, not exceeding 44 letters, should be supplied for running headlines.

The general arrangement of the paper should be as follows: Introduction; Methods; Results; Discussion; Summary of results; References. Tables, explanations of text-figures or plates, and keys to the abbreviations used on figures should be on separate sheets (see below).

PLATES should only be included where they are absolutely essential, as in the reproduction of photomicrographs and similar illustrations demanding very fine detail. Plates are expensive and authors may be required to bear part of the cost if an unduly large number is required. Photographs should be sharp glossy prints, trimmed and mounted on white card and grouped and numbered as they are required to appear in print (oversea contributors may prefer to submit unmounted prints with a sketch to indicate the proposed grouping, in order to reduce postage costs). Please remember that the area of a group cannot exceed 20 by 12·5 cm., after reduction (if any). Lettering is best indicated on an overlay or separate sketch.

TEXT FIGURES should not be more than about twice the size of the finished block, and the thickness of lines and size of points, stipples, etc. determined accordingly. The maximum size of any one group of figures, after reduction, is 20 by 12.5 cm. They must be drawn in Indian ink on white Bristol Board, heavy drawing paper, tracing paper or tracing linen; graph paper ruled in pale blue (but not other colours) is also acceptable. Explanatory lettering should be lightly inserted in blue pencil, with tally lines showing the position of the feature to which it refers, so that the printer may put in the finished lettering. Properly prepared scraperboard drawings may be suitable for reproduction as line drawings, but drawings with gradations of shading (e.g. washdrawings) are not. Please try to avoid creasing or folding illustrations. Legends to illustrations must be given on a separate sheet of paper, but scales of magnification are preferably given alongside the illustrations. Each illustration must have the name of the author and figure number pencilled on the back. Plates and

diagrams should be numbered separately and their approximate positions indicated in the margin of the typescript. It is very convenient if photographs of illustrations are sent with the papers in addition to the original figures; they are essential when the originals are very large, so that they may be sent through the post to referees.

TABLES should be kept to a minimum and the duplication of information in tables and graphs avoided. They should be typed on separate sheets of paper and their approximate positions indicated in the margin of the typescript. Each table should be numbered and be designed to be printed in the normal orientation of the text with the data so grouped as to make the use of rules unnecessary, whenever possible.

NOMENCLATURE. Authors should follow the International Rules of Nomenclature in the names of organisms. When new names are introduced the recommendations of the International Code of Zoological Nomenclature, 1964, should be strictly adhered to, especially with regard to the designation of types and statements as to where such types are deposited.

REFERENCES should be given only when cited in the text and should be listed in alphabetical order at the end of the paper, in the following form:

at the end of the paper, in the following form:
Surname of author(s), initials; year of publication; title of paper; title of journal (abbreviated according to the World List of Scientific Periodicals, 1963–5 ed., and underlined); volume number; first and last page numbers of the work cited. When books are listed, the title of the book should be underlined and it should be followed by the town of publication and the name of the publisher.

Where there is more than one reference to a particular author or group of authors with the same date, references should be distinguished by letters placed after the date, the letter a being used for the first reference cited and so on.

PROOFS sent to authors are already in page form and only essential corrections should be made on them. Excessive alterations, other than corrections of printers' errors, may be disallowed or charged to the author; they are in any case liable to delay publication. Correction should be made using the symbols in British Standard 1219: 1958, or its shortened version B.S. 1219 C: 1958 (British Standards Institution, 2 Park Street, London, W. 1). After correction, the printers' marked proof should be returned to the Editor.

OFFPRINTS. Fifty offprints of each paper are provided free of charge. Additional offprints may be ordered on the form sent out with proofs, provided this is returned within seven days of receipt.

MORE DETAILED INFORMATION on the preparation of manuscripts for publication is to be found in the following:

NUTTALL, G. H. F. (1940). Notes on the Preparation of Papers for Publication in the Journal of Hygiene and in Parasitology, Parasitology 32, 1-62 and J. Hyg. 40, 1-62. (Now out of print and also out of date in some respects.)

The Journal of Physiology (1960). Suggestions to Authors 150, 1-33, Cambridge University Press.

Authors 150, 1-33, Cambridge University Press.

The Royal Society published a second edition of General Notes on the Preparation of Scientific Papers in 1965.

PARASITOLOGY

Volume 58, Part 3 August 1968

CONTENTS

	PAGE
BARON, P. J. On the histology and ultrastructure of Cysticercus longicollis, the cysticercus of Taenia crassiceps Zeder, 1800, (Cestoda, Cyclophyllidea). (With 3 Plates and 6 Figures in the Text)	497
LAVOIPIERRE, M. M. J. and RAJAMANICKAM, C. The skin reactions of two species of Southeast Asian Chiroptera to notoedrid and teinocoptid mites. (With 2 Plates and 9 Figures in the Text)	515
Krishnamurthy, R. A new flagellate of the genus <i>Proteromonas</i> Kunstler, 1883, from an Indian lizard. (With 9 Figures in the Text)	531
TAYLOR, E. W. and THOMAS, J. N. Membrane (contact) digestion in the three species of tapeworm Hymenolepis diminuta, Hymenolepis microstoma and Moniezia expansa. (With 3 Figures in the Text)	535
FERNANDO, S. T. Immunological response of the hosts to <i>Toxocara canis</i> (Werner, 1782) infection. I. Effect of superinfection on naturally infected puppies. (With 3 Figures in the Text)	547
McLaren, Diane J. and Paget, G. E. A fine structural study on the merozoite of <i>Eimeria tenella</i> with special reference to the conoid apparatus. (With 4 Plates and 1 Figure in the Text)	561
Howell, M. J. The life-cycle of <i>Echinoparyphium serratum</i> sp.nov. (Digenea: Echinostomatidae). (With 8 Figures in the Text)	573
Howell, M. J. Excystment and in vitro cultivation of Echinoparyphium serratum. (With 1 Plate and 5 Figures in the Text)	583
NEAL, R. A. and JOHNSON, PAULINE. The virulence to rats of five <i>Entamoeba histolytica</i> -like strains capable of growth at 25 °C and attempts to discover similar strains.	599
THAKUR, AMAR S. and CHENG, THOMAS C. The formation, structure, and histochemistry of the metacercarial cyst of <i>Philophthalmus gralli</i> Mathis & Léger. (With 1 Plate and 2 Figures in the Text)	605
ESCH, GERALD W. Studies on carbohydrates in tissues and coenurus fluid of larval <i>Taenia</i> multiceps from normal and alloxanized mice	619
JARRETT, ELLEN E. E., JARRETT, W. F. H. and URQUHART, G. M. Quantitative studies on the kinetics of establishment and expulsion of intestinal nematode populations in susceptible and immune hosts. <i>Nippostrongylus brasiliensis</i> in the rat. (With 5 Figures in the Text)	625
BARRETT, J. The effect of temperature on the development and survival of the infective larvae of Strongyloides ratti Sandground 1925. (With 4 Figures in the Text)	641
BITAKARAMIRE, P. K. Lymnaea natalensis laboratory culture and production of Fasciola gigantica metacercariae.	653
ROGERS, W. P. Neurosecretory granules in the infective stage of <i>Haemonchus contortus</i> . (With 2 Plates and 1 Figure in the Text).	657
CALLOW, L. L. The infection of Boophilus microplus with Babesia bigemina	663
WHITFIELD, P. J. A histological description of the uterine bell of <i>Polymorphus minutus</i> (Acanthocephala). (With 9 Figures in the Text)	671
Dowding, Valerie M. The formation of the cuticular ridges in the larval pharynx of the blowfly (Calliphora vicina RD.). (With 7 Figures in the Text)	683
Long, P. L. The pathogenic effects of <i>Eimeria praecox</i> and <i>E. acervulina</i> in the chicken. (With 5 Figures in the Text)	691
KIM, KE CHUNG. Two new species of the sucking lice (Hoplopleuridae, Anoplura) from <i>Rattus</i> (Muridae, Rodentia) in Thailand. (With 17 Figures in the Text)	701
HORTON-SMITH, C., LONG, P. L. and LEE, D. L. Observations on the tissue and post-tissue larval development of <i>Ascaridia dissimilis</i> Vigeuras 1931, and a comparison with that of <i>Ascaridia galli</i> , in turkeys. (With 1 Plate)	709
JOYNER, L. P. and PURNELL, R. E. The feeding behaviour on rabbits and in vitro of the Ixodid tick Rhipicephalus appendiculatus Neumann, 1901. (With 3 Figures in the Text)	715
Purnell, R. E. and Joyner, L. P. The development of <i>Theileria parva</i> in the salivary glands of the tick, <i>Rhipicephalus appendiculatus</i> . (With 1 Plate and 1 Figure in the Text)	725
McConnachie, Elspeth W. The growth of an axenic strain of Entamoeba invadens in	
different media. (With 4 Figures in the Text)	733