

CPD Online

- 73 **New from CPD Online**

Cochrane corner

- 74 **Physical health care monitoring for people with serious mental illness**
Graeme Tosh, Andrew V. Clifton, Jun Xia & Margueritte M. White
- 74 **General physical health advice for people with serious mental illness**
Graeme Tosh, Andrew V. Clifton, Jun Xia & Margueritte M. White

Round the corner

- 75 **Commentary Physical monitoring and advice: absence of evidence is not evidence of absence**
Katharine Smith

Editorials

- 78 **Long-term effects of antipsychotics**
Joanna Moncrieff
- 80 **Early schizophrenia: skilful management of medication**
John Cookson
- 85 **Patients need reliable facts and figures about antipsychotics**
Stephen M. Lawrie

Articles

- 88 **High-dose antipsychotics: addressing patients' resistance to physical health monitoring**
Paula Marie Murphy, Andrew Iles & Suja Sreedharan
- 98 **Why are long-acting injectable antipsychotics still underused?**
Peter Bosanac & David Jonathan Castle
- 106 **Psychiatric assessment and treatment of survivors of torture**
Richard M. Duffy & Brendan D. Kelly
- 116 **In-patient services for people with intellectual disability and mental health or behavioural difficulties**
John Devapriam, Alan Rosenbach & Regi Alexander
- 124 **Sleep disorders in children and adolescents**
Gregory Stores
- 132 **Drug and alcohol addiction: do psychosocial treatments work?**
Jason Luty

Cover image

Bosanac & Castle (pp. 98–105) discuss the place of long-acting injectable antipsychotics (LAIs) in the continuum of care. They argue that their use should be considered more often with patients requiring long-term treatment, even early in the course of their illness.
Image: ©iStockphoto/MilosJokic.