

## Regulatory responses to excess zinc ingestion in growing rats

Tomoya Fujimura, Tohru Matsui and Masayuki Funaba\*

Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwakecho, Kyoto 606-8502, Japan

(Submitted 14 April 2011 – Final revision received 27 July 2011 – Accepted 31 July 2011 – First published online 12 September 2011)

### Abstract

The growth of weaning piglets is effectively improved by feeding a high-Zn diet (3000 mg Zn/kg of diet). The present study examined whether feeding a diet supplemented with Zn (1016–3000 mg/kg) for 10 d induces growth benefits in rats. In addition, tissue weight, Zn content of tissues and expression of Zn transporters were examined in these rats. Zn supplementation did not significantly increase body weight. Breaking line model analyses indicated that the weight of the pancreas, the organ most sensitive to excess Zn, significantly decreased with increasing Zn intake beyond 15.2 mg/d. Excess Zn has been suggested to accumulate in the liver, kidney and bone in order to protect the pancreas. Zn concentrations in the plasma, liver, kidney and femur increased with increasing Zn intake up to approximately 30 mg/d, whereas those in the pancreas increased up to 8.4 mg/d and decreased by Zn intake beyond 8.4 mg/d. The expression levels of the Zn transporters *Zip4* and *ZnT1* in the intestinal epithelium were significantly lower in rats fed a diet supplemented with 1016 mg/kg Zn compared to those fed the basal diet. The present study reveals that (1) excess Zn intake does not accelerate growth in rats, but is detrimental to the pancreas, (2) the excess Zn is effectively accumulated in the liver, kidney and bone, without sufficient protection of the pancreas and (3) expression of Zn transporters is down-regulated in response to excess Zn intake.

**Key words:** Excess zinc: Zinc accumulation: Zinc transporters: Growth

Zn is an essential mineral that acts as a cofactor for numerous enzymes and transcription factors<sup>(1)</sup>. The physiological responses to Zn deficiency are well-characterised in mammals<sup>(2–5)</sup>, whereas less information is available regarding the effects of excess Zn intake. The pancreas has been suggested as the tissue most sensitive to excess Zn<sup>(6)</sup>. The National Research Council (NRC)<sup>(7)</sup> proposes that Zn accumulates in tissues such as the liver, kidney and bone, in order to protect other organs from failure induced by Zn accumulation. Zn concentrations were increased in the liver (6-fold) and in the kidney (11-fold) in preruminant calves fed a diet supplemented with 500–700 mg Zn/kg, whereas the increases in Zn concentration in the heart and muscle were relatively smaller<sup>(8)</sup>; similar results were obtained in sheep fed a diet supplemented with 700–2100 mg Zn/kg<sup>(9)</sup>. Furthermore, Zn concentrations in the liver, kidney and bone were higher in rats fed a diet supplemented with 2438 mg Zn/kg compared with growing rats fed a diet containing 38 mg Zn/kg, and the Zn concentration of these tissues plateaued at supplementation levels of 2438–7238 mg Zn/kg<sup>(10)</sup>.

Elevated Zn intake (3000 mg/kg) for a period of 14 d surprisingly induces growth in weaning piglets<sup>(11–13)</sup>. Considering that the Zn requirement for growing pigs is 100 mg/kg<sup>(14)</sup>,

the Zn-induced growth promotion results from the pharmacological effects of excess Zn intake. The physiological basis for these pharmacological effects remains unclear. ZnO possesses antimicrobial properties<sup>(15)</sup>, but several studies suggest that ZnO promotes growth in early-weaned and conventionally weaned pigs, regardless of diarrhoea prevalence or intestinal microbial numbers<sup>(16–18)</sup>.

Zn homeostasis is primarily maintained by regulation of its absorption and secretion. Several Zn transporters of the Slc39 (*Zip*) and Slc30 (*ZnT*) families have been identified. Members of the *Zip* family have been shown to increase the cytosolic Zn concentration, whereas those of the *ZnT* family decrease the cytosolic concentration<sup>(19–22)</sup>. *Zip4* and *ZnT1* are involved in Zn absorption in the small intestine<sup>(23,24)</sup>, whereas *Zip5* is responsible for intestinal Zn secretion<sup>(25)</sup>. In addition, Zn is secreted from the pancreas into the gut by *Zip5* and *ZnT1*<sup>(25,26)</sup>. Zn transporter activities are modulated in response to Zn depletion through alteration of gene expression, transporter translocation, or both<sup>(22)</sup>.

We hypothesised that growth promotion induced by excess Zn intake (3000 mg/kg) is not limited to weaning piglets but is instead observed in other animals, including rats. In addition, it was hypothesised that each tissue grows proportionally in

**Abbreviations:** BW, body weight; *Hprt1*, hypoxanthine phosphoribosyltransferase 1; *Igf-1*, insulin-like growth factor-1; *Mt*, metallothionein; NRC, National Research Council.

\* **Corresponding author:** M. Funaba, fax +81 75 753 6344, email mfunaba@kais.kyoto-u.ac.jp

rats fed diets supplemented with excess Zn. In order to examine these points, we examined body and tissue growth, accumulation of Zn in tissues and expression of Zn transporters in growing rats fed diets supplemented with excess Zn. We specifically examined (1) whether growth promotion induced by excess Zn is observed in growing rats, (2) whether the concept proposed by the NRC regarding the prevention from Zn toxicity is applicable and (3) whether Zn transporter gene expression is altered in response to excess Zn ingestion. Our results indicate that excess Zn ingestion did not enhance the growth performance in growing rats, but in fact decreased pancreatic weight. Unexpectedly, the gene transcript levels of both the intestinal Zn transporters involved in Zn absorption and those involved in secretion were decreased in rats fed diets with higher Zn contents.

## Materials and methods

### Animals and diets

The experiments were approved by the Kyoto University Animal Experiment Committee (20-19). A total of twenty-eight male specific pathogen-free Sprague–Dawley rats aged 4 weeks were housed individually in stainless-steel cages under constant conditions (24°C, 50% humidity) with a fixed light–dark cycle (lights on from 05.00 to 19.00 hours). Because excess Zn-induced growth promotion in weaning pigs is not necessarily due to the antimicrobial effects as described earlier, we used specific pathogen-free rats. After a 5 d acclimatisation period of feeding the basal diet (24 mg Zn/kg) shown in Table 1, rats were randomly assigned to receive diets with differing Zn concentrations. The requirement of Zn in growing rats is 12 mg/kg<sup>(27)</sup>; the basal diet contained twice as much Zn<sup>(27)</sup>. Zn content in the diet recommended

for growing rats by AIN<sup>(28)</sup> is 30 mg/kg. All groups were allowed free access to food and distilled water for the 10 d study period. The diets were prepared by addition of ZnO to a Zn-deficient diet at the expense of glucose to provide a 24, 1016, 2008 or 3000 mg Zn/kg diet, and the actual measured content was 23.8, 1050, 2090 or 3200 mg Zn/kg diet, respectively. Since this study tested responses to dietary Zn status, i.e. excess intake as well as deficiency, egg-white was used as a protein source. D-Biotin was added to the basal diet due to the high avidin content of egg-white in order to prevent biotin deficiency. Body weight (BW) and feed consumption were measured every day.

At the end of the 10 d experimental period, rats were killed by bleeding from the abdominal aorta under isoflurane anaesthesia. Tissues (the liver, kidney, pancreas, spleen, small intestine, testis, gastrocnemius muscle, femur and perirenal fat pad) were collected and weighed. Blood collected with a heparinised syringe was centrifuged at 2500 g for 30 min at 4°C to obtain the plasma. The intestine was flushed with saline and scraped with slide glass to obtain the intestinal epithelium. Other tissues were rinsed in saline, immediately frozen in liquid N<sub>2</sub> and stored at –80°C until analysis.

### Determination of mineral contents

After wet-ash digestion of diets, plasma, and tissues with trace-element grade nitric acid and H<sub>2</sub>O<sub>2</sub>, Zn concentrations in diets, plasma and tissues were measured by atomic absorption spectrometry (AA-6600F; Shimadzu, Kyoto, Japan). The analytical accuracy of the Zn determination was confirmed by analysis of a certified reference material from bovine liver (standard reference material 1577b, National Institute of Standards and Technology, Gaithersburg, MD, USA).

### RNA extraction and quantitative RT-PCR

Total RNA was isolated from the small intestine epithelium, pancreas and liver using TRIzol reagent (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's protocol. Recovered RNA was used as a template for RT using random primers (ABI high-capacity complementary DNA reverse transcription kit; Applied Biosystems, Foster City, CA, USA). The quantitative RT-PCR was carried out using an SYBR premix Ex Taq II kit (TaKaRa, Otsu, Japan) in a Roter-Gene 6000 instrument (Corbett Research, Mortlake, Australia). PCR was performed as follows: an initial denaturation step of 10 s at 95°C, followed by forty cycles of 5 s at 95°C and 20 s at 60°C. The dissociation (melting) curve of quantitative RT-PCR products was subsequently examined by changing the ramp temperature from 60°C to 94°C. Each sample showed a single peak, suggesting that the expected PCR products were obtained. The PCR primers used to detect *Zip4*, *Zip5*, *ZnT1*, metallothionein-1a (*Mt-1a*), *Mt-2a*, insulin-like growth factor-1 (*Igf-1*) and hypoxanthine phosphoribosyltransferase 1 (*Hprt1*) were as follows: 5'-AAC CCA CCA GGG AGG AGA-3' and 5'-TTC TGG AAA CCC CTG CTT C'-3' for *Zip4*; 5'-CCT CGG GCC TAG ACC TCT T-3' and 5'-AGC TGG GAA CCA TTC AGA CA-3' for *Zip5*; 5'-AAC ACC AGC AAT TCC AAC G-3' and 5'-CCA CTG


**Table 1.** Ingredients of the basal diet

	g/kg diet
Glucose	634.96
Egg-white powder	200
Maize oil	100
Cellulose powder	20
Vitamin mixture*	10
D-Biotin	0.01
Mineral mixture†	35
ZnO (77.3% Zn)	0.03

\* The vitamin mixture (g/kg) contains: nicotinic acid, 3.000; Ca pantothenate, 1.600; pyridoxine-HCl, 0.0700; thiamine-HCl, 0.600; riboflavin, 0.600; folic acid, 0.20; D-biotin, 0.020; vitamin B<sub>12</sub> (cyanocobalamin) (0.1% in mannitol), 2.500; vitamin E (all-*rac*- $\alpha$ -tocopheryl acetate) (500 mg/g), 15.00; vitamin A (all-*trans*-retinyl palmitate) (150 mg/g), 0.800; vitamin D<sub>3</sub> (cholecalciferol) (10 mg/g), 0.250; vitamin K (phyloquinone), 0.075; powdered sucrose, 974.655.

† The mineral mixture (g/kg) contains: CaCO<sub>3</sub>, 357.00; KH<sub>2</sub>PO<sub>4</sub>, 196.00; K<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).H<sub>2</sub>O, 70.78; NaCl, 74.00; K<sub>2</sub>SO<sub>4</sub>, 46.60; MgO, 24.00; Fe(C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>), 6.06; MnCO<sub>3</sub>, 0.63; CuCO<sub>3</sub>, 0.30; KIO<sub>3</sub>, 0.01; Na<sub>2</sub>O<sub>3</sub>Se, 0.01 025; (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O, 0.00 795; NaSiO<sub>2</sub>.9H<sub>2</sub>O, 1.45; CrK(SO<sub>4</sub>)<sub>2</sub>.12H<sub>2</sub>O, 0.275; LiCl, 0.174; H<sub>3</sub>BO<sub>3</sub>, 0.0815; NaF, 0.0635; NiCO<sub>3</sub>, 0.0318; NH<sub>4</sub>VO<sub>3</sub>, 0.0066; powdered sucrose 222.676.





**Fig. 1.** Time-course changes in body weight (BW) in growing rats. Rats were fed diets containing various concentrations of Zn for 10 d. BW was plotted against experimental days. Values are means with their standard errors represented by vertical bars ( $n$  7). —○—, 24 mg Zn/kg; —△—, 1016 mg Zn/kg; —◆—, 2008 mg Zn/kg; —■—, 3000 mg Zn/kg.

GAT CAT CAC TTC TCA A-3' for *ZnT1*; 5'-CAC CAG ATC TCG GAA TGG AC-3' and 5'-GCA GCA GCT CTT CTT GCA G-3' for *Mt-1a*; 5'-ACC TCC TGC AAG AAA AGC TG-3' and 5'-ACT TGT CCG AAG CCT CTT TG-3' for *Mt-2a*; 5'-GGA CGC TCT TCA ATT CGT GT-3' and 5'-CTT CAG CGG AGC ACA GTA CA-3' for *Igf-1*; and 5'-GAC CGG TTC TGT CAT GTC G-3' and 5'-ACC TGG TTC ATC ATC ACT AAT CAC-3' for *Hprt1*. Gene transcript levels in each sample were determined using the relative standard curve method. The level of gene transcripts was expressed as a ratio relative to *Hprt1* mRNA, with the level in rats fed the basal diet set to 1.

**Statistical analyses**

Data are expressed as the least square means with their standard errors. All analyses were performed using SAS (SAS Institute, Cary, NC, USA)<sup>(29)</sup>. The data on BW and feed


intake were subjected to the MIXED procedure. Each rat was determined an experimental unit and measurements of the same rat on different days were considered repeated measures. The statistical model included the effects of diet, experimental day and the interaction between both. In addition, the effects of dietary Zn on tissue weight were analysed with the general linear model procedure. Furthermore, when tissue Zn concentrations or weights were plotted against daily Zn intake, a breaking point of daily Zn intake indicating a plateau was explored using the NLIN procedure. A model with 1 breaking point and no limit of its slope value before or after the point was applied. When the model was significant, and when the slope was not significantly different from 0, the breaking point was further determined by application of the model with 1 breaking point with 0 as the slope after the point. Differences were considered significant at  $P < 0.05$ .

**Results**

*Body and tissue growth*

Time-course changes in BW showed an insignificant effect of diet, but the effect of the interaction between diet and experimental day was significant, suggesting that the effect of excess Zn on BW depended on the length of the treatment period (Fig. 1). This indicates that, unlike in piglets, a diet supplemented with 3000 mg Zn/kg does not have a beneficial effect on BW gain in rats. As for daily feed intake, both the diet effect and the interaction between diet and experimental day were not statistically significant (Fig. 2(A)). In addition, feed efficiency, i.e. weight gain per feed intake, was not significantly affected by the diet, the experimental day or the interaction between both (Fig. 2(B)).

Typical symptoms of Zn toxicity are vomiting and gastrointestinal dysfunction<sup>(30)</sup>, and the pancreas is the organ most sensitive to Zn toxicity<sup>(6)</sup>. Two rats fed a diet supplemented with 3000 mg Zn/kg excreted soft stools for the last 3 d of the experimental period, although the other rats did not exhibit any symptoms throughout the study. The weight of the pancreas relative to BW was lower in rats fed a diet



**Fig. 2.** Time-course changes in daily feed intake and feed efficiency in growing rats. Rats were fed diets containing various concentrations of Zn for 10 d. (A) Daily feed intake and (B) feed efficiency were plotted against experimental days. Values are means with their standard errors represented by vertical bars ( $n$  7). —○—, 24 mg Zn/kg; —△—, 1016 mg Zn/kg; —◆—, 2008 mg Zn/kg; —■—, 3000 mg Zn/kg.

**Table 2.** Relative tissue weight of rats fed the diets supplemented with excess zinc (Mean values with their standard errors, *n* 7)

Dietary Zn (mg/kg diet)...	24	1016	2008	3000	SEM	<i>P</i>
Tissue weight (mg/g BW)						
Liver	45.9	45.0	47.8	48.5	1.4	NS
Kidney*	4.57	4.51	4.86	4.77	0.15	NS
Pancreas	6.19 <sup>a</sup>	6.57 <sup>a</sup>	5.18 <sup>b</sup>	5.19 <sup>b</sup>	0.28	<0.003
Spleen	2.51	2.47	2.42	2.80	0.10	NS
Testis*	5.16	5.30	5.56	5.42	0.23	NS
Gastrocnemius muscle*	0.98	0.96	0.95	0.90	0.04	NS
Femur*	3.81	3.75	3.99	3.74	0.09	NS
Perirenal fat*	4.93	4.24	3.50	3.43	0.51	NS

<sup>a,b</sup> Mean values with unlike letters were significantly different ( $P < 0.05$ ).

\* Average weight of tissues from both left and right parts.

supplemented with 2008 mg Zn/kg or 3000 mg Zn/kg compared to those fed a diet supplemented with 24 mg Zn/kg or 1016 mg Zn/kg, whereas no significant effects of the diet on the weight of the liver, kidney, spleen, testis, gastrocnemius muscle, femur and perirenal fat were detected (Table 2). Plotting the pancreas weight against daily Zn intake revealed a decrease in rats that ingested Zn at levels above 15.2 mg/d (Fig. 3).

#### Plasma and tissue zinc concentration


Plasma and tissue Zn concentrations were higher in rats fed diets supplemented with more than 1016 mg Zn/kg than in those fed the basal diet (data not shown). To examine the relationship between Zn intake and the Zn concentrations in detail, plasma and tissue Zn concentrations were plotted against daily Zn intake (Fig. 4). Zn concentrations in the plasma, liver, kidney and femur increased linearly with increasing intake of Zn up to 31.3, 37.1, 28.3 and 35.1 mg/d, respectively, and eventually reached a plateau (Fig. 4(A)–(D)). These results suggest that Zn efficiently accumulates in the plasma, liver, kidney and femur in response to increased Zn intake and that the capacity to retain Zn is limited. In view of the higher proportion of skeletal muscle weight relative to total BW, approximately 60% of body Zn is stored in the muscle tissue<sup>(31)</sup>; however, the Zn concentration in the muscle was not significantly altered by increasing Zn intake (Fig. 4(E)).

The relationship between the Zn concentration in the pancreas and daily Zn intake also indicated that the breaking point was 8.4 mg/d; the pancreatic Zn concentration increased up to this point (Fig. 4(F)). In contrast to the plasma, liver, kidney and femur, the Zn concentration in the pancreas linearly decreased in rats that ingested Zn above the breaking point.


#### Expression of zinc transporters, metallothionein and insulin-like growth factor-1

The gene transcript levels of the transporters involved in intestinal absorption and secretion of Zn were examined by quantitative RT-PCR (Fig. 5). Expression of *Zip4* in the intestinal epithelium, which is responsible for the uptake of Zn across


the mucosal membrane<sup>(24)</sup>, was significantly lower in rats fed a diet supplemented with 1016 mg Zn/kg than in those fed the basal diet, with no differences among dietary groups with higher Zn contents (Fig. 5(A)). The expression of *ZnT1* in the intestinal epithelium, which promotes Zn transport from the cytosol of epithelial cells to the portal vein<sup>(23)</sup>, was also down-regulated in rats fed diets supplemented with 1016–3000 mg Zn/kg (Fig. 5(B)). *Zip5* is located at the basolateral membrane of the intestinal epithelium and promotes Zn transport from the portal vein to epithelial cells<sup>(25)</sup>. The expression of *Zip5* was down-regulated in rats fed diets supplemented with 1016–3000 mg Zn/kg; the expression in rats fed a diet supplemented with 2008 mg Zn/kg was higher than that in rats fed a diet containing 1016 mg Zn/kg (Fig. 5(C)). No significant differences were detected in the expression of *ZnT1* and *Zip5*, transporters responsible for Zn secretion into the gut<sup>(25,26)</sup>, in the pancreas (Fig. 5(D) and (E)).




**Fig. 3.** Effects of excess Zn intake on the weight of the pancreas in growing rats. Rats were fed diets with various concentrations of Zn for 10 d. The weight of the pancreas relative to body weight (BW) was plotted against average daily intake of Zn. The breaking point of daily Zn intake on the pancreas weight was calculated, and shown in the figure by an arrow. O, 24 mg Zn/kg;  $\Delta$ , 1016 mg Zn/kg;  $\blacklozenge$ , 2008 mg Zn/kg;  $\blacksquare$ , 3000 mg Zn/kg.



**Fig. 4.** Relationship between Zn intake and plasma and tissue concentrations of Zn in growing rats. Rats were fed diets with various concentrations of Zn for 10 d. Zn concentrations in the (A) plasma, (B) liver, (C) kidney, (D) femur, (E) gastrocnemius muscle and (F) pancreas were plotted against average daily intake of Zn. The breaking point of daily Zn intake on plasma and tissue concentrations of Zn was calculated, and indicated in the figure by an arrow. ○, 24 mg Zn/kg; △, 1016 mg Zn/kg; ◆, 2008 mg Zn/kg; ■, 3000 mg Zn/kg. Note that there was no break point on Zn concentration in the gastrocnemius muscle.



**Fig. 5.** Gene expression of Zn transporters in the small intestinal epithelium and the pancreas of rats. Rats were fed diets supplemented with various concentrations of Zn for 10 d. Gene expression of (A) *Zip4*, (B) *ZnT1* and (C) *Zip5* in the small intestine, and (D) *ZnT1* and (E) *Zip5* in the pancreas was examined by quantitative RT-PCR. The transcription levels were expressed as ratios to hypoxanthine phosphoribosyltransferase 1 with the level in rats fed the basal diet set to 1. Values are means with their standard errors represented by vertical bars ( $n$  7). <sup>a,b,c</sup> Mean values with unlike letters were significantly different ( $P < 0.05$ ).




**Fig. 6.** Gene expression of metallothionein (*Mt*) in the small intestinal epithelium, liver and pancreas of rats. Rats were fed diets supplemented with various concentrations of Zn for 10 d. Gene expression of (A, C, E) *Mt-1a* and (B, D, F) *Mt-2a* in the (A, B) small intestine, (C, D) liver and (E, F) pancreas was examined by quantitative RT-PCR. The transcription levels were expressed as ratios to hypoxanthine phosphoribosyltransferase 1 with the level in rats fed the basal diet set to 1. Values are means with their standard errors represented by vertical bars (*n* 7). <sup>a,b,c</sup> Mean values with unlike letters were significantly different (*P* < 0.05).

*Mt* is involved in Zn homeostasis, and *Mt* expression is induced by several metals including Zn<sup>(32,33)</sup>. Expression of *Mt-1a* has been shown to change in parallel with that of *ZnT1* in response to Zn exposure in cultured hepatoma cells and fibroblasts<sup>(34)</sup>; we therefore evaluated *Mt* expression in the present study. The expression levels of *Mt-1a* and *Mt-2a* in the intestinal epithelium were higher in rats fed diets supplemented with 2008 or 3000 mg Zn/kg than in those fed the basal diet (Fig. 6(A) and (B)); *Mt-1a* expression was higher in rats fed a diet supplemented with 2008 mg Zn/kg than in those fed a diet containing 3000 mg Zn/kg. The expression of *Mt-1a* and *Mt-2a* in the liver was higher in rats fed diets supplemented with 1016–3000 mg Zn/kg than in those fed the basal diet; the expression increased in a dose-dependent manner (Fig. 6(C) and (D)). The regulatory expression of hepatic *Mt-1a* and *Mt-2a* in response to excess Zn intake is similar to the changes in Zn accumulation in the liver (Fig. 4(B)); this reflects the fact that Zn accumulated in the liver is incorporated into *Mt-1a* and *Mt-2a*, which buffer excess Zn to provide protection from Zn toxicity<sup>(33)</sup>. No significant changes in *Mt* expression were detected in the pancreas (Fig. 6(E) and (F)).

A previous study suggested that the increased expression of *Igf-1* and its receptor are responsible for excess Zn-induced growth promotion in piglets<sup>(13)</sup>. We evaluated expression of *Igf-1* in the intestinal epithelium and in the liver, one of the major *Igf-1*-producing organs<sup>(35)</sup>. Expression of *Igf-1* was not significantly different among groups, irrespective of the tissues analysed (Fig. 7).

### Discussion

The present study examined the short-term effects of extremely high Zn intake on body and tissue weight; tissue Zn concentration; and the expression of Zn transporters, *Mt*, and *Igf-1* in growing rats. Our results revealed that, in contrast to studies in piglets, excess intake of Zn did not have any beneficial effects on growth, but rather induced a decrease in weight of the pancreas. The NRC<sup>(7)</sup> proposed that excess Zn predominantly accumulates in organs such as the liver, kidney and bone in order to protect the pancreas, the organ most sensitive



**Fig. 7.** Gene expression of insulin-like growth factor-1 (*Igf-1*) in the small intestinal epithelium and liver of rats. Rats were fed diets supplemented with various concentrations of Zn for 10 d. Gene expression of *Igf-1* in the (A) small intestine and (B) liver was examined by quantitative RT-PCR. The transcription levels were expressed as ratios to hypoxanthine phosphoribosyltransferase 1 with the level in rats fed the basal diet set to 1. Values are means with their standard errors represented by vertical bars (*n* 7).

to excess Zn. The present results basically support this model: Zn concentrations in the liver, kidney and femur proportionally increased with increasing Zn intake up to 28–37 mg/d. However, the weight of the pancreas was lower in rats that ingested more than 15 mg/d Zn. This indicates that the amount of Zn intake required to induce atrophy of the pancreas is smaller than that required to reach a plateau in Zn accumulation in other tissues. Thus, the present results obtained in rats ingesting extremely high concentrations of Zn for a short period suggest the imperfect buffering capacity of the liver, kidney and bone against excess Zn ingestion. Furthermore, our results suggest that Zn accumulation in the skeletal muscle, the tissue that stores the majority of Zn in the body, is not affected by dietary Zn intake.

Zn depletion-induced *Zip4* expression in the small intestine is well-characterised<sup>(36–38)</sup>. By contrast, less information is available regarding the expression of Zn transporters in animals fed excess Zn. Expression of *ZnT1* and *ZnT2* in the small intestine was significantly higher in rats fed a diet containing 180 mg Zn/kg for 2 weeks than in those fed a diet containing 30 mg Zn/kg<sup>(39)</sup>. The present study indicates that excess Zn ingestion clearly down-regulates the mRNA expression of the Zn transporters involved in intestinal Zn absorption, *Zip4* and *ZnT1*. The decrease in *Zip4* expression was particularly evident; the gene transcript level of *Zip4* in rats fed a diet supplemented with 1016 mg Zn/kg was only around 5% of that in rats fed the basal diet, whereas the *ZnT1* mRNA level of rats fed a diet containing 1016 mg Zn/kg was around 35% of that in rats fed the basal diet. These results suggest that in addition to a system detecting Zn depletion, intestinal cells also have a system to sense excess Zn, and they partly regulate Zn absorption through transcriptional inhibition of Zn transporters.

*ZnT1*, *Mt-1a* and *Mt-2a* are transcriptionally regulated by metal-responsive transcription factor-1, a Zn-sensing transcription factor<sup>(40)</sup>, and the expression of both genes increases in response to Zn exposure in cultured cells<sup>(34,41,42)</sup>; however, excess Zn ingestion (>2008 mg Zn/kg) caused the down-regulation of *ZnT1* mRNA expression in the intestinal epithelium but up-regulation of *Mt-1a* and *Mt-2a* mRNA expression. An unidentified additional regulatory mechanism of *ZnT1* expression is probably involved in rats fed a diet with extremely high Zn contents.

The capacity for Zn accumulation in the liver, kidney and bone in response to excess Zn ingestion was limited. Rats that ingested 28–37 mg Zn/d could not accumulate additional Zn in these tissues; this Zn level corresponded to ingestion of a diet supplemented with 2008 mg Zn/kg. These results suggest stimulation of Zn secretion by feeding diets containing more than 2008 mg Zn/kg, or inhibition of Zn absorption, or both. Although *Zip5* plays a role in Zn secretion across the intestinal mucosa<sup>(25)</sup>, the gene transcript level of *Zip5* in the intestinal epithelium was not elevated but instead decreased in rats fed a diet supplemented with 1016 mg Zn/kg compared to those fed the basal diet. Thus, it is unlikely that the *Zip5*-mediated Zn secretion is increased in rats fed diets with high Zn content. The pancreas is the major organ for endogenous Zn secretion into the gut<sup>(43)</sup>. Expression levels

of pancreatic *Zip5* and *ZnT1*, which are expressed predominantly in the acinar cells of the pancreas<sup>(25,26)</sup>, were not increased in response to the ingestion of diets supplemented with excess Zn. These results suggest that the increased Zn secretion from the pancreas is not responsible for the limited Zn accumulation in the liver, kidney and bone.

Intestinal Mt levels are inversely related to the rate of Zn absorption<sup>(44)</sup>, and it has been suggested that Mt inhibits Zn absorption<sup>(45)</sup>. Thus, the present finding that expression of *Mt-1a* and *Mt-2a* was up-regulated in the intestinal epithelium of rats fed diets supplemented with 2008 or 3000 mg Zn/kg may reflect an inhibition of intestinal Zn absorption in these rats. However, the expression of intestinal *Mt-1a* and *Mt-2a* was not significantly higher in rats fed the diet supplemented with 3000 mg Zn/kg than in those fed the diet containing 2008 mg Zn/kg. Thus, the limit of tissue Zn accumulation in rats fed the diet supplemented with 2008 mg Zn/kg could not be explained by the Mt-induced inhibition of Zn absorption. Although gene transcript levels of intestinal *Zip4* were not further decreased in rats fed diets containing more than 1016 mg Zn/kg in this study, processing and translocation of *Zip4* are modified in response to changes in Zn status<sup>(46,47)</sup>. Thus, post-translational modifications may be responsible for the defence against Zn ingestion in diets containing more than 2008 mg Zn/kg.

In piglets, short-term feeding of a diet supplemented with 3000 mg Zn/kg effectively enhanced BW gain<sup>(11–13)</sup>, and this feeding regimen is applied in practice on pig farms. Li *et al.*<sup>(13)</sup> suggested that the excess Zn ingestion stimulates Igf-1-mediated signalling, which enhances the villous height of the small intestinal mucosa resulting in growth promotion of piglets. Considering that *Igf-1* expression in the small intestine and in the liver was not significantly increased in response to excess Zn ingestion, the inability to stimulate the Igf-1 axis may be one of the reasons why body growth was not accelerated in rats fed a diet supplemented with 3000 mg Zn/kg.

The present study clarified that growing rats have defence mechanisms against excess Zn ingestion; in addition to the effective accumulation of excess Zn in the liver, kidney and bone, the down-regulated mRNA expression of Zn transporters involved in intestinal absorption contributes to the protection of the pancreas against excess Zn-mediated adverse effects. Furthermore, up-regulation of Mt expression in the small intestine results in the inhibition of Zn absorption. These multiple defences may contribute to the relative tolerance to excess Zn ingestion.

### Acknowledgements

The present study received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. T. F., T. M. and M. F. designed the experiments; T. F. performed the experiments; T. F., T. M. and M. F. analysed the data and wrote the paper. All authors discussed the results and approved the manuscript in its final version. The authors declare no conflict of interest.

## References

1. McCall KA, Huang C & Fierke CA (2000) Function and mechanism of zinc metalloenzymes. *J Nutr* **130**, 1437S–1446S.
2. Giugliano R & Millward DJ (1984) Growth and zinc homeostasis in the severely Zn-deficient rat. *Br J Nutr* **52**, 545–560.
3. Park JH, Grandjean CJ, Hart MH, *et al.* (1986) Effect of pure zinc deficiency on glucose tolerance and insulin and glucagon levels. *Am J Physiol* **251**, E273–E278.
4. Fairweather-Tait S & Hurrell RF (1996) Bioavailability of minerals and trace elements. *Nutr Res Rev* **9**, 295–324.
5. Sandstead HH, Frederickson CJ & Penland JG (2000) History of zinc as related to brain function. *J Nutr* **130**, 496S–502S.
6. Sutomo FX, Woutersen RA & Van den Hamer CJ (1992) Effects of elevated zinc intake on the copper metabolism and the pancreas of the mouse. *J Trace Elem Electrolytes Health Dis* **6**, 75–80.
7. National Research Council (2005) *Mineral Tolerance of Animals*, 2nd ed. Washington, DC: National Academic Press.
8. Jenkins KJ & Hidiroglou M (1991) Tolerance of the pruruminant calf for excess manganese or zinc in milk replacer. *J Dairy Sci* **74**, 1047–1053.
9. Henry PR, Littell RC & Ammerman CB (1997) Effect of high dietary zinc concentration and length of zinc feeding on feed intake and tissue zinc concentration in sheep. *Anim Feed Sci Technol* **66**, 237–245.
10. Ansari MS, Miller WJ, Neathery MW, *et al.* (1976) Zinc metabolism and homeostasis in rats fed a wide range of high dietary zinc levels. *Proc Soc Exp Biol Med* **152**, 192–194.
11. Hill GM, Mahan DC, Carter SD, *et al.* (2001) NCR-42 Committee on Swine Nutrition. Effect of pharmacological concentrations of zinc oxide with or without the inclusion of an antibacterial agent on nursery pig performance. *J Anim Sci* **79**, 934–941.
12. Mavromichalis I, Peter CM, Parr TM, *et al.* (2000) Growth-promoting efficacy in young pigs of two sources of zinc oxide having either a high or a low bioavailability of zinc. *J Anim Sci* **78**, 2896–2902.
13. Li X, Yin J, Li D, *et al.* (2006) Dietary supplementation with zinc oxide increases Igf-I and Igf-I receptor gene expression in the small intestine of weanling piglets. *J Nutr* **136**, 1786–1791.
14. National Research Council (1998) *Nutrient Requirements of Swine*, 10th ed. Washington, DC: National Academic Press.
15. Hill GM, Mahan DC, Carter SD, *et al.* (2001) Effect of pharmacological concentrations of zinc oxide with or without the inclusion of an antibacterial agent on nursery pig performance. *J Anim Sci* **79**, 934–941.
16. Poulsen HD (1995) Zinc oxide for weaning piglets. *Acta Agric Scand A Anim Sci* **45**, 159–167.
17. Jensen-Waern M, Melin L, Lindberg R, *et al.* (1998) Dietary zinc oxide in weaned pigs – effects on performance, tissue concentrations, morphology, neutrophil functions and faecal microflora. *Res Vet Sci* **64**, 225–231.
18. Katouli M, Melin L, Jensen-Waern M, *et al.* (1999) The effect of zinc oxide supplementation on the stability of the intestinal flora with special reference to composition of coliforms in weaned pigs. *J Appl Microbiol* **87**, 564–573.
19. Guerinot ML (2000) The ZIP family of metal transporters. *Biochim Biophys Acta* **1465**, 190–198.
20. Taylor KM & Nicholson RI (2003) The LZT proteins; the LIV-1 subfamily of zinc transporters. *Biochim Biophys Acta* **1611**, 16–30.
21. Kambe T, Yamaguchi-Iwai Y, Sasaki R, *et al.* (2004) Overview of mammalian zinc transporters. *Cell Mol Life Sci* **61**, 49–68.
22. Palmiter RD & Huang L (2004) Efflux and compartmentalization of zinc by members of the SLC30 family of solute carriers. *Pflugers Arch* **447**, 744–751.
23. McMahon RJ & Cousins RJ (1998) Regulation of the zinc transporter ZnT-1 by dietary zinc. *Proc Natl Acad Sci U S A* **95**, 4841–4846.
24. Wang K, Zhou B, Kuo YM, *et al.* (2002) A novel member of a zinc transporter family is defective in acrodermatitis enteropathica. *Am J Hum Genet* **71**, 66–73.
25. Dufner-Beattie J, Kuo YM, Gitschier J, *et al.* (2004) The adaptive response to dietary zinc in mice involves the differential cellular localization and zinc regulation of the zinc transporters ZIP4 and ZIP5. *J Biol Chem* **279**, 49082–49090.
26. Liuzzi JP, Bobo JA, Lichten LA, *et al.* (2004) Responsive transporter genes within the murine intestinal-pancreatic axis form a basis of zinc homeostasis. *Proc Natl Acad Sci U S A* **101**, 14355–14360.
27. National Research Council (1995) *Nutrient Requirements of Laboratory Animals*, 4th ed. Washington, DC: National Academic Press.
28. Reeves PG (1997) Components of the AIN-93 diets as improvements in the AIN-76A diet. *J Nutr* **127**, 838S–841S.
29. SAS Institute (2001) *SAS User's Guide: Statistics (Ver. 9-2)*. Cary, NC: SAS Institute.
30. Fosmire GJ (1990) Zinc toxicity. *Am J Clin Nutr* **51**, 225–227.
31. Jackson MJ (1989) Physiology of zinc: general aspects. In *Zinc in Human Biology*, pp. 1–14 [CF Mills, editor]. Berlin: Springer-Verlag.
32. Palmiter RD (1994) Regulation of metallothionein genes by heavy metals appears to be mediated by a zinc-sensitive inhibitor that interacts with a constitutively active transcription factor, MTF-1. *Proc Natl Acad Sci U S A* **91**, 1219–1223.
33. Mocchegiani E, Malavolta M, Costarelli L, *et al.* (2010) Zinc, metallothioneins and immunosenescence. *Proc Nutr Soc* **69**, 290–299.
34. Langmade SJ, Ravindra R, Daniels PJ, *et al.* (2000) The transcription factor MTF-1 mediates metal regulation of the mouse ZnT1 gene. *J Biol Chem* **275**, 34803–34809.
35. Ohlsson C, Mohan S, Sjögren K, *et al.* (2009) The role of liver-derived insulin-like growth factor-I. *Endocr Rev* **30**, 494–535.
36. Dufner-Beattie J, Wang F, Kuo YM, *et al.* (2003) The acrodermatitis enteropathica gene ZIP4 encodes a tissue-specific, zinc-regulated zinc transporter in mice. *J Biol Chem* **278**, 33474–33481.
37. Weaver BP, Dufner-Beattie J, Kambe T, *et al.* (2007) Novel zinc-responsive post-transcriptional mechanisms reciprocally regulate expression of the mouse Slc39a4 and Slc39a5 zinc transporters (Zip4 and Zip5). *Biol Chem* **388**, 1301–1312.
38. Jou MY, Hall AG, Philipps AF, *et al.* (2009) Tissue-specific alterations in zinc transporter expression in intestine and liver reflect a threshold for homeostatic compensation during dietary zinc deficiency in weanling rats. *J Nutr* **139**, 835–841.
39. Liuzzi JP, Blanchard RK & Cousins RJ (2001) Differential regulation of zinc transporter 1, 2, and 4 mRNA expression by dietary zinc in rats. *J Nutr* **131**, 46–52.
40. Laity JH & Andrews GK (2007) Understanding the mechanisms of zinc-sensing by metal-response element binding transcription factor-1 (MTF-1). *Arch Biochem Biophys* **463**, 201–210.
41. Yang M, Kroft SH & Chitambar CR (2007) Gene expression analysis of gallium-resistant and gallium-sensitive lymphoma cells reveals a role for metal-responsive transcription factor-1, metallothionein-2A, and zinc transporter-1 in modulating the





- antineoplastic activity of gallium nitrate. *Mol Cancer Ther* **6**, 633–643.
42. Jackson KA, Valentine RA, McKay JA, *et al.* (2009) Analysis of differential gene-regulatory responses to zinc in human intestinal and placental cell lines. *Br J Nutr* **101**, 1474–1483.
  43. McClain CJ (1990) The pancreas and zinc homeostasis. *J Lab Clin Med* **116**, 275–276.
  44. Hoadley JE, Leinart AS & Cousins RJ (1988) Relationship of  $^{65}\text{Zn}$  absorption kinetics to intestinal metallothionein in rats: effects of zinc depletion and fasting. *J Nutr* **118**, 497–502.
  45. Davis SR, McMahon RJ & Cousins RJ (1998) Metallothionein knockout and transgenic mice exhibit altered intestinal processing of zinc with uniform zinc-dependent zinc transporter-1 expression. *J Nutr* **128**, 825–831.
  46. Kim BE, Wang F, Dufner-Beattie J, *et al.* (2004)  $\text{Zn}^{2+}$ -stimulated endocytosis of the mZIP4 zinc transporter regulates its location at the plasma membrane. *J Biol Chem* **279**, 4523–4530.
  47. Kambe T & Andrews GK (2009) Novel proteolytic processing of the ectodomain of the zinc transporter ZIP4 (SLC39A4) during zinc deficiency is inhibited by acrodermatitis enteropathica mutations. *Mol Cell Biol* **29**, 129–139.