
Index

- $\partial_x = x \cdot (\partial/\partial x)$, 245
 ∇ , divergence, 242
~, linear equivalence, xv
 $\sim_{\mathbb{Q}}$, \mathbb{Q} -linear equivalence, xv
 $\sim_{\mathbb{R}}$, \mathbb{R} -linear equivalence, 434
 \equiv , numerical equivalence, xv
 \equiv , identity of sequences or polynomials, 187
 $[\otimes]$, tensor product for divisorial sheaves, 129
 \dashrightarrow , rational map, xiv
 \rightarrow , morphism, xiv
 $\{ \}$, fractional part, xv
 $\lfloor \rfloor, \lceil \rceil$, rounding down or up, xv
 $\widehat{}$, completion, xvii, 397
 \leq , for sequences, 187
 \leq , for polynomials, 187
 $a(\ , \ , \)$, discrepancy of divisor, 420
 $a_\ell(\ , \ , \) := a(\ , \ , \) + 1$, log discrepancy, 420
 \mathbb{A}_k^n or \mathbb{A}_x^n , affine n -space, xiv
 $\mathbb{A}^n/m(a_1, \dots, a_n)$, quotient singularity, 27
two-dimensional, 248
Adjunction, 424
inversion of, 424
AEnv(), affine envelope, 435
AFI, Alexeev–Filipazzi–Inchiostro, 320
functor, $\mathcal{AFI}(\ , \)$, 321
Alexeev stable, 234
good moduli theory, 235
Ample
fiber-wise, 193
 \mathbb{R} -divisor, 439
relatively, xv
strongly, 326
Approximation of \mathbb{R} -divisors, 436, 437
Ass(), associated points or subschemes, 370
Asymptotic Riemann–Roch, 439
 $\text{Aut}_{\mathcal{S}}(\), \mathbf{Aut}_{\mathcal{S}}(\)$, xvii, 337
finite for stable families, 338
Base change
dualizing sheaf, 100, 102
K-flat, 281
notation, xvii
pluricanonical, 112
Bertini theorem
C-flatness, 290
divisorial support, 278
flatness, 400
generically Cartier family, 153
hulls, 377
husks, 355
inverse, local stability, 64, 184
K-flatness, 263
local stability, 64
relative hull, 351
 S_m , 377
Big, divisor, xvi
Birational
transform, xv
fiberwise, 11
map, xiv
Boundary, xvi
Boundedness, 16
marked pairs, 332

- set of sheaves, 360
 strong, 224
 weak, 224
- $C_a(X, L)$, affine cone, 81
 Calabi–Yau pair, xvi
 Canonical
 algebra, 427
 class, 17, 419
 class, relative, 61, 133
 divisor, 17
 line bundle, 17
 model, xvi, 23, 427
 model of resolutions, xvi
 model, existence, 428
 model, nef slc case, 318
 model, simult. numerical criterion, 186
 model, simult. of resolutions, 185
 model, simultaneous, 185
 modification, xvi, 428
 modification, simultaneous, 188
 ring, 23, 26
 ring, not finitely generated, 36
 sheaf, 17, 61
 sheaf, absolute, 419
 sheaf, relative, 419
 singularity, 24, 421
 surface singularity, list, 68
- Canonically
 embedded family, 333
 polarized family, 332
- Cartier
 divisor, 146
 divisor, relative, 150
 divisor, valuative criterion, 156
 generically \sim pull-back, 140, 154
 generically \sim , relative, 152
 index, xv
 non- \sim locus, 152
- Categorical
 moduli as \sim quotient, 334, 335
 quotient by group action, 334
- Cayley–Chow
 correspondence, 176
 correspondence, Mumford divisors, 176
 correspondence, over fields, 171
 family, 120
 form, 171
 hypersurface, 171, 284
 hypersurface, flag, Grassmann, incidence,
 product versions, 284, 285
- inverse, scheme-theoretic, 177
 type hypersurface, 172
 $C^m\mathcal{ESch}(\cdot)$, $C^m\mathcal{E}^s\mathcal{MSch}(\cdot)$, functor and
 moduli of canonically embedded,
 marked schemes, 333
- Center
 log \sim , 422
 log canonical \sim , 422
 of a divisor, xv
 $C^m\mathcal{ESP}(\cdot)$, $C^m\mathcal{ESP}(\cdot)$, functor and moduli of
 m -canonically embedded, stable pairs,
 222, 333, 337
- C-flat, 281
 Bertini theorem, 290
 locally \sim , 288
 stably \sim , 281
- $\text{Ch}(\cdot)$, Cayley–Chow hypersurface, 171
 Ch_{gr} , Ch_{in} , Ch_{fl} , Ch_{pr} , versions of
 Cayley–Chow hypersurface, 284
 $\text{Ch}_{sch}^{-1}(\cdot)$, Cayley–Chow inverse, 177
 Chow
 equations, ideal of, 173
 hull of Mumford divisor, 175
 hull, of a cycle, 174
 variety, 5
 CM, Cohen–Macaulay, xv, 372
 coeff(\cdot), set of coefficients, 418
 coeff $_E(\cdot)$, coefficient of E in, 418
- Coefficient
 of a prime divisor, 418
 vector of marking, 310
 floating, 317
 generic, 216
 major, 216
 standard, 216
 Cohen–Macaulay, CM, xv, 372
 Completion, $\widehat{\cdot}$, xvii, 397
 Component-wise dominant, 155
 Cone
 affine, 81
 deformation to, 39, 82
 Continuous choice, 221
 Contraction, xiv
 crepant, 318
 simultaneous, 323
 simultaneous, crepant, 323
 $\text{CoSp}(\cdot)$, \mathbb{Q} -vector space spanned by
 coefficients, 434
 $C^m\mathcal{P}^s\mathcal{MSch}(\cdot)$, functor of canonically
 polarized, marked schemes, 333
 Crepant, contraction, 318

- Curve
 stable, 10
 stable extension, 11
 Curvilinear scheme, 402
 Cycle
 degree of, 171
 effective, 170
 fundamental \sim , [], 170
 geometrically reduced, 171
 on a scheme, 170
 width of, 174
 Cyclic cover, 426
- $\Delta^{>c}, \Delta^=c, \Delta^{$, 418
 Δ_c or Δ_c^{div} , divisorial fiber, 108
 \mathbb{D} , unit disc, 32
 Decomposition
 locally closed, 415
 partial, 415
 Deformation, 237
 \mathbb{W}_{\sim} , 247
 \mathbb{Z}_{\sim} , 247
 hypersurface singularities, 76
 KSB- \sim , 247
 locally trivial, 237
 nonalgebraic, 31
 of quotients, 73
 Demi-normal, 430
 open condition, 390
 Depth, 371
 along a subscheme, 371
 and flatness, 374
 and push forward, 372
 of a sheaf, 371
 of slc scheme, 425
 semicontinuity, 380
 Descent, 338
 and functorial polarization, 339
 for flat, projective morphism, 340
 for rigid, projective morphism, 341
 Dévissage, 381
 Diff(), different, 423
 Different, 423
 properties of \sim , 424
 Differentiation, 238
 cohomological, 238
 Discrepancy
 log \sim , 420
 of a divisor, 420
 Discrete choice, 220
 Divergence, ∇ , 242
- Divisor
 big, xvi
 canonical, 17
 Cartier, 146
 Cartier, relative, 150
 Cartier, valuative criterion, 156
 generically \mathbb{Q} - or \mathbb{R} -Cartier, relative, 141
 generically Cartier, 146
 generically Cartier, relative, 152
 Mumford, xv, 146
 Mumford, relative, 175
 Mumford, universal family, 179
 on a scheme, xv
 over a scheme, xv
 \mathbb{Q}_{\sim} , 147
 \mathbb{R}_{\sim} , 147
 reduced, xv
 Weil, 147
- Divisorial
 fiber, Δ_c or Δ_c^{div} , 108
 log terminal, 421
 pull-back, 141
 restriction, D_t or D_t^{div} , xvii, 106
 sheaf, 129
 sheaf, flat family, 129
 sheaf, generically flat, 130
 sheaf, mostly flat, 130
 sheaf, valuative criterion, 155
 subscheme, 146
 subscheme, family of \sim , 149
 support and Fitting ideal, 275
 support, DSupp(), 274
 support, Bertini theorem, 278
 support, final definition, 276
 dlt, divisorial log terminal, 421
 is CM, 425
 DSupp(), divisorial support, 274
 Du Bois singularity, 97
 cohomology and base change, 98
 Du Val singularity, 68
 Dual graph, 67
 Dualizing
 sheaf, 17, 61
 sheaf, base change, 102
 sheaf, construction, 101
 sheaf, other deformations, 65
 sheaf, relative, 419
- Elementary étale, 66
 emb(), embedded subsheaf, 370
 Embedded point, 370

- Embedding, locally closed, 414
 $\mathcal{E}^s\mathcal{MSch}(\)$, $\mathcal{E}^s\mathcal{MSch}(\)$ functor and moduli of embedded, marked schemes, 331
Enough one-parameter families, 416
Envelope, affine or linear, 435
Etale, elementary, 66
 $\text{Ex}(\)$, exceptional set, xv
- F^{**} , reflexive hull, 348
 $F^{[**]}$, hull or S_2 -hull, 348
 F^H , relative hull, 350
Family
algebraic, 120
canonically embedded, 333
canonically polarized, 332
Cartier, normal base, 141
Cayley–Chow, 120
divisorial sheaves, flat, 129
divisorial sheaves, mostly flat, 130
divisorial subschemes, 149
generically Cartier divisors, 152
Hilbert–Grothendieck, 121
locally stable, 59, 118
marked pairs, 311
mostly flat of line bundles, 193
non-projective, 22
of polarized schemes, 327
one-parameter, 59
pairs, 59, 138
polarized with K-flat divisors, 330
stable, 89
stable over smooth base, 167
stable, extension of, 169
universal, 121
varieties, 58
well-defined, 140
well-defined, reduced base, 141
- Fano pair, xvi
Fiber, divisorial, Δ_c or Δ_c^{div} , 108
Fiber-wise ample, 193
Field of moduli, 52
hyperelliptic curve, 53
Fine moduli space, 7
universal family, 341
Fitting ideal, 273
Flat
family, divisorial sheaves, 129
generically, 130
mostly, 130
FlatCM($\$), flat and CM locus, 276
- Flatness
associated points, 398
Bertini theorem, 400
curvilinear fibers, 403
Hironaka’s theorem, 409
is open, 374
nodal fibers, 406
relative codimension ≥ 3 , 410
relative codimension 0, 199, 401
relative codimension 1, 405
relative codimension 2, 408
residue field extension, 400
with reduced fibers, 198
- Flattening decomposition, 127
Floating coefficient, 317
Formally K-flat, 281
Framing, projective, 329
Free group action, 335
Frobenius power, $I^{[q]}$, 173
Full subscheme, 294
Functorial polarization, 339
Fundamental cycle $[\]$, 170
- General type, xiv, 23
Generically
Cartier divisor, 146
flat, 130
flat and pure, 276
 \mathbb{Q} -or \mathbb{R} -Cartier divisor, relative, 141
- Genus, sectional, 123
Geometric quotient
by free group action, 337
by group action, 336
existence, 336
Geometrically injective, 414
Grothendieck–Lefschetz theorem, 116
- Henselisation, 66
strict, 67
Hilbert
function, of ω_X , 132
function, of divisorial sheaf, 132
functor, 6
 \sim -Grothendieck family, 121
scheme, 6
 \sim -to-Chow map, 124
 $\text{Hilb}(\)$, Hilbert scheme, 122
 Hilb^{str} , strongly embedded part of Hilb, 329
 $\text{Hom}_X(\ , \)$, $\mathcal{H}\text{om}_X(\ , \)$, $\text{Hom}_S(\ , \)$, xvii, 360
Homeomorphism, universal, 412

- $\mathcal{H}ull()$, Hull(), functor and moduli space of universal hulls, 363
- Hull, 130
- $S_2 \sim$, 348
 - algebraic spaces, 368
 - Bertini theorem, 351, 377
 - of a sheaf, 348
 - pull-back, 130
 - pure, 348
 - reflexive, 348
 - relative, 350
 - universal, 353
 - universal, characterization, 353
 - universal, fine moduli space, 363
- Hurwitz formula, 425
- $\mathcal{H}usk()$, Husk(), functor and moduli of husks, 357
- Husk, 355
- algebraic spaces, 368
 - Bertini theorem, 355
 - quotient, 355
 - quotient, relative, 356
 - relative, 356
 - tight, 355
- Hypersurface
- Cayley–Chow, 171, 284
 - Cayley–Chow, flag, Grassmann, incidence, product versions, 284, 285
- K-flatness, 295
- $I(,)$, intersection form, 321
- $I(,)$ intersection numbers, 187
- $I^{[q]}$, Frobenius power, 173
- Index
- Cartier \sim , xv
 - of ω_X , 247
 - of a variety, xv
- Intersection number, xv
- Inversion of adjunction, 424
- $\text{Isom}_{\mathbb{S}}(,)$, $\mathbf{Isom}_{\mathbb{S}}(,)$, xvii, 337
- finiteness of, 90, 338
- Isotrivial family, 52
- K_X , canonical class, 17, 419
- $K_{X/C}$, relative canonical class, 61
- $\kappa(X)$, Kodaira dimension, xiv, 23
- $\mathcal{K}\mathcal{D}iv()$, $\mathcal{K}\mathcal{D}iv()$, functor and moduli of K-flat divisors, 261
- K-flat, 259
- additive, 262
 - base change, 281
- Bertini theorem, 263
- equals stable C-flat, 283
- family, polarized, 330
- flat implies \sim , 262
- formal nature of, 283
- formally \sim , 281
- functor of \sim pull-backs, 293
- hypersurface singularities, 295
- implies C-flat, 282
- linear equivalence, 262
- locally \sim , 281
- multiplicative, 262
- over reduced base, 262
- push forward of, 262
- reasons for definition, 260
- seminormal curves, 301
- Kodaira dimension, xiv, 23
- jump of, 29–31
- Kodaira lemma, 440
- KSB, Kollar–Shepherd-Barron, 229
- good moduli theory, 230–232
 - stable, 229, 230
 - stable, major coefficients, 232
- KS_B-deformation, 247
- KSBA, Kollar–Shepherd-Barron–Alexeev
- good moduli theory, 308, 313, 315, 316
 - stable strong form, 316
 - stable, general coefficients, 315
 - stable, rational coefficients, 313
- $L^{[m]}$, reflexive power of divisorial sheaf, 129
- lc, log canonical, 421
- center, 422
- LEnv(), linear envelope, 435
- LEnv $_{\mathbb{Z}}$ (), integral points of LEnv, 314
- Lexicographic order, \leq , 187
- Lie derivative, 239
- Linear
- equivalence, \sim , xv
 - $\mathbb{Q}\sim$ equivalence, $\sim_{\mathbb{Q}}$, xv
 - $\mathbb{R}\sim$ equivalence, $\sim_{\mathbb{R}}$, 434
 - system, 4
- Link, 158
- Local
- morphism, 155
 - numerical criterion of \sim stability, 184
 - Picard group, 116
 - stability, Bertini, 64
 - stability, reduced base, 118
 - stability, representable, reduced base, 143

- Locally
 - C-flat, 288
 - closed decomposition, 415
 - closed embedding, 414
 - closed partial decomposition, 415
 - K-flat, 281
 - stable, 28
 - stable morphism, 59, 118
 - stable pair, 58
 - stable, equivalent conditions for, 142
 - stable, KSB version, 135
 - stable, reduced base, 118
- Locus
 - flat or flat-CM, 276
 - non-Cartier, 152
- Log big, 429
- Log canonical, 27
 - lc, 421
 - center, 422
 - modification, 428
- Log center, 422
 - mld bounds, 422
- Log discrepancy, 420
- Log resolution, xvi
- Major coefficient, 216
- Map, rational, xiv
 - birational, xiv
- Marked
 - family of ~ pairs, 311
 - pair, with divisors, 310
 - reasons for, 309
- Marking, of pair or family, 310, 311
- Matsusaka inequality, 440
- $\mathcal{MDiv}(\cdot)$, $\text{MDiv}(\cdot)$, functor and moduli of
 - Mumford divisors, 175
- Minimal log discrepancy, mld, 421
- mld, minimal log discrepancy, 421
- Model, canonical, xvi, 23
- Modification
 - canonical, xvi, 428
 - finite, 410
 - log canonical, 428
 - semi-log-canonical, slc, 211
 - simultaneous, canonical, 188
- Moduli
 - boundary, 217
 - embedded pairs, 221
 - embedded varieties, 221
 - enough one-parameter families, 416
 - field of, 52
- interior, 217
- representable, 13
- separated, 15
- Moduli space
 - categorical, 8
 - categorical quotient, 334, 335
 - coarse, 8
 - fine, 7
 - fine for universal hull, 363
 - genus 2 curves, 45
 - husks, 357
 - hypersurfaces, 38, 40
 - irreducible components proper, 169
 - KSBA, exists, 308
 - non-separated, 18, 21, 41, 48
 - projectivity of, 225
 - quotient by group action, 333
 - quotient husks, 357
 - reduced version, 138
 - stable varieties, 120
- Moduli theory
 - Alexeev, 235
 - good, 226
 - KSBA, 230
 - KSBA, standard coeffs, 231
 - KSBA, rational coeffs, 313
 - KSBA, real coeffs, 315
 - KSBA, strong form, 316
 - V^+ , 234
- Monomorphism, 414
- $\text{Mor}(\cdot, \cdot)$, $\text{Mor}_S(\cdot, \cdot)$, xvii, 337
- Morphism, xiv
 - dominant, 155
 - Hilbert-to-Chow, 124
 - locally stable, 59
 - locally stable, reduced base, 118
 - pure dimensional, 104
 - scheme, Mor , 337
 - small, xv
 - stable, 13
- Morse lemma, 390
- Mostly flat, 130
 - divisorial sheaf, 130
 - family of line bundles, 193
 - S_2 sheaf, 130
- Mumford divisor, xv, 146
 - along subscheme, 146
 - Cayley–Chow correspondence, 176
 - Chow hull of, 175
 - flat, 227
 - functor and moduli, 175

- relative, 175
- relative class group, 265
- universal family, 179

- Nagata openness criterion, 375
- Nakai–Moishezon criterion, 441
- Nef, xv
- Negativity lemma, 444
- Node, 430
 - deformation of, 407
- Noether normalization
 - étale version, 395
 - fails for affine morphism, 394
 - local version, 396
- Norm, 117
- Normal, 411
 - crossing, simple, xvi
 - pair, 411
- Numerical
 - criterion for relative line bundles, 184
 - criterion of local stability, 184
 - criterion of stability, 181, 183
 - criterion of simult. canonical model, 186
 - equivalence, \equiv , xv
 - pull-back, 442, 443
- Numerically
 - log canonical, 163
 - polarized, 328
 - \mathbb{Q} -Cartier, 163
 - \mathbb{R} -Cartier, 163
 - \mathbb{R} -Cartier, lc modification of, 428
 - relatively trivial, 163
 - semi-log-canonical, 163
 - slc is slc, 164
- $\omega_{X/S}$, relative dualizing sheaf, 419
- ω_X , canonical or dualizing sheaf, 17
- $\omega_X^{[m]}$, reflexive power of canonical sheaf, 18
- Obstruction theory, 218

- \mathbb{P}_k^n or \mathbb{P}_x^n , projective n -space, xiv
- Pair, xvi, 418
 - Calabi–Yau, xvi
 - family of, 59, 138
 - family, marked, 311
 - Fano, xvi
 - locally stable, 58
 - marked with divisors, 310
 - normal, 411
 - rigid and universal family, 341
 - seminormal, 411
- stable, 58
- weakly normal, 411
- well-defined family, 140
- Partial decomposition, 415
- PGL, group scheme, 333
- $\text{Pic}(\cdot)$, $\mathbf{Pic}(\cdot)$, Picard group and scheme, xvii
- $\text{Pic}^{\text{loc}}(\cdot)$, $\mathbf{Pic}^{\text{loc}}(\cdot)$, local Picard group and scheme, 116, 164
- Picard group, xvii
 - for smooth morphisms, 159
 - local, 116, 164
- $\mathcal{P}^s\mathcal{M}\mathcal{S}ch(\cdot)$, functor of polarized, marked schemes, 331
 - stack version, 332
- Poincaré residue map, 423
- Pointed scheme, 330
- Polarization
 - family of schemes, 327
 - functorial, 339
 - numerical, 328
 - scheme, 327
 - strong, 327
- Potentially, slc or ..., 99
- Pre-polarization, 327
- Preserve residue fields, 414
- Projection
 - approximation of, 279
 - various versions, 280
- Projective
 - framing, 329
 - moduli space, 225
- Proper
 - group action, 335
 - valuative \sim , 15
- $\mathcal{P}^s\mathcal{S}ch(\cdot)$, functor of polarized schemes, 327
 - étale sheafification of, 328
 - stack version, 330
- Pull-back
 - C-flat, 294
 - Cartier, 154
 - divisorial, 141
 - generically Cartier, 140
 - hull \sim , 130
 - K-flat, 293
 - locally stable, representable, 160
 - numerical, 442, 443
 - \mathbb{Q} - or \mathbb{R} -Cartier divisors, 141
 - stable, representable, 160
 - Weil-divisor, 139
- Pure
 - quotient, 147, 371

- relatively, 365
- scheme, 371
- sheaf, 370
- vertically, 352
- Purely log terminal, plt, 421
- $q^{[*]}(D)$, pull-back (generically Cartier), 140
- $q_{W\text{div}}^*(D)$, Weil-divisor pull-back, 139
- \mathbb{Q} -Cartier divisor, xv
 - valuative criterion, 156
- \mathbb{Q} -divisor, xv, 147
- $Q\mathcal{Husk}(\cdot)$, $Q\text{Husk}(\cdot)$, functor and moduli of quotient husks, 357
- Quasi-étale, 62
- $Quot(\cdot)$, $Quot(\cdot)$, functor and scheme, 359
- Quot-scheme, 359
- Quotient
 - categorical, by group action, 334
 - geometric, by free group action, 337
 - geometric, by group action, 336
 - geometric, existence, 336
 - husk, 355
 - singularity, 27
- $R(\cdot)$, canonical ring, 23, 26
- \mathcal{R} , restriction or Poincaré residue map, 423
- Rational
 - double point, 68
 - map, xiv
 - singularity, xv
- \mathbb{R} -Cartier divisor, 434
 - class, 440
 - valuative criterion, 156
- \mathbb{R} -divisor, xv, 147, 434
 - ample, 439
 - convex approximation, 436, 437
 - depth of pluricanonical sheaf, 438
- \mathbb{R} -line bundle, 440
- Reduced
 - normal form, 230
 - divisor, xv
- Regular sequence, 371
- Relatively
 - ample, xv
 - generically Cartier, 141
 - isomorphic, 364
 - pure, 365
- Representable
 - C-flat pull-back, 294
 - Cartier pull-back, 154
 - flat, divisorial pull-back, 154
- flatness, 127
- functor, 126
- hull of divisorial sheaves, 131
- invertible hull of sheaves, 131
- K-flatness, 261, 293
- local stability, 119, 160
- local stability, reduced base, 143
- moduli theory, 13
- pull-backs, 126
- stability, 119, 160
- stability, over reduced base, 143
- Residue map, 423
- Resolution, xv
 - dual graph of \sim , 67
 - $\log \sim$, xvi
 - of Du Val singularities, 68
- Restriction
 - divisorial, D_t or D_t^{div} , xvii, 106
 - map, 107, 353
- Riemann-Roch, asymptotic, 439
- Rigid, scheme or pair, 337
 - and universal family, 341
- Ring, canonical, 23, 26
- S_2 , Serre's condition, 371
 - divisorial sheaf, 129
 - family of varieties, 118
 - for families, 107
 - for restriction, 106
 - Hilbert-to-Chow map, 124
 - mostly flat, 130
- Semi-log-canonical, slc, 28, 431
 - and depth, 425
 - modification, 211
- Seminormal, 411
 - K-flatness for \sim curves, 301
 - pair, 411
- Separatedness
 - for husks, 358
 - for stable maps, 90, 338
 - moduli spaces, 15
 - valuative criterion, 15
- Serre's condition S_m , 371
 - along a subset, 105
- Seshadri criterion, 441
- Sheaf
 - canonical, 17
 - divisorial, 129
 - dualizing, 17
- Simple normal crossing, snc, xvi

- Singularity
 canonical, 24, 421
 cyclic quotient $\mathbb{A}^2/\frac{1}{n}(1, q)$, 27, 248
 dlt, 421
 Du Bois, 97
 Du Val, 68
 klt, 421
 lc, log canonical, 27, 421
 log terminal, 421
 plt, 421
 potentially slc or ..., 99
 quotient, 27
 slc, semi-log-canonical, 28
 terminal, 421
 slc, semi-log-canonical, 28, 431
 characterization using normalization, 431
 depth, 425
 potentially \sim , 99
 S_m , Serre's condition, 371
 along a subset, 105
 Bertini theorem, 377
 is open, 374
 Small morphism, xv
 snc, simple normal crossing, xvi
 $\mathcal{SP}(\cdot)$, $\mathcal{SP}(\cdot)$, functor and moduli of stable,
 marked pairs, 137, 223, 308
 $\mathcal{SP}^{\text{rigid}}(\cdot)$, $\mathcal{SP}^{\text{rigid}}(\cdot)$, functor and moduli of
 rigid pairs, 341
 SSupp, scheme-theoretic support, 274
 Stability
 automatic in codimension ≥ 3 , 62, 115, 184
 local, representable, 119
 numerical criterion, 181, 183
 representable, 119
 representable over reduced base, 143
 Stabilization functor, 34, 35
 over nodal curves, 36
 Stable
 Alexeev \sim , 234
 Alexeev–Filipazzi–Inchiostro, 320
 curve, 10
 equivalent conditions for, 142
 extension for curves, 11
 extension, weak, 96
 family over smooth base, 167
 family, 1 parameter, 89
 family, extension of, 169
 KSB, 135, 229, 230
 KSB, major coefficients, 232
 KSBA, general coefficients, 315
 KSBA, rational coefficients, 313
 KSBA, strong form, 316
 locally \sim , 28
 morphism, 13
 one parameter family, 59
 pair, 58
 $V^+ \sim$, 233
 variety, 28
 Stably C-flat, 281
 equals K-flat, 283
 independence of embedding, 293
 Standard coefficient, 216
 Stratum, of an snc pair, 422
 Strongly ample, 326
 Subscheme, divisorial, 146
 Support
 divisorial, 274
 divisorial, final definition, 276
 scheme-theoretic, 274
 $\mathcal{SV}(\cdot)$, $\mathcal{SV}(\cdot)$, functor and moduli of stable
 varieties, 120
 Tight husk, 355
 tors(\cdot), torsion subsheaf, 370
 Trace map, 101
 Transform, birational, xv
 Tree, 67
 Twig, 67
 Univ(\cdot), universal family, 121
 Universal
 family, 121
 family for rigid pairs, 341
 family of flat Mumford divisors, 227
 homeomorphism, 412
 hull, 353
 hull, characterization, 353
 Universally flat, 353
 V^+ , strict Viehweg stability, 233
 V-deformation, 247
 Valuative criterion
 for \mathbb{Q} -Cartier divisor, 156
 for flat, divisorial sheaf, 155
 for relative Cartier divisor, 156
 locally closed embedding, 415
 morphism, 413
 section, 413
 Valuative-proper, 15
 stable map, 90
 Vanishing theorem
 Ambro–Fujino version, 429
 Fujita version, 440

- Variety, xiv
general type, xiv, 23
stable, 28
Vertically pure, $\text{vpure}()$, 352
Volume, 16, 385
and push-forward, 385
birational models, 387, 388
finite maps, 388
perturbations, 389
 $\text{vpure}()$, vertically pure, 352
- W-deformation, 247
- Weakly normal, 411
pair, 411
Weil divisor, 147
pull-back, 139
relative, 148
Well-defined
family, 140
family, reduced base, 141
Width of a cycle, 174
- \mathbb{Z} -divisor, xv

