

patients in a tertiary-care teaching hospital. A computer screen displayed, at the time of physician order entry, an adaptation of the CDC guidelines for appropriate vancomycin use. The main outcome measures were the frequency of initiation and renewal of vancomycin therapy, as well as the duration of therapy prescribed on a per prescriber basis.

Compared with the control group, intervention physicians wrote 32% fewer orders (11.3 vs 16.7 orders per physician; $P=.04$) and had 28% fewer patients for whom they either initiated or renewed an order for vancomycin (7.4 vs 10.3 orders per physician; $P=.02$). In addition, the duration of vancomycin therapy attributable to physicians in the intervention group was 36% less than that of control physicians (26.5 vs 41.2 days; $P=.05$). Analysis of pharmacy data confirmed a decrease in the overall hospital use of intravenous vancomycin during the study period.

The authors concluded that implementation of computer-guided physician order entry decreased vancomycin use. Computerized guidelines represent a promising tool for changing prescribing practices.

FROM: Shojania KG, Yokoe D, Platt R, Fiskio J, Ma'luf N, Bates DW. Reducing vancomycin use utilizing a computer guideline: results of a randomized controlled trial. *J Am Med Assoc* 1998;279:554-562.

MRSA Bloodstream Infection and Mortality

Moreira and coinvestigators from the Universidade Federal de São Paulo, Brazil, have reported a study to identify the attributed mortality rate of bloodstream hospital infection by methicillin-resistant *Staphylococcus aureus* (MRSA) and its effect on length of hospital stay. In a case-control study conducted in a 660-bed, tertiary-care teaching hospital in São Paulo, Brazil, 71 adult patients with hospital-acquired MRSA bacteremia diagnosed between January 1, 1991, and September 30, 1992, and 71 MRSA-free controls were matched on age, gender, underlying disease, surgical procedure, same risk time, and admission date.

MRSA accounted for 73% of hospital bloodstream infections involving *Staphylococcus aureus*. Mortality rates were 56% (40/71) for cases and 11% (8/71) for controls. The attributable mortality rate was 45% (OR=17.0; CI₉₅, 3.58-202.26; $P=.000001$). The median length of hospital stay was 32.6 days for the cases and 29.8 for the controls ($P=.32$).

The authors concluded that a high proportion of

Staphylococcus aureus bacteremia involved MRSA. Nosocomial bloodstream infection with MRSA provides a high level of mortality independently from the patients' base disease, without increasing their hospital length of stay.

FROM: Moreira M, Medeiros EA, Pignatari AC, Wey SB, Cardo DM. The effect of nosocomial bloodstream infection by *Staphylococcus aureus* resistant to oxacillin on the mortality and the length of hospitalization. *Rev Assoc Med Bras* 1998;44:263-268.

New Assay for Rapid Detection of MRSA

A multiplex polymerase chain reaction (PCR), involving detection of the *mecA* and *femB* genes, was combined with a novel immunoassay system capable of detecting specific PCR products. The resulting PCR-immunoassay was evaluated in comparison with conventional microbiological techniques used in the routine diagnostic laboratory for the rapid identification of methicillin-resistant *Staphylococcus aureus* (MRSA), either in pure culture or in overnight broth cultures obtained following enrichment of patient screening swabs. Among the 480 purified isolates of staphylococci and 246 enrichment broths examined, only one false-negative result was obtained by PCR, compared with 18 false-negative results obtained by conventional methodology and demonstrated by further conventional examination. Five demonstrable false-positive results were obtained by conventional methodology, compared with a possible 10 by the PCR-immunoassay, although it was not certain that these 10 PCR results were true false positives, as, by definition, MRSA could not be isolated by conventional methodology.

The results indicated that the routine diagnostic laboratory was encountering difficulties in identifying MRSA correctly and that the conventional microbiological techniques lacked sensitivity. Overall, the PCR technique was more accurate and sensitive than conventional methodology in detecting MRSA, and results were available within 24 hours of screening swabs arriving in the laboratory, compared with a minimum of 48 to 72 hours by conventional techniques. The immunoassay system added to the usefulness of the method by allowing the detection of specific PCR products within 5 minutes of completing the PCR, without the normal additional step of agarose gel electrophoresis.

FROM: Towner KJ, Talbot DC, Curran R, Webster CA, Humphreys H. Development and evaluation of a PCR-based immunoassay for the rapid detection of methicillin-resistant *Staphylococcus aureus*. *J Med Microbiol* 1998; 47:607-613.



NINTH ANNUAL SCIENTIFIC MEETING

April 18 – 20, 1999 Hyatt Regency Embarcadero San Francisco, California



SATURDAY, APRIL 17, 1999

WORKSHOP 1

Update on Antimicrobial Resistance

WORKSHOP 2

Emerging Nosocomial Fungi

WORKSHOP 3

DNA Typing Techniques in Infection Control

WORKSHOP 4

How to Apply Clinical Epidemiology and Statistical Process Control Methods to Measure and Improve Quality of Care

APRIL 18-20, 1999

PLENARY SESSIONS:

BIOTERRORISM

1. The Threat
2. Public Health System Response
3. The Hospital Response

EMERGING INFECTIOUS HAZARDS IN THE HEALTHCARE SETTING

1. Infectious Risks & XenoTransplantation
2. Infectious Risks of Animals in the Healthcare Setting
3. The Hazards of Prions in the Healthcare Setting

IS THE ENVIRONMENT HAZARDOUS TO YOUR HEALTH?

1. Airborne Micro-organisms
2. Water
3. Inanimate Environment

BIOMATERIAL RELATED INFECTIONS

1. Pathogenesis
2. The Diagnostic Dilemma
3. We Should Use Therapeutic Antimicrobial Agents Incorporated Into or Onto Biomaterials to Prevent Infections-Point/Counter Point

**REGISTRATION DEADLINE:
APRIL 2, 1999**

SYMPOSIA:

HOT ISSUES OF INFECTION CONTROL IN SPECIALIZED PEDIATRIC AREAS

1. New Challenges in the Very Low Birth Weight Infant
2. Emerging Problems in Immunocompromised Hosts
3. Emerging Controversies in the Pediatric Intensive Care Areas

CUTTING-EDGE CHANGES IN THE OPERATING ROOM

1. Surgical Site Infections in Minimally Invasive Surgery (MIS)
2. Environmental issues in the Operating Room and their Impact on Surgical Site Infections
3. Surgical Site Infection Prevention in Latin America

QUALITY ASPECTS RELATED TO THE USE OF MEDICAL DEVICES

1. Introductory Remarks: Systems are changing, where can they be improved?
2. How does industry design, evaluate, and demonstrate the safety and efficacy of a medical device?
3. How do regulatory agencies ensure release of a safe medical device?
4. Institutional epidemiologists' role(s) in evaluating medical devices
5. Panel Discussion & Concluding Remarks

CONTROVERSIES IN HAND WASHING

1. Review of the Data
2. Compliance Obstacles and Measures to Increase Compliance in Traditional and Non-Traditional Areas
3. Scientific Basis and Controversies Related to Waterless Systems

MEET THE CONSULTANT BREAKFASTS

Seven Meet the Consultant Sessions will be offered on the Monday and Tuesday of the Scientific Meeting. You can consult with specialists in your field of interest.

For additional information regarding the SHEA Annual Meeting, please contact:

SHEA Meetings Department
19 Mantua Road
Mt. Royal, NJ 08061
Telephone: (609) 423-7222, ext. 350
Fax: (609) 423-3420
Email: meetings@tmg.smarthub.com