



## Effects of sleep fragmentation on appetite and related hormone concentrations over 24 h in healthy men

Hanne K. J. Gonnissen\*, Rick Hursel, Femke Rutters, Eveline A. P. Martens and Margriet S. Westerterp-Plantenga

Department of Human Biology, Maastricht University, PO Box 616, 6200 MD, Maastricht, The Netherlands

(Submitted 18 November 2011 – Final revision received 13 April 2012 – Accepted 13 April 2012 – First published online 8 June 2012)

### Abstract

In addition to short sleep duration, reduced sleep quality is also associated with appetite control. The present study examined the effect of sleep fragmentation, independent of sleep duration, on appetite profiles and 24 h profiles of hormones involved in energy balance regulation. A total of twelve healthy male subjects (age 23 (SD 4) years, BMI 24.4 (SD 1.9) kg/m<sup>2</sup>) completed a 24 h randomised crossover study in which sleep (23.30–07.30 hours) was either fragmented or non-fragmented. Polysomnography was used to determine rapid-eye movement (REM) sleep, slow-wave sleep (SWS) and total sleep time (TST). Blood samples were taken at baseline and continued hourly for the 24 h period to measure glucose, insulin, ghrelin, leptin, glucagon-like peptide 1 (GLP-1) and melatonin concentrations. In addition, salivary cortisol levels were measured. Visual analogue scales were used to score appetite-related feelings. Sleep fragmentation resulted in reduced REM sleep (69.4 min compared with 83.5 min;  $P < 0.05$ ) and preservation of SWS without changes in TST. In fragmented *v.* non-fragmented sleep, glucose concentrations did not change, while insulin secretion was decreased in the morning, and increased in the afternoon ( $P < 0.05$ ), and GLP-1 concentrations and fullness scores were lower ( $P < 0.05$ ). After dinner, desire-to-eat ratings were higher after fragmented sleep ( $P < 0.05$ ). A single night of fragmented sleep, resulting in reduced REM sleep, induced a shift in insulin concentrations, from being lower in the morning and higher in the afternoon, while GLP-1 concentrations and fullness scores were decreased. These results may lead to increased food intake and snacking, thus contributing to a positive energy balance.

**Key words:** Sleep fragmentation: Rapid-eye movement sleep: Insulin: Glucagon-like peptide 1: Fullness

During the last decade, the prevalence of obesity has increased worldwide to epidemic proportions. Obesity is considered to be a major health problem, as it is associated with type 2 diabetes, CVD and certain cancers<sup>(1)</sup>. Efforts to understand the causes of obesity have focused on a positive energy balance that occurs when energy intake exceeds energy expenditure<sup>(2,3)</sup>. The observation of a rapid decline in sleep duration parallel to the rapid rise in body weight has drawn attention to sleep deprivation as another possible contributor to a positive energy balance and therefore sleep may be a factor in the aetiology of obesity<sup>(4–8)</sup>.

Appetite regulation involves both tonic (long-term) and episodic (short-term) hunger and fullness signals. Tonic signals include leptin, which is secreted by adipose tissue, and insulin, which is secreted by the  $\beta$  cells in the pancreas. Episodic signals are triggered in response to the ingestion of food from the stomach (e.g. ghrelin) and intestines (e.g. glucagon-like peptide 1 (GLP-1)). Resistance to these hunger and satiating hormones results in overeating, leading to overweight and

obesity<sup>(9,10)</sup>. Experimental sleep restriction has been shown to make an impact on many of these hormones. Spiegel *et al.*<sup>(11)</sup> showed that a restriction of time in bed to 4 h compared to an extended bedtime of 10 h decreases leptin levels known to suppress appetite, and increases ghrelin levels that promote hunger. In addition, sleep restriction was associated with increased hunger and appetite, especially for energy-dense foods with high carbohydrate content<sup>(11)</sup>. Moreover, the change in the ratio of ghrelin:leptin between the two conditions was strongly correlated to the change in hunger rating<sup>(11)</sup>. Sleep restriction may also affect glucose and insulin metabolism. Sleep restriction to only 4 h of sleep reduced glucose tolerance relative to sleep recovery of 12 h of sleep<sup>(12)</sup> and reduced insulin sensitivity relative to normal sleep duration of 8.5 h of sleep<sup>(13)</sup>. Even sleep restriction to 5.5 h of time in bed per night for 2 weeks<sup>(14)</sup> or 5.0 h of time in bed for 1 week<sup>(15)</sup> showed similar reductions on insulin sensitivity. These experimental studies that restricted the time available for sleep observed significant changes in

**Abbreviations:** GLP-1, glucagon-like peptide 1; REM, rapid-eye movement; SWS, slow-wave sleep; TST, total sleep time; VAS, visual analogue scale.

\* **Corresponding author:** H. K. J. Gonnissen, fax +31 43 3670976, email hkj.gonnissen@maastrichtuniversity.nl

hormones involved in appetite regulation in a direction that would increase food intake. Thus, the link between sleep and appetite may be the modulation of appetite-related hormones.

However, besides sleep duration, sleep disturbance can also affect sleep quality by changing the composition of sleep in terms of sleep stages. Human sleep is not a homogeneous state but is composed of rapid-eye movement (REM) sleep and non-REM sleep. The non-REM sleep can be further divided into four stages of progressively deeper sleep. Stages 3 and 4 of non-REM sleep are also named slow-wave sleep (SWS). Different relationships may exist between distinct sleep stages and body weight regulation<sup>(16)</sup>. For example, the initiation of SWS coincides with hormonal changes that affect glucose regulation. Tasali *et al.*<sup>(17)</sup> showed that selective suppression of SWS, without affecting total sleep time (TST), increased the risk of type 2 diabetes. The design of the study by Tasali *et al.*<sup>(17)</sup> was quite extreme, reducing SWS by nearly 90%. However, more research is needed about the influence of sleep fragmentation, resulting in decreased sleep quality, on appetite control.

Therefore, the aim of the present study was to test the hypothesis that sleep fragmentation may lead to disturbed appetite control in a similar way as shorter sleep duration increases appetite. We thus examined the effect of sleep fragmentation, independent of sleep duration, on appetite profiles and 24 h profiles of hormones involved in energy balance regulation (glucose, insulin, ghrelin, leptin, GLP-1, melatonin and cortisol) in healthy men.

## Subjects and methods

### Subjects

A total of twelve healthy male subjects with a mean age of 23 (SD 4) years and with a mean BMI of 24.4 (SD 1.9) kg/m<sup>2</sup> participated in the present study. They were recruited via advertisements on notice boards at Maastricht University. The subjects underwent an initial screening including measurements of body weight and height and completed a questionnaire related to health, use of medication, smoking behaviour, alcohol consumption, sleeping behaviour and food allergies. All subjects were in good health, non-smokers, not using medication and, at most, moderate alcohol consumers (<10 drinks a week). Baseline characteristics of the subjects are presented in Table 1. Body composition was measured using the <sup>2</sup>H dilution technique<sup>(18,19)</sup>.

All procedures were carried out with adequate understanding and subjects provided written informed consent at the start of the first test day. Subject recruitment started in June 2009 and the study was conducted between September 2009 and May 2010. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Medical Ethical Committee of Maastricht University Medical Centre. The study was registered in the Dutch Trial Register (registration number NTR1919). The power calculation of the present study was based on a sleep deprivation study<sup>(20)</sup>. In the reference study, a crossover design with twelve subjects was performed and

**Table 1.** Subject characteristics

(Mean values and standard deviations, *n* 12)

Characteristics	Mean	SD
Age (years)	23	4
BM (kg)	79.7	5.3
Height (cm)	180.9	4.4
BMI (kg/m <sup>2</sup> )	24.4	1.9
FFM (kg)	62.4	6.7
FM (kg)	16.2	5.8
%BF	20.5	6.8
Self-reported sleep duration (h/night)	8.2	1.0
Self-reported sleep latency (min)	17.3	7.6

BM, body mass; FFM, fat-free mass; FM, fat mass; %BF, percentage of body fat.

resulted in significant differences between 4 h of sleep compared to 8 h of sleep<sup>(20)</sup>. The metabolic units, used in the present study, ensure a highly controlled situation in which subjects are in a stable environment.

### Study design

Recently, Hursel *et al.*<sup>(21)</sup> reported a respiration chamber study in which they determined the effects of sleep fragmentation on energy expenditure and substrate oxidation in fifteen healthy men. In a subgroup of these participants, we performed a separate experiment in a metabolic ward where regular blood sampling, also overnight, was possible. The data presented in the present paper are obtained from different nights compared to those reported in Hursel *et al.*<sup>(21)</sup>. The two experiments were separated by at least 1 week and the order was randomised across the subjects.

The study had a randomised, single-blinded, crossover design. Subjects participated in two 24 h metabolic unit experiments, separated by at least 1 week and at most 6 weeks; and 2 days before the experiment, subjects were asked to sleep for 8 h during the nights according to their habitual sleep duration (self-reported sleep duration, Table 1). During both phases, subjects arrived at the university at 18.00 hours and dinner was served at 18.30 hours. At 20.00 hours, electrodes for electroencephalogram, electromyogram and electrooculogram recordings were applied according to standardised criteria<sup>(22)</sup>. At 20.30 hours, a canula was placed into a dorsal hand vein of the contralateral arm. Blood samples, for the determination of glucose, insulin, ghrelin, leptin, GLP-1 and melatonin, were taken at baseline (21.00 hours) and continued hourly for the 24 h period. An experimenter entered the room to collect these blood samples, without disturbing the subjects. This was checked with the polysomnography data. Furthermore, saliva samples were collected every hour during the day (21.00–23.00 and 08.00–20.00 hours) to determine cortisol levels. The subjects were allowed to sleep from 23.30 to 07.30 hours ('non-fragmented' condition) or were woken up several times during the night ('fragmented' condition). Lights were switched off automatically at 23.30 hours and switched on at 07.30 hours, resulting in 8 h of sleep opportunity in a dark and quiet environment. The order of the two conditions was randomised across the subjects to prevent

any order effects. Wake-up calls, varying in frequency between 500 and 2000 Hz and intensity between 40 and 110 dB, occurred approximately every 90 min. The subjects confirmed waking up since they had to put off the alarm, which took about 2 min. The alarm was near the bed so that subjects did not need to get up and the lights were kept off. The next day, standardised meals were served at 08.30, 13.30 and 18.30 hours. At 20.00 hours, the canula and the electrodes were removed. Artificial light intensity during the day (07.30–23.30 hours) was always above 400 lux (Energy Saver, Tornado E27, 900 lumen; Philips Lighting).

### Sleep recordings

To ensure that sleep fragmentation was successful, polysomnographic recordings were obtained throughout the 24 h by using BrainRT (OSG BVBA). Sleep is recorded in occipital and central regions of the brain. Surface electrodes were used to record the electroencephalographic signals; two central (C3–A2 and C4–A1) and two occipital (O1–A2 and O2–A1), bilateral electrooculograms and submental electromyograms. All records were visually scored in 30-s epochs according to Rechtschaffen & Kales<sup>(22)</sup> by the same experienced person blinded to the experimental condition. Sleep parameters of interest were REM sleep, SWS and TST. Sleep latency is defined as the time to fall asleep. Daytime naps were not allowed.

### Energy intake

During their stay and 2 d before the experiment, subjects were fed in energy balance. The energy content of the food was tailored to the energy requirements of each subject by calculating BMR with the Harris & Benedict<sup>(23)</sup> equation that uses sex, age, height and weight. To estimate the total energy requirement at home, the BMR was multiplied with a physical activity index of 1.75 estimated by means of a computer simulation program<sup>(24)</sup>. The total energy requirement in the metabolic unit was estimated by multiplying the BMR with a physical activity index of 1.35. Energy intake was divided over the meals as 20% for breakfast, 40% for lunch and 40% for dinner. The macronutrient composition of the diet was 12/55/33% of energy (protein/carbohydrate/fat). Participants were required to finish each meal and were not allowed to eat additional food. Water was freely available during the whole experiment.

### Questionnaires

The appetite questionnaire was composed of visual analogue scale (VAS, in mm) questions on subjective feelings of hunger, fullness and desire to eat. Opposing extremes of each feeling were described at either end of a 100 mm horizontal line, and subjects marked the line to indicate how they felt at that moment. These questionnaires were completed every hour and before and after each meal.

Aspects of mood were assessed using the Dutch translation of the Profile of Mood States. This questionnaire contains

seventy adjectives that are rated on a five-point scale, anchored by 'much like this' to 'much unlike this' and is divided into five categories (depression, tension, confusion, fatigue and anger). An increase in Profile of Mood States scores is associated with a worsening in mood<sup>(25)</sup>. Mood profiles were assessed before each meal.

Anxiety was assessed using the Dutch translation of the State-Trait Anxiety Inventory. This questionnaire is composed of twenty questions rated on a four-point scale, ranging from 'much like this' to 'much unlike this'. An increase in State-Trait Anxiety Inventory scores is associated with an increase in anxiety<sup>(26)</sup>. State anxiety was assessed every hour and before and after each meal.

### Blood and saliva samples

Blood was distributed into EDTA tubes for glucose, insulin, leptin and active ghrelin measurements. For active GLP-1 measurement, blood was collected in EDTA tubes with added dipeptidyl peptidase IV inhibitor. Plasma was obtained by centrifugation at 4°C for 10 min at 3000 rpm. For melatonin measurement, blood was collected in clot tubes, containing 'Silica Clot Activator'. Blood in the clot tubes was allowed to clot for 30 min and was centrifuged at 3000 rpm, 4°C for 10 min to obtain serum. All samples were stored at –80°C until analysis. Plasma glucose concentrations were determined using the hexokinase method (Glucose HK 125 kit; ABX Diagnostics). Insulin and plasma active ghrelin concentrations were measured by RIA (Linco Research, Inc.). Plasma active GLP-1 samples were analysed using ELISA (EGLP-35K; Linco Research, Inc.). Plasma leptin concentrations were measured using the human leptin RIA-kit (Linco Research, Inc.). Serum melatonin concentrations were determined by RIA (IBL International GmbH).

To determine cortisol levels, saliva samples were collected using cotton swabs (Salivettes, Sarstedt). Subjects were instructed to gently chew on the swab for 1 min. Cotton swabs were then transferred to the plastic containers and stored at –20°C until analysis. Salivary cortisol concentrations were measured at the laboratory of Professor Dr C. Kirschbaum, Dresden University of Technology, Germany. After thawing, saliva samples were centrifuged at 3000 rpm for 10 min. Luminescence immunoassay (IBL) with intra- and inter-assay precision of 2.5 and 4.7%, respectively, was used to measure salivary cortisol concentrations.

### Statistical analysis

Data are presented as means with their standard errors, unless otherwise indicated. AUC were calculated using the trapezoidal method. ANOVA with repeated measures was carried out to determine the conditional effects of fragmented *v.* non-fragmented sleep and the effects of time, on measurements of glucose, insulin, ghrelin, leptin, GLP-1, melatonin and cortisol levels. If relevant, differences between fragmented and non-fragmented sleep condition were analysed per time point. To assess the strength of the within-subject relationship between changes in VAS scores for fullness and



GLP-1 concentrations, we calculated, separately for each subject, regression slopes and  $R^2$  values for the regression of fullness scores on GLP-1 concentrations. We presented the mean and 95% CI for the mean and interquartile ranges (Q1–Q3) of the observed slopes and  $R^2$  values. Student's one-sample  $t$  tests were used to test whether the means of the regression slopes were different from zero. All tests were two-sided and the level for establishing significant differences was taken at  $P < 0.05$ . Data were analysed using SPSS 11 (SPSS, Inc.).

## Results


### Sleep

There were no significant differences between the non-fragmented and fragmented nights in sleep latency, time awake, TST and sleep stage 1 (Table 2). Subjects woke up approximately five times during the fragmented night. Although the amount of SWS was reduced in the fragmented night compared with the non-fragmented night, this difference was not statistically significant (Table 2). REM sleep was significantly shorter during the fragmented night compared with the non-fragmented night ( $P < 0.05$ , Table 2). Consequently, sleep stage 2 was significantly longer in the fragmented night compared with the non-fragmented night ( $P < 0.001$ , Table 2).

### Glucose and insulin concentrations

However, 24 h glucose concentrations did not differ between the fragmented sleep and non-fragmented sleep nights (Fig. 1(a), Table 3). There was an overall effect of time on glucose levels ( $P < 0.001$ ).

There was no overall effect of sleep fragmentation on 24 h insulin concentrations. Postprandial insulin secretion changed significantly differently after the fragmented *v.* non-fragmented sleep nights. Following fragmented sleep, the postprandial rise in insulin was significantly lower after breakfast and higher after dinner, while after non-fragmented sleep the postprandial rise in insulin was significantly higher after breakfast and lower after dinner (Fig. 1(b), time  $\times$  condition effect,  $F = 4.9$ ;  $P < 0.05$ ). There was an overall effect



**Fig. 1.** Mean (a) glucose and (b) insulin concentrations in fragmented sleep (■) and non-fragmented sleep (▲) conditions ( $n$  12). \* $P < 0.05$  (analyses per time point). ANOVA (repeated measures) showed a significant time  $\times$  condition effect ( $F = 4.9$ ,  $P < 0.05$ ). ↓ indicates meal times.

of time on insulin levels ( $P < 0.001$ ). Analysis per time point revealed that following a night of fragmented sleep, post-breakfast insulin concentrations were lower, while post-dinner insulin concentrations were higher compared with a night of non-fragmented sleep (Fig. 1(b),  $P < 0.05$ ).

### Ghrelin, leptin and melatonin concentrations

There were no significant differences in the 24 h ghrelin, leptin and melatonin concentrations between the fragmented sleep and non-fragmented sleep nights (AUC, respectively: 67 058

**Table 2.** Sleep parameters during non-fragmented and fragmented nights

(Mean values with their standard errors,  $n$  12)

Parameters	Non-fragmented night		Fragmented night		$P^*$
	Mean	SEM	Mean	SEM	
Sleep latency (min)	22.7	4.7	20.3	3.7	0.63
Wake (min)	30.0	8.2	27.9	9.3	0.75
Total sleep time (min)	428.4	11.7	433.2	9.7	0.45
Stage 1 (min)	18.8	3.9	19.0	3.3	0.97
Stage 2 (min)	244.0	9.8	273.9	10.5	$< 0.001$
SWS (min)	82.0	9.0	70.9	9.1	0.17
REM sleep (min)	83.5	7.9	69.4	6.0	$< 0.05$

SWS, slow-wave sleep; REM, rapid-eye movement.

\* Differences between non-fragmented and fragmented nights (ANOVA – repeated measures).

**Table 3.** Blood and saliva parameters in the non-fragmented sleep and fragmented sleep conditions

(Mean values with their standard errors,  $n$  12)

Parameters	Non-fragmented night		Fragmented night	
	Mean	SEM	Mean	SEM
Glucose (mmol/l)	5.6	0.1	5.6	0.1
Insulin (pmol/l)	30.6	2.4	31.2	2.5
Ghrelin (ng/l)	52.9	6.2	48.4	5.9
Leptin ( $\mu$ g/l)	4.2	0.6	4.7	0.8
Melatonin (ng/l)	38.5	4.2	38.6	5.4
GLP-1 (pmol/l)	4.4	0.9	4.1	0.7
Cortisol (nmol/l)	8.6	0.5	8.5	0.5

GLP-1, glucagon-like peptide 1.

(SEM 28 841) *v.* 73 173 (SEM 8678) ng/l × min; 6426 (SEM 1063) *v.* 5718 (SEM 866) µg/l × min; 54 739 (SEM 7855) *v.* 56 465 (SEM 6486) ng/l × min, Table 3). There was an overall effect of time on ghrelin, leptin and melatonin concentrations ( $P < 0.001$ ).

### Glucagon-like peptide 1 concentrations and appetite profile

There was no overall effect of sleep fragmentation and no time × condition interaction effect on 24 h GLP-1 concentrations. GLP-1 concentrations were different over time ( $P < 0.001$ ). Since previous studies<sup>(20,27)</sup> showed an increased food intake and snacking during the afternoon or evening, we performed a closer inspection of the GLP-1 data from the afternoon. Analyses of AUC during the afternoon and per time point revealed that following a night of fragmented sleep, GLP-1 concentrations were significantly lower in the afternoon (Fig. 2(a) and (b);  $P < 0.05$ ) than after a night of non-fragmented sleep.

In addition, there was also an effect of time on fullness and desire-to-eat ratings ( $P < 0.001$ ). Analysis per time point revealed that in the afternoon following a night of fragmented sleep, subjects were less full (Fig. 2(c);  $P < 0.05$ ) than after a night of non-fragmented sleep. After dinner, desire-to-eat ratings were significantly higher during the fragmented sleep condition than during the non-fragmented sleep condition (24.6 (SEM 8.3) *v.* 14.9 (SEM 6.6) mm VAS;  $P < 0.05$ ).


In both the non-fragmented sleep and fragmented sleep conditions, GLP-1 concentrations changed synchronously with VAS fullness scores (Fig. 3). The regression slopes and  $R^2$  values were significantly higher in the non-fragmented sleep condition (Table 4).

### Cortisol concentrations and mood state

With respect to stress-related parameters, there was a time × condition interaction effect on cortisol concentrations ( $P < 0.01$ ,  $F = 2.26$ ), peak values were higher in the morning following a night of non-fragmented sleep, whereas evening cortisol levels remained higher following a night of fragmented sleep (Fig. 4). Cortisol concentrations were different over time ( $P < 0.001$ , Fig. 4). Analysis per time point revealed that cortisol concentrations were significantly lower in the morning after the fragmented sleep night than after the non-fragmented sleep night (Fig. 4,  $P < 0.01$ ). In the evening, cortisol concentrations were significantly higher in the fragmented sleep condition compared with the non-fragmented sleep condition (Fig. 4,  $P < 0.05$ ).

There was an overall time effect ( $P < 0.05$ ) and a time × condition interaction effect ( $P < 0.05$ ,  $F = 5.0$ ) on the Profile of Mood States scores for the category 'fatigue'. The morning after a night of fragmented sleep, Profile of Mood States scores for 'fatigue' were significantly higher than after a night of non-fragmented sleep (24.7 (SEM 1.3) *v.* 20.5 (SEM 1.4);  $P < 0.05$ ).


There were no significant differences in the State-Trait Anxiety Inventory scores between the fragmented sleep and non-fragmented sleep nights (AUC: 21 604 (SEM 1202) *v.* 19 534 (SEM 877)).



**Fig. 2.** Mean glucagon-like peptide 1 (GLP-1) concentrations in fragmented sleep (■) and non-fragmented sleep (▲) conditions ( $n = 12$ ). Data are presented in lines (a) and bars (b, AUC from 14.00 to 20.00 hours). Mean visual analogue scale (VAS) fullness scores (c) in fragmented sleep (■) and non-fragmented sleep (▲) conditions ( $n = 12$ ). \* $P < 0.05$  (analyses per time point). ↓ indicates meal times.

### Discussion

In the present study, sleep fragmentation resulted in the preservation of SWS and substitution of REM sleep with sleep stage 2 without changes in TST and time awake. Sleep fragmentation occurred less frequently compared to Hursel *et al.*<sup>(21)</sup> (five *v.* seven times), which explains why the alterations in sleep parameters were less pronounced in this study compared to the previous study. Both studies show a reduction in SWS and REM sleep, which are regarded as the restorative



**Fig. 3.** Mean glucagon-like peptide 1 (GLP-1) concentrations (black lines) and Visual Analogue Scale (VAS) fullness scores (grey lines) in non-fragmented sleep (a, ▲) and fragmented sleep (b, ■) conditions (*n* 12).

parts of sleep and the sleep parameters related to health. However, in the present study, the reduction in SWS did not reach significance. Although there were no significant differences in time awake, sleep stage 1 and SWS in our present study, we were able to show an effect of altering the amount of REM sleep without changing TST, thereby enabling us to investigate the effect of sleep quality.

Our results show that a night of fragmented sleep is accompanied by a change in the pattern of postprandial insulin secretion over the day without changes in glucose secretion. Several studies have already indicated that sleep restriction in healthy subjects induces a reduction in insulin sensitivity<sup>(13–15)</sup>. In addition, sleep fragmentation across all stages was associated with a decrease in insulin sensitivity<sup>(28)</sup>. Tasali *et al.*<sup>(17)</sup> even showed that selective suppression of SWS could worsen the reduced insulin secretion in normal people. The present study shows significant differences in insulin release only on two specific time points; therefore, the evidence is too limited to draw a firm conclusion about insulin sensitivity from these data. Surprisingly, analysis of insulin secretion per time point revealed that following a night of fragmented sleep, insulin concentrations were less increased after breakfast. Although non-significant, also glucose concentrations after breakfast were slightly lower following a night of fragmented sleep than following a night of non-fragmented sleep. We speculate that more glycogen has been used during the fragmented night due to higher activity-induced energy expenditure during the night for putting off the alarm<sup>(21,29,30)</sup>. The overall effect of time on glucose concentrations is caused by its meal-related secretion pattern. Glucose concentrations were low before each meal and peaked after each meal.

The higher insulin secretion in the afternoon after fragmented sleep may imply that a vulnerable condition appears in the afternoon. Previous studies<sup>(20,27,31–33)</sup> showed indeed an increased food intake and an increased intake of energy from snacks after partial sleep deprivation. In relation to appetite, GLP-1 concentrations and fullness scores were decreased in the afternoon following a night of fragmented sleep. The overall effect of time on GLP-1 concentrations is caused by its meal-related secretion pattern. GLP-1 concentrations were low before each meal and peaked after each meal. In agreement with previous studies<sup>(9,34)</sup> GLP-1 concentrations changed synchronously with VAS fullness scores. However, the present study shows that the correlation between GLP-1 concentrations and VAS fullness scores decreases after fragmented sleep. After dinner, the appetite effect continued by showing a larger desire to eat. So, it seems that a vulnerable condition following sleep fragmentation occurs in the afternoon. During the 48 h respiration chamber experiment with the same subjects, we observed an increase in carbohydrate oxidation


**Table 4.** Observed slopes and  $R^2$  values for the within-subject relationships between visual analogue scale fullness scores and glucagon-like peptide 1 concentrations in the non-fragmented sleep and fragmented sleep conditions

(Mean values with their standard errors; 95% confidence intervals and interquartile ranges (IQR), *n* 12)

	Non-fragmented sleep night				Fragmented sleep night				<i>P</i> †
	Mean	SEM	CI	IQR	Mean	SEM	CI	IQR	
Slope	10.0*	1.4	7.2, 12.7	6.0–2.3	7.3*	1.1	5.1, 9.6	4.2–0.3	< 0.05
$R^2$	0.5	0.05	0.4, 0.6	0.4–0.6	0.3	0.06	0.2, 0.4	0.1–0.4	< 0.01

\* Means of the regression slopes were significantly different from zero ( $P < 0.0001$ , Student's one-sample *t* test).

† Differences between non-fragmented sleep and fragmented sleep nights (ANOVA – repeated measures).



**Fig. 4.** Mean cortisol concentrations in fragmented sleep (■) and non-fragmented sleep (▲) conditions ( $n$  12). \* $P < 0.05$  (analyses per time point). ANOVA (repeated measures) showed a significant time  $\times$  condition effect ( $F = 2.26$ ,  $P < 0.01$ ). ↓ indicates meal times.

when sleep was fragmented<sup>(21)</sup>. In addition, Nedeltcheva *et al.*<sup>(35)</sup> reported that the respiratory quotient was increased for short sleepers, which implies less fat oxidation and more carbohydrate oxidation. We speculate that these results may indicate a glycogen depletion that leads to hypoglycaemia and thereby may underscore a vulnerable condition leading to increased food intake and snack consumption.

Both epidemiological and experimental observations of lower leptin and higher ghrelin concentrations during the day with short sleep duration<sup>(36,37)</sup> have led to the hypothesis that sleep deprivation may affect energy balance through such hormone changes. The lack of changes in ghrelin and leptin during the day after a night of fragmented sleep in our study may be explained by our experimental design investigating the effect of short-term sleep fragmentation independent of sleep duration. The overall effect of time on leptin and melatonin concentrations is caused by their circadian secretion pattern, while the overall effect of time on ghrelin concentrations is due to its meal-related secretion pattern. Leptin and melatonin concentrations increased during the night and decreased during the day. Ghrelin concentrations peaked before each meal and decreased immediately after each meal.

Because sleep fragmentation may affect several neuroendocrine signals involved in the control of substrate utilisation, we also examined cortisol concentrations. The sharp morning rise and the steep fall to low evening levels are two characteristic features of the human daily cortisol rhythm. Other studies<sup>(38,39)</sup> have previously demonstrated that both morning and evening cortisol levels are modulated by even a single night of reduced sleep. Corresponding with these studies, the present results indicate that after a single night of fragmented sleep, cortisol levels were significantly reduced after awakening and were elevated in the evening.

The strengths of our study included the standardised protocol with detailed sleep scoring using polysomnography. Subjects were fed in energy balance 2 d before and during the experiment. Moreover, blood samples were taken every hour

for 24 h. In addition, a refinement that our study shows is in connecting the glucose metabolism and appetite-related parameters to the different sleep stages. This suggests that TST is not primarily indicative of sleep quality, which consists of REM sleep and SWS. Thus, affecting TST, such as previous studies have demonstrated, may affect glucose metabolism and appetite-related parameters since sleep quality will be reduced. Our observation on reduced sleep quality is based upon reduced REM sleep while SWS is preserved. Tasali *et al.*<sup>(17)</sup> showed that the suppression of SWS adversely affected glucose homeostasis. However, assessing SWS suppression and SWS recovery only partly explains the effect of sleep quality on glucose metabolism because sleep restriction also results in a reduction of REM sleep<sup>(40,41)</sup>. A possible limitation of the present study is the mild sleep fragmentation, in contrast to studies examining the effect of sleep restriction and sleep recovery. However, mild sleep fragmentation is more likely to occur in everyday life with its street noise and cell phones that are not switched off during the night. Our study indicates that even mild sleep fragmentation during a single night may contribute to a vulnerable condition in the afternoon, accompanied by increased insulin secretion, lower GLP-1 concentrations and diminished fullness. Another limitation might be that the acute effects from this study cannot be extrapolated to chronic effects since adaptations might occur. Therefore, more research over the long term is necessary to establish the initial observations. Finally, there was the instance of an experimenter entering the room softly to take blood samples. We had anticipated that this might be a limitation, but we decided to use this procedure because of safety concerns. Therefore, after each night it was checked whether this procedure disturbed the sleep pattern, and no acute changes in sleep phases coincided with entering the room. In addition, this procedure occurred in both conditions and only the night with fragmented sleep showed a reduction of REM sleep.

In the present study, sleep fragmentation led to reduced sleep quality by reducing the amount of REM sleep. Several studies<sup>(42–44)</sup> suggest that REM sleep may have a role in metabolism and obesity. A population-based study found an association between reduced amount of REM sleep and central obesity in women<sup>(42)</sup>, while a cross-sectional study indicated that reduced REM sleep time was associated with elevated BMI in children and adolescents<sup>(44)</sup>. The significant association between reduced REM sleep and overweight suggests that REM sleep loss may alter the balance of energy intake and energy expenditure. It has been shown that sleeping metabolic rate is significantly higher in REM sleep<sup>(45)</sup> and diminished sleeping metabolic rate is again associated with elevated BMI<sup>(46)</sup>. Moreover, the present study shows that also endocrine changes are sensitive to reduced REM sleep. Increased insulin concentrations and decreased GLP-1 concentrations in the afternoon due to REM sleep loss decreased fullness scores, which in turn could lead to increased food intake and snacking if subjects had been allowed *ad libitum* food intake. In addition to an increase in total energy intake, the proportional intake of specific macronutrients, such as carbohydrates or fat, could have increased. It has been already

shown that sleep restriction increased the appetite for energy-dense nutrients with high carbohydrate content, including sweets, salty snacks and starchy foods<sup>(11)</sup>. In addition, Nedeltcheva *et al.*<sup>(27)</sup> found that bedtime restriction was associated with increased consumption of energy from snacks with higher carbohydrate content. While in another study, six nights of 4 h in bed was associated with more energy consumption, mostly due to increased fat consumption<sup>(33)</sup>. Beside the homeostatic system, also the hedonic system plays a role in food intake regulation. Previous animal studies showed that REM sleep deprivation produced a decline in motivation for food reward<sup>(47)</sup>. So, also in humans, reduced REM sleep could lead to a decrease in the reward values of food which may shift food choice to higher reward and higher energy foods and thereby contributing to a positive energy balance. Moreover, Van Cauter *et al.*<sup>(48)</sup> observed an inverse relationship between nadir cortisol levels and REM sleep. Thus, the elevation of the evening cortisol levels in the fragmented sleep condition of this study may be due to the decreased amount of REM sleep during the fragmented sleep. Raised cortisol concentrations in the evening are thought to reflect an impairment of the negative-feedback control of the hypothalamo–pituitary–adrenal axis that has been related to obesity<sup>(49)</sup>. Earlier studies using extreme forms of sleep deprivation may reflect the effect of decreases in the subject-specific amount of SWS and REM sleep, thereby implying a reduction in sleep quality. Therefore, they cannot distinguish between the effect of reduction of TST or of reduction of sleep quality. In our study, we reduced sleep quality without reducing TST and report that reduced sleep quality already affects insulin, GLP-1, cortisol concentrations and fullness scores. Therefore, we suggest that the quality of sleep is more important than the duration of sleep. This also may explain why not all people who sleep less than 6 h are overweight/obese, as they may have a sufficient amount of REM sleep.

In summary, a single night of fragmented sleep, resulting in reduced REM sleep, induced a shift in insulin concentrations; from being lower in the morning and higher in the afternoon while GLP-1 concentrations and fullness scores were decreased. Together, these results may create a vulnerable condition in the afternoon, which could lead to increased food intake and snacking, thus contributing to a positive energy balance.

### Acknowledgements

The authors' responsibilities relating to the present study were as follows: H. K. J. G., R. H., F. R. and M. S. W.-P. designed the study; H. K. J. G., R. H., F. R. and E. A. P. M. conducted research and analysed the data; H. K. J. G. wrote the manuscript; and M. S. W.-P. supervised the execution of the study and reviewed the manuscript. None of the authors had a personal or financial conflict of interest, or any other conflict of interest that could be perceived as prejudicing the impartiality of the research reported. This research did not receive any specific grant or funding from any organisation in the public, commercial or not-for-profit sector.

### References

1. Haslam DW & James WP (2005) Obesity. *Lancet* **366**, 1197–1209.
2. Hansen K, Shriver T & Schoeller D (2005) The effects of exercise on the storage and oxidation of dietary fat. *Sports Med* **35**, 363–373.
3. Curioni CC & Lourenco PM (2005) Long-term weight loss after diet and exercise: a systematic review. *Int J Obes (Lond)* **29**, 1168–1174.
4. Van Cauter E & Knutson KL (2008) Sleep and the epidemic of obesity in children and adults. *Eur J Endocrinol* **159**, Suppl. 1, S59–S66.
5. Gangwisch JE (2009) Epidemiological evidence for the links between sleep, circadian rhythms and metabolism. *Obes Rev* **10**, Suppl. 2, 37–45.
6. Marshall NS, Glozier N & Grunstein RR (2008) Is sleep duration related to obesity? A critical review of the epidemiological evidence. *Sleep Med Rev* **12**, 289–298.
7. Knutson KL & Van Cauter E (2008) Associations between sleep loss and increased risk of obesity and diabetes. *Ann N Y Acad Sci* **1129**, 287–304.
8. Nielsen LS, Danielsen KV & Sorensen TI (2011) Short sleep duration as a possible cause of obesity: critical analysis of the epidemiological evidence. *Obes Rev* **12**, 78–92.
9. Blundell JE, Levin F, King NA, *et al.* (2008) Overconsumption and obesity: peptides and susceptibility to weight gain. *Regul Pept* **149**, 32–38.
10. Adam TC & Westerterp-Plantenga MS (2005) Glucagon-like peptide-1 release and satiety after a nutrient challenge in normal-weight and obese subjects. *Br J Nutr* **93**, 845–851.
11. Spiegel K, Tasali E, Penev P, *et al.* (2004) Brief communication: sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* **141**, 846–850.
12. Spiegel K, Leproult R & Van Cauter E (1999) Impact of sleep debt on metabolic and endocrine function. *Lancet* **354**, 1435–1439.
13. Donga E, van Dijk M, van Dijk JG, *et al.* (2010) A single night of partial sleep deprivation induces insulin resistance in multiple metabolic pathways in healthy subjects. *J Clin Endocrinol Metab* **95**, 2963–2968.
14. Nedeltcheva AV, Kessler L, Imperial J, *et al.* (2009) Exposure to recurrent sleep restriction in the setting of high caloric intake and physical inactivity results in increased insulin resistance and reduced glucose tolerance. *J Clin Endocrinol Metab* **94**, 3242–3250.
15. Buxton OM, Pavlova M, Reid EW, *et al.* (2010) Sleep restriction for 1 week reduces insulin sensitivity in healthy men. *Diabetes* **59**, 2126–2133.
16. Somers VK, Dyken ME, Mark AL, *et al.* (1993) Sympathetic-nerve activity during sleep in normal subjects. *N Engl J Med* **328**, 303–307.
17. Tasali E, Leproult R, Ehrmann DA, *et al.* (2008) Slow-wave sleep and the risk of type 2 diabetes in humans. *Proc Natl Acad Sci U S A* **105**, 1044–1049.
18. Westerterp KR, Wouters L & van Marken Lichtenbelt WD (1995) The Maastricht protocol for the measurement of body composition and energy expenditure with labeled water. *Obes Res* **3**, Suppl. 1, 49–57.
19. Schoeller DA, van Santen E, Peterson DW, *et al.* (1980) Total body water measurement in humans with <sup>18</sup>O and <sup>2</sup>H labeled water. *Am J Clin Nutr* **33**, 2686–2693.

20. Brondel L, Romer MA, Nougues PM, *et al.* (2010) Acute partial sleep deprivation increases food intake in healthy men. *Am J Clin Nutr* **91**, 1550–1559.
21. Hursel R, Rutters F, Gonnissen H, *et al.* (2011) Effects of sleep fragmentation in healthy men on energy expenditure, substrate oxidation, physical activity, and exhaustion measured over 48 h in a respiratory chamber. *Am J Clin Nutr* **94**, 804–808.
22. Rechtschaffen A & Kales A (1968) *A Manual of Standardized Terminology, Techniques and Scoring System for Sleep Stages of Human Subjects*. Bethesda, MD: National Institute of Neurological Diseases and Blindness.
23. Harris JA & Benedict FG (1918) A biometric study of human basal metabolism. *Proc Natl Acad Sci U S A* **4**, 370–373.
24. Westerterp KR, Donkers JH, Fredrix EW, *et al.* (1995) Energy intake, physical activity and body weight: a simulation model. *Br J Nutr* **73**, 337–347.
25. Albrecht RR & Ewing SJ (1989) Standardizing the administration of the profile of mood states (POMS): development of alternative word lists. *J Pers Assess* **53**, 31–39.
26. Tenenbaum G, Furst D & Weingarten G (1985) A statistical reevaluation of the STAI anxiety questionnaire. *J Clin Psychol* **41**, 239–244.
27. Nedeltcheva AV, Kilkus JM, Imperial J, *et al.* (2009) Sleep curtailment is accompanied by increased intake of calories from snacks. *Am J Clin Nutr* **89**, 126–133.
28. Stamatakis K & Punjabi NM (2009) Effects of sleep fragmentation on glucose metabolism in normal subjects. *Chest* **137**, 95–101.
29. Kuipers H, Keizer HA, Brouns F, *et al.* (1987) Carbohydrate feeding and glycogen synthesis during exercise in man. *Pflugers Arch* **410**, 652–656.
30. Schrauwen P, van Marken Lichtenbelt WD, Saris WH, *et al.* (1997) Role of glycogen-lowering exercise in the change of fat oxidation in response to a high-fat diet. *Am J Physiol* **273**, E623–E629.
31. Bosy-Westphal A, Hinrichs S, Jauch-Chara K, *et al.* (2008) Influence of partial sleep deprivation on energy balance and insulin sensitivity in healthy women. *Obes Facts* **1**, 266–273.
32. Weiss A, Xu F, Storfer-Isser A, *et al.* (2010) The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep* **33**, 1201–1209.
33. St-Onge MP, Roberts AL, Chen J, *et al.* (2011) Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr* **94**, 410–416.
34. Lemmens SG, Martens EA, Kester AD, *et al.* (2011) Changes in gut hormone and glucose concentrations in relation to hunger and fullness. *Am J Clin Nutr* **94**, 717–725.
35. Nedeltcheva AV, Kilkus JM, Imperial J, *et al.* (2010) Insufficient sleep undermines dietary efforts to reduce adiposity. *Ann Intern Med* **153**, 435–441.
36. Taheri S, Lin L, Austin D, *et al.* (2004) Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* **1**, e62.
37. Chaput JP, Despres JP, Bouchard C, *et al.* (2008) The association between sleep duration and weight gain in adults: a 6-year prospective study from the Quebec Family Study. *Sleep* **31**, 517–523.
38. Omisade A, Buxton OM & Rusak B (2010) Impact of acute sleep restriction on cortisol and leptin levels in young women. *Physiol Behav* **99**, 651–656.
39. Leproult R, Copinschi G, Buxton O, *et al.* (1997) Sleep loss results in an elevation of cortisol levels the next evening. *Sleep* **20**, 865–870.
40. Banks S & Dinges DF (2007) Behavioral and physiological consequences of sleep restriction. *J Clin Sleep Med* **3**, 519–528.
41. Van Dongen HP, Maislin G, Mullington JM, *et al.* (2003) The cumulative cost of additional wakefulness: dose–response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* **26**, 117–126.
42. Theorell-Haglow J, Berne C, Janson C, *et al.* (2010) Associations between short sleep duration and central obesity in women. *Sleep* **33**, 593–598.
43. Horne J (2009) REM sleep, energy balance and 'optimal foraging'. *Neurosci Biobehav Rev* **33**, 466–474.
44. Liu X, Forbes EE, Ryan ND, *et al.* (2008) Rapid eye movement sleep in relation to overweight in children and adolescents. *Arch Gen Psychiatry* **65**, 924–932.
45. Fontvieille AM, Rising R, Spraul M, *et al.* (1994) Relationship between sleep stages and metabolic rate in humans. *Am J Physiol* **267**, E732–E737.
46. Zhang K, Sun M, Werner P, *et al.* (2002) Sleeping metabolic rate in relation to body mass index and body composition. *Int J Obes Relat Metab Disord* **26**, 376–383.
47. Hanlon EC, Andrzejewski ME, Harder BK, *et al.* (2005) The effect of REM sleep deprivation on motivation for food reward. *Behav Brain Res* **163**, 58–69.
48. Van Cauter E, Leproult R & Plat L (2000) Age-related changes in slow wave sleep and REM sleep and relationship with growth hormone and cortisol levels in healthy men. *JAMA* **284**, 861–868.
49. Bjorntorp P (2001) Do stress reactions cause abdominal obesity and comorbidities? *Obes Rev* **2**, 73–86.