NOTES FOR AUTHORS

The *Bulletin of Entomological Research* publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the *Bulletin* is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

Page Format. The *Bulletin* is printed in a two-column format (column width of 80 mm) with a text area of 170×225 mm.

Text. Papers should be typed, on one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

When the paper has been accepted word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with fewer type-setting errors.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words. A short title should be provided as a running head.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or land-scape (if absolutely necessary) format. Table captions should be typed on a separate sheet.

Illustrations. Copies only of artwork should be submitted. The original illustrations should accompany the paper after acceptance and revision. Text figures, line drawings, computer-generated figures and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and captions should be typed on a separate sheet in

the following format:

Figs 23–26. Figs 23–24, <u>Urophora</u> eggs: 23, <u>U. hispanica</u>; 24, <u>U. stigma</u>. Figs 25–26, spermathecae: 25, <u>U. maura</u>; 26, <u>U. stigma</u>; scale lines = 0.05 mm.

Voucher specimens. The deposition of voucher specimens should be considered where appropriate.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 <u>in</u> Waage, J. & Greathead, D. (<u>Eds</u>) <u>Insect parasitoids</u>. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) <u>Ecological methods with</u> particular reference to the study of insect populations. 2nd edn. 524 pp. London, Chapman & Hall

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, Metopolophium dirhodum (Walker) (Hemiptera: Aphididae), in the UK. Bulletin of Entomological Research 79, 373–380.

Citation of authors in the text should appear in the form: Polaszek (1990) or (Polaszek, 1990). More than one author should be cited in chronological order as: (Holloway et al., 1987; Walker & Huddleston, 1988).

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. Three copies of the manuscript and artwork should be submitted to:

The Editors
Bulletin of Entomological Research
International Institute of Entomology
56, Queen's Gate
London
SW7 5JR, UK.

MARCH 1993 Vol. 83(1)

Bulletin of Entomological Research

Guest Editorial: Schofield, C.J. The politics of malaria vector control
Chiasson, H. & Hill, S.B. Population density, development and behaviour of <i>Diopsis longicornis</i> and <i>D. apicalis</i> (Diptera: Diopsidae) on rice in the Republic of Guinée
Cudjoe, A.R., Neuenschwander, P. & Copland, M.J.W. Interference by ants in biological control of the cassava mealybug <i>Phenacoccus manihoti</i> (Hemiptera: Pseudococcidae) in Ghana
Daly, J.C. & Fisk, J.H. Expression of pyrethroid resistance in adult <i>Helicoverpa armigera</i> (Lepidoptera: Noctuidae) and selective mortality in field populations
Hargrove, J.W. & Packer, M.J. Nutritional states of male tsetse flies (Glossina spp.) (Diptera: Glossinidae) caught in odour-baited traps and artificial refuges: models for feeding and digestion
Hobbs, S.E. & Hodges, G. An optical method for automatic classification and recording of a suction trap catch
Holter, P., Sommer, C., Grønvold, J. & Madsen, M. Effects of ivermectin treatment on the attraction of dung beetles (Coleoptera: Scarabaeidae and Hydrophilidae) to cow pats
Ito K, Sugiyama, H., Nik Mohd. Noor b. Nik Salleh & Min, C.P. Effects of lunar phase on light trap catches of the Malayan black rice bug, Scotinophara coarctata (Heteroptera: Pentatomidae)
Khayrandish, A. & Wood, R.J. Organophosphorus insecticide resistance in a new strain of Culex quinquefasciatus (Diptera: Culicidae) from Tanga, Tanzania
Khayrandish, A. & Wood, R.J. A multiple basis for insecticide resistance in a strain of Culex quinquefasciatus (Diptera: Culicidae) from Muheza, Tanzania, studied as resistance declined
Kluge R.L. & Caldwell P.M. The biolopgy and host specificity of Pareuchaetes aurata aurata (Lepidoptera: Arctiidae), a 'new association' biological control agent for Chromolaena odorata (Compositae)
Miller, R.H., El Masri, S. & Al Jundi K. Plant density and wheat stem sawfly (Hymenoptera: Cephidae) resistance in Syrian wheats
Mills, N.J. Observations on the parasitoid complexes of budmoths (Lepidoptera: Tortricoidea) on larch in Europe
Moore, R., Clarke, R.T. & Creer, S. An insect sorting device to be used in conjunction with insect suction samplers
Passerini, J. & Hill, S.B. Field and laboratory trials using a locally produced neem insecticide against the Sahelian grasshopper, Kraussaria angulifera (Orthoptera: Acrididae), on millet in Mali
Peiris, H.T.R. & Hemingway, J. Characterization and inheritance of elevated esterases in organophosphorus and carbamate insecticide resistant Culex quinquefasciatus (Diptera: Culicidae) from Sri Lanka
Pleass, R.J., Armstrong, J.R.M., Curtis, C.F., Jawara, M. & Lindsay, S.W. Comparison of permethrin treatments for bednets in The Gambia
Book Reviews

© C-A-B International, 1993

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.