
SUPPLEMENT TO
“CHEMICAL AND BOTANICAL GUIDE TO LICHEN PRODUCTS”

by

Chicita F. Culberson

This supplement to the book *Chemical and Botanical Guide to Lichen Products*, published in 1969 by the University of North Carolina Press, Chapel Hill, summarizes the information from an additional 295 papers, primarily from 1965 to 1968 but including many from 1969 as well. This four-year summary of the rapidly growing field of lichen chemistry and chemosystematics is nearly one third the length of Dr. Culberson's original book that summarized all the literature of the science from its origin.

From the first reviews published of the book . . .

“Taxonomists will find this book a gold mine of information that will take years to work over.”—MASON E. HALE, JR., *THE BRYOLOGIST* (Spring, 1970).

“The present work is . . . especially valuable since it is so authoritative and covers a series of molecules of outstanding biosynthetic and chemotaxonomic interest.”—JEFFREY B. HARBOURNE, *Phytochemistry* (1970).

“This is a good book which will probably be the standard text on the subject for some time to come.”—DAVID C. SMITH, *Nature* (March, 1970).

“This unique book . . . will be an invaluable time-saver for lichen taxonomists and investigators of natural products.”—*Choice* (Association of College and Research Libraries) (March, 1970).

“This is a remarkable book. Dr. Culberson, who is already well known as a natural products chemist, has proved herself to be a very careful scholar as well as an astute organizer of a very ungainly amount of literature.”—CECIL FOX, *Naturaliste Canadien* (1970).

“. . . ein Musterbeispiel für exakte Literaturarbeit . . . für jede botanische und chemische Bibliothek.”—G. FOLLMANN, *Excerpta Botanica* (February, 1970).

“This book is unique . . . indispensable.”—VERNON AHMADJIAN, *Quarterly Review of Biology* (June, 1970).

“. . . a landmark in the history of our knowledge of the lichen substances.”—F. A. STAFLEU, *Taxon* (June, 1970).

200 pages, paperback.—\$4.00, postpaid.

Available exclusively from the American Bryological and Lichenological Society

Send orders with payment to

LEWIS E. ANDERSON

Department of Botany

Duke University

Durham, North Carolina 27706, USA

PART I: GENERAL INDEX*

- Acanthomyops niger*, soredia on, 256
Acer, Lichens on, 305
 — *pseudoplatanus*, lichens on, 145, 156, 157
 —, pH of bark, 341
 Acid rock crevices and overhangs, lichens of, 309–322
Ahmadjian, V., 234–247, 259–267, 268–270
 Air pollution and lichens, 105, 106, 110–113, 114, 117, 142, 149, 172
 —, in relation to lichens and fungi, 337–349
 Algae, carbohydrate release, 330–331
 —, transfer, 327
 —, carbon dioxide fixation, 330
 —, presence or absence in fungi, 350–362
 —, starch levels, 330
Alnus, lichens on, 152
Alternaria sp., 342
 Aluminium ions, intolerance of, 4
 Amino acids in lichens, 271–282
 —, quantitative determination, 283–288
Ammophila, lichens on, 7, 12
 Analysis methods for lichens, 272
 Animals, dispersal of soredia by, 256
 Ant, Soredia on, 256
 Anthropogenic factors and lichens, 116–118
 Apothecia, 357
 —, discharge of ascospores from, 57–65
Armeria, lichen on, 317
 Ascocarp structure, 350–362
 Ascomyces, 350–362
 Ascospores, discharge of, 57–65
 —, excretion by rotifers, 74–75
 —, germination of, 74–75
 Ascus, bitunicate, 354, 355
 —, dehiscence mechanisms, 357
 —, structure, 350–362
 —, unitunicate, 354, 355–357
 Atranorin, 106, 135, 162
 —, formula, 253
 Bailey, R. H., 57–65, 73, 256
 Barbatic acid, Formula, 253
Betula, Buffer capacity, 341
 —, lichens on, 73, 224
 Blakeney Point, lichens of, 1–15
 Book Review, *Chemical and Botanical Guide to Lichen Products* (Culberson), 370–371
 —, *The Lichen Genus Cladonia in North America* (Thomson), 196–197
Botrytis cinerea, 342
 Brenan, J. P. M., 257
 British Mycological Society, 323
 Brittany coast, oil pollution, 55–56
 Brown, D. H., 1–15
 —, R. M., 1–15
 Calcicole plants, 4, 9
 Calcium-requiring species, 4
Calothrix, 96, 97
 —, carbohydrate release, 331
 Carbohydrate in lichens, 331
 Carbon dioxide fixation in algae, 330
Ceratostomella ulmi, resistance to SO₂, 342
Cercospora beticola, 339
 Chemistry of lichens, 248–255, 271–282, 283–288
 Chemotypes, 254
Chiodectin, 194
Chiodectonic acid, 194
Chloramine-T solution, 249
Chodat's nutrient agar, 289
 Chromatographic properties of lichen substances, 249
 Chromatography methods, 272
Cladophora, 141
 Clean-air and return of lichens, 110
 Climate, differences of, 4–5
Coccomyxa, 216, 330
 —, carbohydrate release, 331
 Cold resistance of phycobionts, 289–293
 Communities of lichens, 363
Acarosporion fuscatae, 105, 111, 173, 177–178
Aspicilietum cinereum, 173, 177
Buellietum punctiformis, 173, 176
Caloplacetum heppianae, 105, 111, 173, 174, 175, 178, 179, 180
 — *murorum*, 173, 180, 181
 — *phloginae*, 173, 174
Caloplacion decipientis, 173, 190
Candelariellum corallizae, 298
Chaenothecetum, 173, 179
Cladonietum coniocraeae, 173, 179
Cladonion coniocraeae, 173, 179
Collematetum tuniformis, 173, 180–181
Collemation tuniformis, 173, 180–181
Conizaeoidion, 173–174, 179
Dirinetum repandae, 194
Graphidion, 182
Gyalectetum jenensis, 173, 180
Lecanoretum calcareae, 173, 174, 180
 — *dispersae*, 105, 173, 179–180
 — *pityreiae*, 105, 111, 173–174, 175, 176, 178
Lecanorion calcareae, 173, 180
 — *dispersae*, 173, 179–180
Lecideetum crustulatae, 173, 178
Leprarietum, 173, 179
Leprarion, 173, 179

*Compiled by S. A. Manning, F.L.S., L.I.BIOL Registered Indexer of the Society of Indexers.

- Parmelietum furfuraceae*, 105, 115, 173, 175, 176–177, 178, 182
 — *omphalodis*, 173, 177, 178–179
Physciaetum caesiae, 173, 175
Physodion, 111, 142, 176–177, 179, 345
Psoretum ostreatae, 173, 174
Ramalinetum fastigatae, 115, 142, 173, 175–176, 179
Roccellion oceanicum, 194
*Umbilicariion cylindrica*e, 105, 168, 173, 177, 178–179
Xanthorion, 173, 174–176, 180
- Competition and lichen distribution, 5, 71
 Confluentic acid, 194
 Controlled environment lichen growth chamber, 83–87
Corticium centrifugum, 145
Corylus avellana, lichens on, 66–72, 78, 224
 Cover analysis of lichens, 365–366
Crataegus, Lichens on, 115, 129, 131, 145, 256
 — *monogyna*, lichens on, 174
 Cryptogamic collections at Kew and British Museum, 257
 Culture of lichens, 259–267
- Damage to lichens by pollutants, 339
 Decarboxythamnolic acid, chromatographic properties, 249
 —, formula, 253
 Decontamination measures and lichens, 55–56
Dichothrix sp., 88, 94, 95
 — *baueriana*, 90, 92, 93
 Dickinson, C. H., 66–72
Diplocarpon rosae, 339, 342
 — and SO₂, 343–344
 Discharge of ascospores, 57–65
 Distribution records of lichens:
 Africa, 21–30
 Antarctica, 24
 Argentina, 22, 26, 28–30, 88–98, 290–291
 Australia, 23, 27, 30, 88–98
 Bolivia, 22, 27, 29, 30
 Brazil, 23, 28
 British Isles, 34–54, 184–185, 218–233, 248–255, 297–308, 309–322
 Aberdeenshire, 207–208, 251–252 ;
 Angus, 184–186, 214, 250–251 ;
 Argyllshire, 248, 251 ; Berkshire, 184, 186, 298 ; Buckinghamshire, 230 ; Caernarvonshire, 208, 250–251 ; Cardiganshire, 251 ; Carmarthenshire, 184, 186 ; Cornwall, 55–56 ; Cumberland, 227, 250 ; Derbyshire, 105–193, 251–252 ; Devon, 3–5, 8–10, 47, 250, 298 ; Dorset, 3–5, 10, 227 ; Dumbarton, 298 ; Dumfries, 298 ; Durham, 250 ; Easterness, 250–251 ; East Lothian, 250 ; Glamorgan, 74–75 ; Gloucestershire, 256 ; Hampshire, 368 ; Hertfordshire, 230 ; Ireland, 184, 186 ; Antrim, 298 ; Clare, 66–72 ; Cork, 227, 230 ; Down, 52, 298 ; Galway, 184, 186 ; Kerry, 78, 81, 208 ; Leitrim, 44–45 ; Tyrone, 298 ; Wicklow, 251 ; Isle of Man, 230 ; Kent, 184, 186 ; Kincardine, 299 ; Kirkcudbrightshire, 251 ; Lancashire, 250 ; Leicestershire, 184–185, 252, 303 ; Lincolnshire, 183–184 ; London, 184, 186 ; Merioneth, 208, 250, 253, 303 ; Mid Lothian, 45 ; Montgomeryshire, 251 ; Mull, 227, 298 ; Norfolk, 1–15, 302 ; Northumberland, 184–185, 251–252 ; Pembrokeshire, 227 ; Perthshire, 208, 227, 250–251 ; Ross-shire, 250–252 ; Shetlands, 184, 186 ; Shropshire, 250, 253 ; Somerset, 10, 47, 53, 184, 186 ; Suffolk, 184, 186 ; Surrey, 230, 252 ; Sussex, 134, 230 ; Sutherland, 42, 250–251 ; Westerness, 47, 250–251, 298 ; Westmorland, 181, 208 ; Wiltshire, 298, 303–304 ; Worcestershire, 182 ; Yorkshire, 42, 50, 250–252
 Canada, 99–104, 184, 209, 211–212
 Canary Islands, 77–82
 Chile, 22, 77–82
 China, 23–24
 Colombia, 77–82
 Crete, 184
 Finland, 35, 205, 207–208, 303
 France, 26, 55–56, 227, 291, 301, 303, 305, 307
 Germany, 42, 184, 290–291, 303, 305
 Greenland, 294–296
 Iceland, 73, 184
 India, 184
 Iran, 28
 Italy, 290
 Juan Fernandez Islands, 77–82
 Mexico, 21–22, 24, 29
 Morocco, 26
 New Zealand, 229
 Norway, 205, 207–208, 271–282, 283–288
 Spain, 23–24, 28, 290–291
 Sudan, 25
 Sweden, 206–208, 227, 299–300, 301
 Switzerland, 27, 73–74, 294
 U.S.A., 21–30, 77–82, 88–98, 184, 209–212, 223, 225, 229, 260, 271
 Venezuela, 23
 West Indies, 24, 77–82
 Yugoslavia, 194
- Divaricatic acid, 106, 156
 —, chromatographic properties, 249
 —, formula, 253
 Dobbs, G., 323–325
 Domí scale, 173
 Dune systems, 3–5
- Ecological changes in lichen populations, 337

- Environmental requirements of lichen, 83–87
Erysiphe graminis, sucrose absorption, 332
Erythrin, 194–195
 Extraction procedure, 272
- Fagara mayu*, lichen on, 78
Fagus, lichens on, 132, 171, 224, 301–302
 — *sylvatica*, lichens on, 175
 Field studies in lichenology, 363–369
 Fir, lichens on, 145
 Fires and lichens, 113, 115, 152
 Follmann, G., 194–195
Fraxinus, lichens on, 224
 — *excelsior*, lichens on, 125–126, 132, 142, 145–146, 148, 155–158, 175–177, 179, 341
 —, pH of bark, 341
 Fresh-water species of *Verrucaria*, 34–54
 Fruiting of *Cladonia*, 7, 9
 Fumarpotocetaric acid, 106, 125, 132, 135
 Fungi, lichenized and non-lichenized genera, 350–362
 —, phycotrophic, 323–325
 —, polyols in tissues infected by, 333
 —, symbiotic associations with other plants, 326–336
- G.A.O-T solution, 249
 Garrett, R. M., 57–65
 G.E. solution, 249
Gloeocapsa, 52
 Glomelliferic acid, 304
 Growth chamber for lichens, 83–87
 —, rates in lichens, 32–33
 Gyrophoric acid, 21, 23
- Haematomic acid, 249
 Harris, R. C. 77–82
 Hawksworth, D. L., 105–193, 248–255, 370–371
 Haynes, F. N., 363–369
 Heikkila, H., 259–267
 Henssen, A., 88–98, 99–104
Heterococcus caespitosus, 37
 Hormocystangia in *Lempholemma*, 99–104
 Huneck, S., 194–195
Hyalococcus, Carbohydrate release, 331
 — *dermatocarponis*, 266
 Hypoprotocetaric acid, 106, 163
Hysterium pulicare, 342
- Ilex*, lichens on, 224
 — *aquifolium*, lichens on, 80
 Imshaug, H. A., 77–82
 Interascal tissue, 353–354
 Introduction of lichens by man, 73
 Isidia as vegetative propagules, 214–217
- Jacobs, J. B., 268–270
 Jahns, H. M., 199–213
 James, P. W., 309–322
- Kappen, L., 289–293
 Kershaw, K. A., 83–87, 214–217
- Key to *Candelariella*, 299
 — *Verrucaria* (fresh-water species), 40
 — lichens of shaded acid rock crevices, 311–321
 — *Arthopyrenia* (sect. *Acrocordia*), 221
 — *Pilophorus*, 204
 Kinraide, W. T. B., 234–247
- Lamb, I. M., 73–74
 Lange, O. L., 289–293
Larix, lichens on, 142
 Laundon, J. R., 196–197, 297–308
 Leaching, rate of, 4
 Lead mining, lichens on spoil tips, 117, 132
 Lecanoric acid, 194, 234
Leptiphyphantes tenuis, 256
 Lewis, D. H., 326–336
 Lichen products, review of guide to, 370–371
 Lichenicolous lichen, 303
Lichenonconium lecanoracearum, 145
 Light (spore discharge and, 64
Limonium, lichens on, 11, 14
Lycopodium spores, trapping of, 256
- Magnusson, A. H., *Acarospora* studies, 16–31
 Mazaedia, 357, 360
 Measurement of lichen growth rates, 32–33
 Medulla as storage organ, 346
Melampsora cerastii, 342
 — *lini*, 334
 Melzer's reagent, formula, 39
 Microclimate and epiphyte distribution, 66–72
Microcoleus chthonoplastes, 10
 Microcrystal tests, 249
Microsphaera alphitoides, 342
 Millbank, J. W., 83–87, 214–217
 Molluscan slime, soredia on, 256
 Monosaccharides, quantitative determination, 283–288
 Morgan-Huws, D. I., 363–369
 Mortality of lichens, 55–56
 Mycorrhizas, 328–329
Myrcenugenia schulzei, lichens on, 80
Myrmecia, carbohydrate release, 331
 — *pyriformis*, 37
- Ninhydrin-positive compounds in lichens, 271–282
 Nitrogen in lichens, 273, 286–287
 Nitrogen dioxide and lichens, 339
 Norstictic acid, 194
Nostoc, 10, 99, 101–103, 201, 330
 —, carbohydrate release, 331, 346
 — *muscorum*, CO₂ fixation, 330
Nothomyrcia fernandeziana, lichens on, 80
- Oil pollution and lichens, 55–56
 Olivetoric acid, 106, 162
 Ontogeny of *Pilophorus*, 199

- Paper partition chromatography, 249
 Paraphyses, 353
Paraphysoides, 353–354
 Paraphysoid tissue, 353–354
Parietin, 129, 131
 Periphyses, 353
Petrithecia, 357
Persea, lichens on, 80
Philodina (rotifer) on lichen, 74–75
 Photophobous lichens, 309–322
Phragmidium spp., 342
 Phycobiont, cold resistance, 289–293
 —, pollutant damage, 347
 Physodic acid, 162
Phytophthora infestans, SO₂ and, 343, 345
Pinus sylvestris, lichens on, 225
 —, pH of bark, 341
Pleurococetum vulgaris, 173–174
Pleurococcus viridis, 173–174
 Pollutants, atmospheric, 338
 —, removal of, 83–87
 —, water, 182
 Pollution studies, zoning criteria in, 340–342
 Polysaccharides of lichens, 283–288
Populus, lichens on, 6, 11, 12, 14, 224
 — *alba*, lichens on, 301–302
 — *tremula*, lichens on, 278, 301–303
 Precipitation and spore discharge, 64
 Protein fraction of lichen tissue, 283–288
Protocetraric acid, 163
Protococcus viridis, 176
 Pseudoparaphyses, 353–384
Pseudopleurococcus incrustans, 37
 Psoromic acid, 194
Puccinia graminis, 334
 — *poarum*, 328–329, 331–332
 — *striiformis*, 328
 Pyatt, F. B., 74–75
 Pyrenocarpous lichens, 34–54, 218–233

Quercus, lichens on, 115, 125, 128, 131–132, 142, 145–146, 151–152, 154, 156, 158–159, 171–172, 179, 224, 341
 — *robur*, lichens on, 175, 177, 179, 368
 — —, pH values of bark, 367–368

 Ranwell, D. S., 55–56
 Relative humidity and lichens, 113
 Reproduction in *Cladonia*, 7
 Rhizocarpic acid, 20
Rhizoctonia tuliporum, sensitive to SO₂, 342
 Richardson, D. H. S., 350–362
 Rock climbing and lichens, 117
 Ross, R., 257
 Rotifers on lichen, 74–75

Saccharomyces ellipoideus and sulphurous acid, 343
 Salazinic acid, 155
Salix, lichens on, 128, 146, 151, 154, 156, 158
Samucus, lichens on, 115, 145, 159, 172
 Saunders, P. J. W., 337–349

Sclerotium delphinii, sensitive to SO₂, 342
Scytonema, carbohydrate release, 331
 Shaded acid rock crevices and overhangs, lichens of, 309–322
 Sociology of lichens, 105, 111, 115, 142, 168, 172–183
 Soil-type and lichens, 113–114
 Solberg, Y. J., 271–282, 283–288
Sorbus aucuparia, lichens on, 171
 Soredia, dispersal by animals, 256
 Sørensen coefficient, 183
 Spiders' web, soredia on, 256
 Spore-to-spore synthesis, 259–267
 Squamules, ultrastructure, 268–270
 Starch levels in lichen algae, 330
Stichococcus pyrenoglobuli, 269
 — *diplosphaera*, 259, 261, 268
 — *mirabilis*, 259, 261–262
 Stigmatidin, 194
Stigonema, 201
Suaeda, lichens on, 6, 7, 10–14
 Substratum stability, 4–5
 Sugars in lichens, 271–282
 Sugar alcohols in lichens, 271
 Sulphur requirement of lichen, 73
 — dioxide, 338–347
 —, experimental exposure of lichens and fungi to, 342
 Swinscow, T. D. V., 34–54, 218–233, 294–296
 Symbiotic associations between fungi and other plants, 326–336
 — fungi, carbohydrate nutrition of, 327–334
 Synthesis of lichens, 259–267

 Taxonomic revision of *Acarospora*, subgenus *Xanthothallia*, 16–31
 Taxonomy, integration for lichenized and non-lichenized fungi, 350–362
 — of *Pilophorus*, 199–213
 Temperature and spore discharge, 64
 Thamnolic acid, 137
 —, chromatographic properties, 249
 —, formula, 253
Thelycrania, lichens on, 156
 Thermal damage to lichens, 56
 Thorp, T. K., 66–72
 Topham, P. B., 294–296
Torrey Canyon oil and lichens, 55–56
 Trampling of lichens, 138
Trebouxia as phycobiont, 329–330
 —, carbohydrate release, 331, 346
 —, pyrenoglobuli in, 269
 —, SO₂ and, 345
 —, usnic acid and, 234–247
Trematosphagaria, 342
Trentepohlia, 79, 141, 220, 232
 —, carbohydrate release, 331
Tsuga, lichens on, 153
Tussilago farfara, 328–329, 331–332

Ulmus, lichens on, 128, 132, 140, 142, 145–146, 154, 156–158, 174–176, 224, 305

- *glabra*, lichens on, 176
 Ultrastructure of lichens, 268–270
Uromyces fabae, 328
 — *phaseoli*, 328, 339
Usnic acid, 106, 156, 163
 —, chromatographic properties, 249
 —, effects on *Trebouxia*, 234–247
 —, formula, 253
 —, properties, 234
 Vegetative propagules, 214–217
- , reproduction in *Cladonia*, 7
Ventosic acid, 249
- Water, availability, 4–5
Weber, W. A., 16–31
Woolhouse, H. W., 32–33
- Xerophytic lichens, 309
- Zygosaccharomyces prioranus* and SO₂, 343

PART II: LICHEN INDEX

*Denotes a new combination, a new rank, or a new name.

†Denotes a new taxon.

- Acarospora albida*, 21; *A. amabilis*, 21–22; *A. angolensis*, 21; *A. austro-africana*, 22; *A. bella*, 22, 28, 30; *A. boliviiana*, 22, 28; *A. brouardii*, 22; *A. calviniensis*, 22; *A. catamarcae*, 22; *A. chilensis*, 22; *A. chlorophhana*, 16–31; *A. chrysops*, 22–23, 28–29; *A. citrina*, 23; *A. contigua*, 23; *A. deserticola*, 23; *A. discurrents*, 23; *A. dissipata*, 23; *A. dissipata* var. *separabilis*, 23; *A. epilutescens*, 23; *A. epithallina*, 23; *A. erythrophora*, 23; *A. evoluta*, 23; *A. extenuata*, 24; *A. finkei*, 22, 24, 28; *A. fuscostata*, 5, 124, 177, 178, 235, 237–245, 340; *A. gobiensis*, 24; *A. guadeloupensis*, 24; *A. gwynnii*, 24; *A. heppii*, 5, 124; *A. heuffleriana*, 24–26; *A. hilaris*, 19, 24, 27; *A. immixta*, 24; *A. incertula*, 25; *A. initialis*, 25, 29; *A. initialis* var. *perfectior*, 25; *A. insculpta*, 25; *A. intercedens*, 25; *A. kordofanica*, 25–26; *A. laeta*, 25; *A. laeta* var. *annularis*, 25; *A. laevigata*, 26; *A. lavicola*, 26; *A. lucida*, 26; *A. luderitzensis*, 26–28; *A. malmeana*, 25–26; *A. maroccana*, 26; *A. massiliensis*, 24, 26; *A. meridionalis*, 26–29; *A. microcarpa*, 26; *A. microcarpa* var. *dispersella*, 26; *A. negligens*, 27; *A. novae-hollandiae*, 22, 27; *A. novomexicana*, 23, 25, 27–28, 30; *A. ocellata*, 27; *A. ochrophana*, 27; *A. oculata*, 27; *A. örtendahlii*, 27; *A. oxytona*, 25, 27; *A. oxytona* f. *rugulosa*, 27; *A. oxytona* var. *substerilis*, 28; *A. placidiiformis*, 28; *A. punae*, 28; *A. radicata*, 28; *A. regnellianum*, 22, 28; *A. rhodesiae*, 28; *A. rimulosa*, 28; *A. rubicunda*, 28; *A. samoensis*, 28; *A. schleicheri*, 16–31; *A. scrobiculata*, 25; *A. smaragdula*, 187; *A. smaragdula* var. *lesdainii*, 5, 124; *A. socialis*, 25, 28; *A. staphiana*, 28; *A. steineri*, 24, 28–29; *A. subalbida*, 29; *A. subcontigua*, 29; *A. subcontigua* var. *invadens*, 29; *A. subtarsa*, 28–29; *A. sulphurata*, 19, 22–23, 25, 29; *A. sulphurata* f. *inornata*, 29; *A. tenuis*, 21, 26–27, 29; *A. terestriss*, 29; *A. texana*, 29; *A. theleomma*, 29; *A. viridula*, 30; *A. weldenii*, 30; *A. wilsonii*, 27, 30; *A. xanthophhana*, 29–30; *A. xanthophhana* f. *tenuior*, 30
- Acrocordia alba*, 223; *A. biformis*, 228; *A. biformis* var. *conformis*, 228; *A. conformis*, 228; *A. conoidea*, 221; *A. conoidea* var. *macrospora*, 225, 227; *A. conoidea* var. *salweii*, 225; *A. epipolaea*, 221; *A. Garovagliai*, 221; *A. gemmata*, 223; *A. gemmata* var. *sphaerooides*, 223; *A. glauca*, 223–224; *A. ligustica*, 221–223; *A. macrospora*, 221–223; *A. monensis*, 230; *A. polycarpa*, 228; *A. Salweii*, 225; *A. scotofora*, 228; *A. sphæroides*, 223; *A. tersa*, 228
- Alectoria bicolor*, 185; *A. capillaris*, 273, 275; *A. chalybeiformis*, 124; *A. fremontii*, 125; *A. fuscescens*, 112, 124–125, 273, 277, 284–285, 287; *A. fuscescens* var. *albosorediosa*, 125; *A. jubata* complex, 6, 125; *A. lanestris*, 125, 185; *A. nadvornikiana*, 285, 287; *A. nigricans*, 273, 277, 281, 285, 287–288; *A. nigricans* f. *subchalybeiformis*, 125; *A. ochroleuca*, 273, 277, 281, 284–285, 287, 304–305; *A. pubescens*, 178, 185; *A. sarmentosa*, 273, 277, 285–286; *A. sarmentosa* subsp. *sarmentosa*, 291 (cold resistance)
- Anaptychia ciliaris*, 6, 125; *A. fusca*, 56; *A. fusca* (chemistry), 271, 273, 277–279, 281; *A. leucomelaena* (cold resistance), 290
- Anthracothecium hibernicum*, 77–82; *A. pyrenuloides*, 77–82
- Arthonia* sp. in dry crevices, 311; *A.*

- arthonioides*, 318 ; **A. atlantica* P. James nom. nov., 318 ; *A. cinnabarinata*, 68–69 ; *A. dendritica*, 318 ; *A. lobata*, 309, 311 ; *A. lurida*, 125 ; *A. punctiformis*, 125 ; *A. radiata*, 68–69, 125 ; *A. spadicea*, 125
- Arthopyrenia alba*, 125, 223 ; *A. antecellans*, 126 ; *A. argilospora*, 126 ; *A. biformis*, 228–230 ; *A. caesia*, 126, 231–232 ; *A. conformis*, 228 ; *A. conoidea*, 126, 186, 221–223 ; *A. conoidea* var. *suzae*, 223, 228 ; *A. epipolaea*, 221 ; *A. fallax*, 68–69, 126, 355 (ascus illustrated) ; *A. gemmata*, 223–225, 230 ; *A. gemmata* var. *glauca*, 224 ; *A. glauca*, 224 ; *A. halodystes*, 55–56 ; *A. ligustica*, 221 ; *A. macrospora*, 221 ; *A. monensis*, 230–232 ; *A. punctiformis*, 68, 126 ; *A. salweyi*, 225–228 ; *A. saxicola*, 126, 232 ; *A. sphaeroides*, 223 ; *A. subglobosa*, 228 ; *A. tera*, 228
- Aspicilia phaeops*, 321
- Astrothelciaceae*, first report from Europe, 77–82
- Bachmannia maurula*, 50
- Bacidia* species, 182 ; *B. ciurinella*, 126, 187 ; *B. herbarum*, 313 ; *B. inundata*, 6 ; *B. lignaria*, 14, 126 ; *B. melaena*, 126 ; *B. muscorum*, 6, 126–127 ; *B. rubella*, 127 ; *B. sabuletorum*, 127 ; *B. scopulicola*, 313 ; *B. violacea*, 217
- Baeomyces acicularis*, 208–210 ; *B. roseus*, 127 ; *B. rufus*, 113, 127, 182 ; *B. rufus* f. *subsquamulosus*, 127
- Belonia russula*, 316
- Biatorella moriformis*, 14
- Biatorina chalybeia*, 7 ; *B. lenticularis*, 7
- Botrydina vulgaris*, 127, 182
- Bryopogon prostratostoeulos*, 125
- Buellia aethalea*, 6 ; *B. alboatra*, 127, 310, 317 ; *B. alocizoides*, 164 ; *B. canescens*, 6, 128, 175–176, 187 ; *B. chlorophaea*, 317 ; *colludens*, 13 ; *B. confervoides*, 13 ; *B. epipolia*, 180 ; *B. punctata*, 6, 128, 174–176, 341 (in built-up area) ; *B. myriocarpa*, 6 ; *B. spuria*, 6 ; *B. stellulata*, 6, 128
- Calicium abietinum*, 128 ; *C. hyperellum*, 6 ; *C. viride*, 6, 128
- Caloplaca*, 6–7, 183 ; sect. *Gasparrina*, 175, 180 ; *C. arnoldii*, 128 ; *C. atroflava*, 6 ; *C. aurantia*, 58–61, 63, 128, 181 ; *C. aurantiaca*, 6, 128, 181 ; *C. cerina*, 129 ; *C. chalybea*, 129 ; *C. cirrochroa*, 129 ; *C. citrina*, 6–7, 129, 312 ; *C. citrina* f. *citrina*, 129, 181 ; **C. citrina* forma *phlogina* (Ach.) Hawksworth n. comb., 129, 174 ; *C. citrina* var. *phlogina*, 129 ; *C. decipiens*, 7, 175, 187 ; *C. ferruginea*, 7, 129 ; *C. heppiana*, 7, 128, 129–130, 181, 344 (in built-up area) ; *C. holocarpa*, 6–7, 130, 174, 180 ; *C. incrassata*, 6–7 ; *C. lactea*, 130, 180 ; *C. littorea*, 309, 312 ; *C. luteoalba*, 7, 130 ; *C. marina*, 6–7, 55–56, 186 ; *C. murorum*, 14, 128, 130, 174, 180–181 ; *C. oblitterans*, 312 ; *C. ochracea*, 130 ; *C. phlogina*, 129 ; *C. stillicidiorum*, 14, 129–130 ; *C. tetrica*, 130 ; *C. thallincola*, 56 ; *C. trachyphylla*, 21 ; *C. variabilis*, 131 ; *C. xantholyta*, 131
- Candelaria concolor*, 14
- Candelariella*, key to British species, 299 ; *C. aurella*, 7, 131, 179 ; *C. coralliza*, 131, 297–299 ; *C. medians*, 131, 174–175 ; *C. pulvinata*, 297–299 ; *C. superdistans*, 303 ; *C. vitellina*, 7, 131, 174, 177, 182 ; *C. vitellina* var. *assericola*, 131 ; *C. vitellina* f. *corruscans*, 131 ; *C. vitellina* var. *pulvinata*, 297–299.
- Catillaria chalybeia*, 7, 310, 317, 321 ; *C. graniformis*, 14 ; *C. griffithii*, 7, 131 ; *C. lenticularis*, 7, 131 ; *C. lightfootii*, 185 ; *C. litorella*, 320–321 ; *C. pulverea*, 132
- Catocarpone*, sect. of *Rhizocarpon*, 73–74.
- Cetraria aculeata*, 10 ; *C. aculeata* f. *acanthella*, 10 ; *C. chlorophylla*, 132, 178 ; *C. cucullata*, 273, 284–285, 287 ; *C. delisei*, 273 ; *C. glauca*, 7, 132, 160, 340 (in built-up area) ; *C. islandica*, 113, 132, 187, 273, 275, 277, 281 ; *C. islandica* f. *islandica*, 132 ; *C. islandica* f. *platyna*, 132 ; *C. nivalis*, 273–274, 285, 287, 291 (cold resistance) ; *C. richardsonii*, 271, 273, 277, 281 ; *C. sepincola*, 358 (asci illustrated)
- Chaenotheca ferruginea*, 132, 179
- Chiodescon albidum*, 314, 317 ; *C. cretaceum*, 194–195 (erythrin in) ; *C. myrticola*, 309, 314, 317 ; *C. petraeum*, 309, 314–316 ; *C. sanguineum*, 194 ; *C. sphaerale*, 194 ; *C. stalactinum*, 194
- Cladonia*, sugar absorption, 332 ; *C. alcidonis*, 3, 4, 8 ; *C. alpestris*, 7 ; *C. arbuscula*, 3, 8–10, 113, 132, 197, 273–274 and 284 (chemistry) ; *C. arbuscula*, subsp. *arbuscula* chem. str. 11, 197 ; *C. arbuscula* subsp. *beringiana*, 197 ; *C. bacillaris*, 8, 133 ; *C. bellidiflora*, 273, 284 (chemistry) ; *C. boryi*, 235, 238–243, 245 ; *C. caespiticia*, 8, 133 ; *C. cervicornis*, 133 ; *C. chlorophaea*, 8, 9, 133, 179, 197 ; *C. chlorophaea* f. *lepidophora*, 8 ; *C. coccifera*, 14, 133 ; *C. contiocraea*, 8, 133, 175, 179 ; *C. conista*, 8, 9, 133, 197 ; *C. convoluta*, 291 (cold resistance) ; *C. cornuta*, 134 ; *C. deformis*, 134, Chemistry: 273, 284 ; *C. degenerans* f. *anomoeus*, 134 ; *C. digitata*, 134 ; *C. fimbriata*, 8, 9, 58, 134, 179 ; *C. fimbriata* var. *major*, 8 ; *C. fimbriata* var. *subcornuta*, 10 ; *C.*

- fimbriata* var. *subulata*, 10 ; *C. floerkeana*, 8, 134, 182 ; *C. floerkeana* var. *carcata*, 8 ; *C. foliacea*, 3, 4, 8, 291 (cold resistance) ; *C. foliosa*, 8 ; *C. furcata complex*, 7-9, 134-135 ; *C. furcata* var. *scabriuscula*, 9 ; *C. furcata* var. *spinosa*, 8 ; **C. furcata* ssp. *subrangiformis* (Sandst.) Hawksworth n. comb., 113-114, 117, 135, 290 (cold resistance) ; *C. furcata* var. *subrangiformis*, 135 ; *C. glauca*, 8-10 ; *C. gonecha*, 134 ; *C. gracilis*, 135, 281 (chemistry) ; *C. gracilis* var. *aspera*, 14 ; *C. impexa*, 7, 9, 132, 135 ; *C. leucophaea*, 197 ; *C. macilenta*, 8, 9, 135, 182 ; *C. mediterranea*, 291 (cold resistance) ; *C. mitis*, 9 ; *C. nylanderi*, 291 (cold resistance) ; *C. ochrochlora*, 136 ; *C. pacifica*, 197 ; *C. papillaria*, 136 ; *C. pityrea*, 7, 9, 133, 136 ; *C. pityrea* f. *hololepis*, 9 ; *C. pocillum*, 9, 117, 133, 136 ; *C. polydactyla*, 134, 136 ; *C. pungegens*, 8, 9, 138 ; *C. pyxidata*, 8, 9, 133, 136 ; *C. pyxidata* var. *chlorophaea*, 8 ; *C. rangiferina*, 9, 136 ; chemistry: 273, 277, 281 ; *C. rangiformis*, 8, 9, 114, 117, 135, 137-138 ; *C. rangiformis* var. *foliosa*, 9 ; *C. scabriuscula*, 9, 137 ; *C. squamosa*, 9, 137 ; *C. squamosa* var. *allosquamosa*, 137 ; *C. squamosa* var. *squamosa*, 290 (cold resistance) ; *C. strepsilis*, 137, 187 ; *C. subcervicornis*, 137 ; *C. subrangiformis*, 9, 113-114, 117, 135 ; *C. subsquamosa*, 137 ; *C. subulata*, 8, 10 ; *C. sylvatica*, 8, 132, 197 ; *C. sylvatica* f. *alpestris*, 7 ; *C. sylvatica* f. *tenuis*, 10 ; *C. tenuis*, 10, 117, 137 ; *C. uncialis*, 137-138, 273 (chemistry)
- Clathroporina calcarea*, 138
- Collema*, 3, 10, 180 ; *C. auriculatum*, 111, 138, 181 ; *C. callospurum*, 138 ; *C. ceranoides*, 10 ; *C. crispum*, 138 ; *C. cristatum*, 181 ; *C. cristatum* var. *cristatum*, 138 ; *C. cristatum* var. *marginale*, 138 ; *C. cristatum* var. *marginale* f. *papulosum*, 139 ; *C. fragile*, 111, 139 ; *C. glaucescens*, 10 ; *C. limosum*, 10, 139 ; *C. multipartitum*, 139 ; *C. polycarpon*, 139 ; *C. subnigrescens*, 68 ; *C. tenax*, 291 (cold resistance) ; *C. tenax* var. *ceranoides*, 10 ; *C. tenax* vars. *corallinum*, *tenax*, *vulgare*, 139 ; *C. tuniforme*, 111, 139-140, 181
- Coniocybe furfuracea*, 140, 309, 312, 319
- Cora pavonia*, 290-291 (cold resistance)
- Coriscium viride*, 140
- Cornicularia aculeata*, 3, 10, 113, 117, 140 ; *C. aculeata* var. *hispida*, 140 ; *C. divergens*, 273 (chemistry) ; *C. muricata*, 10, 140 ; *C. normoerica*, 178, 186, 329-330 (phytobiont of) ; *C. tenuissima* var. *hispida* f. *muricella*, 140
- Crocynia camusi*, 305-307
- Cyphelium inquinans*, 140
- Cystocoleus*, 181 ; *C. niger*, 140-141
- Dermatocarpon fluvatile*, 141, 266 ; *D. lachneum*, 141 ; *D. meiophyllum*, 141 ; *D. miniatum*, 266, 332 (polyol uptake) ; *D. miniatum* vars. *complacatum*, *leptophyllum* and *miniatum*, 141
- Dimerella diluta*, 185 ; *D. lutea*, 185
- Diploschistes gypsaceus*, 141 ; *D. scruposus*, 177 ; *D. scruposus* var. *bryophilus*, 10, 142 ; *D. scruposus* var. *scruposus*, 141-142
- Dirina stenhammarii*, 311
- Dirinaceae*, 194
- Encephalographa cerebrina*, 142
- Endocarpon pusillum*, 142, 259-267 (culture and synthesis), 268-270 (ultra-structure)
- Enterographa crassa*, 68, 194, 318 (saxicolous form) ; *E. dendritica*, 318 ; *E. hutchinsiae*, 318 ; *E. venosa*, 194
- Ephebe brasiliensis*, 93
- Evernia prunastri*, 10, 112, 142, 174-176, 291 (cold resistance), 368 (bark acidity and), *E. prunastri* var. *stictoceros*, 10
- Fulglesia*, 131 ; *F. fulgens*, 3, 186
- Gasparrinia*, sect. of *Caloplaca*, 7, 175, 180
- Graphidales*, 195
- Graphina anguina*, 185
- Graphis elegans*, 57-59, 61, 143, 359 (asci illustrated) ; *G. scripta*, 143
- Gyalecta jenensis*, 143
- Haematomma coccineum*, 143, 300 ; *H. coccineum* var. *coccineum*, 254 ; *H. coccineum* var. *fuscocinereum*, 254 ; *H. coccineum* var. *porphyrium*, 254, 301 ; *H. elatinum*, 186 ; *H. lapponicum*, 248 ; *H. leiphaenum*, 300 ; **H. ochroleucum* (Neck.) Laundon, comb. nov., 299-301, 306 ; *H. ochroleucum* var. *ochroleucum*, 309, 312-313, 320 ; **H. ochroleucum* var. *porphyrium* (Pers.) Laundon, comb. nov., 301, 312, 314, 320 ; *H. porphyrium*, 254, 301 ; *H. ventosum*, 178, chemistry: 248-255, 273-275, 277, 281 ; *H. ventosum* subsp. *lapponicum*, 248 ; *H. ventosum* var. *typicum*, 143 ; *H. ventosum* var. *ventosum*, 143, 254
- Hydroverrucaria*, sect. of *Verrucaria*, 35
- Hypogymnia physodes* in built-up area, 340-341
- Hysteriales*, 195
- Icmadophila ericetorum*, 113, 143
- Ionopsis epulotica*, 143

- Lecanactis abietina*, 58–62, 143 ; *L. delimis*, 316 ; *L. dileniana*, 318–319 ; *L. grumulosa*, 316 ; *L. monstrosa*, 316 ; *L. plocina*, 143 ; *L. premnea*, 144 ; *L. stenhammarii*, 144
- Lecania caesia*, 314 ; *L. erysibe*, 11, 144, 320–321 ; *L. erysibe* f. *cinereofusca*, 144 ; *L. erysibe* f. *sorediata*, 144, 175 ; *L. holophaea* var. *glaucospora*, 311 ; *L. hutchinsiae*, 320–321 ; *L. leucospeirea*, 311 ; *L. nylanderiana*, 11 ; *L. prosochoides*, 11 ; *L. prosechoidiza*, 11 ; **L. rupicola* (Nyl.) P. James, comb. nov., 314
- Lecanora achariana*, 144 ; *L. actophila*, 11, 56, 186 ; *L. atra*, 11, 56, 58, 144, 182, 187, 310 ; *L. atroflava*, 6 ; *L. atrynaea*, 144 ; *L. badia*, 11, 144, 177–178 ; *L. bella*, 22 ; *L. caesia*, 314 ; *L. calcarea*, 144 ; *L. campestris*, 11, 58–59, 61–62, 144–145, 181 ; *L. chlorona*, 145 ; *L. chlorotera*, 58–59, 61, 68, 115, 145, 174–175 ; *L. cinerea*, 145, 177 ; *L. currina* f. *incrustans*, 6–7 ; *L. conizaeoides*, 11, 58–59, 61–64, 73, 105, 145, 147, 149, 173–176, 256 (soredia dispersal), 340–341 (in built-up area) ; *L. contorta*, 145–146 ; *L. coralliza*, 297–299 ; *L. crenulata*, 145 ; *L. dispersa*, 11, 58–59, 61, 145–146, 174, 179–181, 340 and 344 (in built-up area) ; *L. dispersa* f. *albescens*, 145–146 ; *L. dispersa* f. *dispersa*, 11 ; *L. distans*, 301 ; *L. epanova*, 146, 313 ; *L. expallens*, 11, 115, 146, 174–176 ; *L. ferruginea* var. *festiva*, 7 ; *L. galactina* var. *dispersa*, 11, 145–146 ; *L. gangeoides*, 146, 310 ; *L. gibbosa*, 11 ; *L. hagenii*, 11, 145–146, 174 ; *L. hagenii* f. *zosterae*, 11 ; *L. helicopis*, 11, 56 ; *L. intricata* var. *intricata*, 146 ; *L. intricata* var. *soralifera*, 146, 177–178, 313 ; *L. muralis*, 11, 146, 175 ; *L. myrinii*, 276–277 (chemistry) ; *L. pallida*, 146 ; *L. phaeops*, 321 ; *L. polytropa*, 58–59, 61–62, 146–147, 177–178, 310 ; *L. populicola*, 301–303 ; *L. praepostera*, 320 ; *L. rupicola*, 178 ; *L. subcarnea*, 320 ; *L. symmictera*, 12 ; *L. tenera*, 320 ; *L. umbrina*, 11 ; *L. umbrinofusca*, 6 ; *L. varia*, 11, 147 ; *L. ventosa* f. *laevigatum*, 250 ; *L. ventosa* f. *pallidior*, 251 ; *L. ventosa* var. *subfestiva*, 251–254 ; *L. verrucosa*, 147, 306 ; *L. verrucosa* var. *argena*, 305 ; *L. vitellina* var. *pulvinata*, 297–299 ; *L. xanthophana*, 29
- Lecidea* species, 186 ; *L. albocaerulescens*, 147 ; 310 ; *L. alocizoides*, 164 ; *L. anthropophila*, 321 ; *L. argena*, 305–306 ; *L. ascerculorum*, 11 ; *L. caesiolepra*, 314 ; *L. callicarpa*, 321 ; *L. callosyne*, 300 ; *L. cinereofusca*, 147 ; *L. coarctata*, 147 ; *L. confluens*, 147 ; *L. contigua*, 178 ; *L. crustulata*, 147, 177, 310 ; *L. cyanea*, 73–74 ; *L. cyathoides*, 310 ; *L. decipiens*, 147 ; *L. demissa*, 148 ; *L. dilutiuscula*, 321 ; *L. dubia*, 12 ; *L. erratica*, 11, 148 ; *L. exanthemitria*, 158 ; *L. expansa*, 11 ; *L. fuscoatra*, 177 ; *L. fuscoatra* var. *fuscoatra*, 148, 178 ; *L. fuscoatra* var. *grisella*, 148 ; *L. gagei*, 321 ; *L. granulosa*, 12, 113, 115, 148, 179, 182 ; *L. henricii*, 321 ; *L. hydrophila*, 148, 187 ; *L. indigula*, 321 ; *L. infidula*, 321 ; *L. jurana*, 148 ; *L. lactea*, 148 ; *L. lactea*, 73–74 ; *L. limitata*, 68, 148 ; *L. lithophila*, 148, 310 ; *L. lucida*, 105, 140, 149, 312 ; *L. lurida*, 147, 149 ; *L. macrocarpa*, 58–59, 61–62, 149, 177–178 ; *L. orosthea*, 313 ; *L. pantherina*, 73–74 ; *L. paucula*, 321 ; *L. pelobotrya*, 321 ; *L. phaeops*, 310, 321 ; *L. plana*, 149 ; *L. poliotides*, 321 ; *L. prominula*, 150 ; *L. pulveracea*, 12 ; *L. quernea*, 149, 175 ; *L. rupicola*, 314 ; *L. rusticella*, 321 ; *L. saxetana*, 301 ; *L. scabra*, 149 ; *L. scalaris*, 149, 174 ; *L. scutellata*, 149–150 ; *L. semipallens*, 321 ; *L. silacea*, 74 ; *L. speirea*, 185 ; *L. stigmatae*, 58, 150 ; *L. stigmatae* f. *subsequens*, 150 ; *L. sulphurea*, 150, 178 ; *L. sylvicola*, 150, 321 ; *L. symmicta*, 12 ; *L. taylorii*, 310, 321 ; *L. templetonii*, 150 ; *L. tenebrosa*, 150 ; *L. tumida*, 150, 181, 310 ; *L. turgidula*, 11 ; *L. uliginosa*, 12, 113, 150, 182 ; *L. viridescens*, 151
- Leiophloea biformis*, 288 ; *L. conoidea*, 221 ; *L. ligistica*, 221 ; *L. praetermissa*, 42–43 ; *L. Salweii*, 225 ; *L. scotophora*, 228 ; *L. sphaeroides*, 224 ; *L. tersa*, 228
- Lejophlebia gemmata*, 223
- Lembidium*, 219 ; *L. conforme*, 228 ; *L. polycarpum*, 228
- Lempholemma*, 97 ; *L. albonigrum*, 101, 103–104 ; *L. cladodes*, 102–103 ; *L. dispansum*, 103 ; *L. fasciculare*, 103 ; *L. fennicum*, 103 ; *L. isidioides*, 103 ; *L. minutulum*, 103 ; *L. radiata*, 103 ; *L. umbellum*, 103 ; +*L. vesiculosum* A. Henssen, n. sp., 99–104 ; **L. wrightii* (Tuck.) A. Henssen, nov. comb., 103
- Lepraria*, 131, 179 ; *L. aeruginosa*, 312 ; *L. candelaris*, 151 ; *L. chlorina*, 312 ; *L. crassissima*, 151 ; *L. incana*, 12, 105, 113, 115, 149, 151, 179, 181, 312 ; *L. leiphaema*, 299–300, 305–306 ; *L. membranacea*, 151 ; *L. xantholyta*, 151
- Leptogium*, 3, 10, 111, 181 ; *L. lichenoides*, 151 ; *L. plicatile*, 151 ; *L. schraderi*, 151 ; *L. sinuatum*, 151 ; *L. turgidum*, 152
- Leptorhaphis epidermidis*, 152
- Lichen argenus*, 305 ; *L. botryoides*, 172 ; *L. cereolus*, 206 ; *L. ciliaris*, 125 ; *L.*

- cinereus*, 172 ; *L. citrinus*, 304 ; *L. coccineus*, 299 ; *L. cornutus*, 134 ; *L. correns*, 172 ; *L. digitatus*, 134 ; *L. farinosus*, 305–306 ; *L. flavidus*, 304–305 ; *L. floridus*, 169 ; *L. fragilis*, 165 ; *L. gemmatus*, 223–224 ; *L. haematomma*, 299 ; *L. haematomma* var. *porphyrius*, 301 ; *L. hirtus*, 169 ; *L. hoffmannii*, 146 ; *L. lacteus*, 158 ; *L. metaleucus*, 223 ; *L. muscorum*, 150 ; *L. ochroleucus*, 299, 304–305 ; *L. perlatus*, 155 ; *L. resupinatus*, 152 ; *L. rigidus*, 304 ; *L. subfuscus*, 145 ; *L. subvirescens*, 304–305 ; *L. sulphureus*, 150 ; *L. ventosus*, 248 ; *L. verrucosus*, 147 ; *L. violaceus*, 127
- Lichina confinis*, 55–56, 94, 97, 186 ; *L. confinis* var. *willeyi*, 88, 90–94 ; *L. pygmaea*, 55–56, 97 ; †*L. rosulans* A. Henssen, n. sp., 89–93, 95–98 ; †*L. tasmanica* A. Henssen, n. sp., 89–98 ; **L. willeyi* (Tuck.) Henssen, nov. comb., 88–98
- Lithoicea aquatilis*, 50 ; *L. elaeomeaena* var. *alpina*, 51 ; *L. eleomelaena*, 45
- Lobaria isidiosa*, 281 ; *L. laetevirens*, 152 ; *L. pulmonaria*, 152, Chemistry: 273, 275, 277, 284–288, Cold resistance: 291 ; *L. scrobiculata*, 273, 275
- Medusolina chilensis*, 194
- Micarea*, 361
- Mycoblastus sanguinarius*, 152, 178, 359 (asci illustrated)
- Nephroma bellum*, 291 (cold resistance) ; *N. laevigatum*, 68, 152 ; *N. lusitanicum*, 68 ; *N. resupinatum*, 152
- Normandina pulchella*, 68, 186
- Ochrolechia androgyna*, 152–153 ; *O. parella*, 14, 56, 153, 158 ; *O. tartarea*, 153 ; *O. turneri*, 153
- Omphalaria wrightii*, 103
- Opegrapha atra*, 57–59, 61, 68, 153 ; *O. calcarea*, 153 ; *O. cesareensis*, 319 ; *O. confluens*, 319 ; *O. gyrocarpa*, 310, 313 ; *O. lithygra*, 310, 319 ; *O. mougeotii*, 153 ; *O. persoonii*, 153 ; *O. rimalis*, 153 ; *O. rufescens*, 153 ; *O. saxatilis*, 153, 310, 319 ; *O. saxicola*, 56, 153, 310, 319 ; *O. saxigena*, 310, 319 ; *O. vulgata*, 68–69, 154 ; *O. zonata*, 313
- Pachyphiale cornea*, 186
- Pannaria rubiginosa*, 68, 186
- Parmelia alpicola*, 283 ; *P. argena*, 305–306 ; *P. caperata*, 12, 83 (light), 154, 271 (sugars), 290–292 (cold resistance), 368 (bark acidity and) ; *P. carporrhizans*, 291 ; *P. centrifuga*, 273, 277 ; *P. ciliata*, 155 ; *P. cirrhata*, 290 ; *P. citrina* var. *phlogina*, 129 ; *P. conspersa*, 154 ; *P. delisei*, 303 ; *P. delisei* var. *isidiascens*, 303 ; *P. elegans*, 186 ; *P. exasperatula*, 340 ; *P. fuliginosa*, 12, 340 ; *P. fuliginosa* var. *laetevirens*, 12 ; *P. furfuracea*, 182 ; *P. furfuracea* var. *ceratea*, 162 ; *P. furfuracea* var. *furfuracea*, 162 ; *P. glabratula* subsp. *fuliginosa*, 12, 154 ; *P. glabratula* subsp. *glabratula*, 12, 115, 154, 175 ; *P. globellifera*, 303 ; *P. haematomma*, 306 ; *P. incurva*, 154, 178, 187 ; *P. isidiolyta*, 303–304 ; *P. loxodes*, 303–304 ; *P. omphalodes*, 154, 178 ; *P. omphalodes* var. *panniformis*, 154 ; *P. perlata*, 68, 155, 368 ; *P. perlata* var. *ciliata*, 155 ; *P. physodes*, 12, 83, 155, 182, 256, 332, 368 (bark acidity) ; *P. physodes* var. *tubulosa*, 12 ; *P. physodes* f. *vittatoides*, 155 ; *P. prolixia* subsp. *delisei* var. *isidiascens*, 303 ; *P. prolixia* var. *globellifera*, 303 ; *P. reticulata*, 155 ; *P. saxatilis*, 12, 33, 155, 177–178, 181–182, 187, 271 (sugars), 277 (chemistry), 340 (SO_2 and), 341 (in built-up area), 346 (sulphur content) ; *P. saxatilis* var. *furfuracea*, 155 ; *P. saxatilis* var. *panniformis*, 155 ; *P. subargentifera*, 304 ; *P. subaurifera*, 12, 84, 155, 175 ; *P. subfusca* var. *distans*, 301 ; *P. subrugosa*, 291 ; *P. sulcata*, 12, 83 (light), 155, 175–176, 339 (pollutants), 340 (in built-up area), 345 (SO_2 and) ; *P. taractica*, 273, 277 ; *P. tiliacea*, 156, 175 ; *P. tubulosa*, 12, 156 ; *P. ventosa* var. *cruenta*, 253 ; *P. ventosa* var. *lepadolemma*, 253–254 ; *P. verruculifera*, 303–304
- Parmeliella plumbea*, 68, 186
- Parmeliopsis ambigua*, 154, 156, 177, 273
- Parmentaria*, First European report, 77–82 ; *P. chilensis*, 77–82 ; *P. ravenelii*, 79
- Patellaria heterodoxa*, 73–74 ; *P. populicola*, 301
- Peltigera*, 4 ; *P. aphthosa*, 83–87, 156, 330 (phytociont of) ; *P. aphthosa* var. *variolosa*, 214–217 ; *P. canina*, 12, 114, 117, 156, 285–288 (chemistry), 355 (ascus) ; *P. horizontalis*, 156 ; *P. polydactyla*, 12, 68, 117, 156, 330 (phytociont), 332 (sugar absorption) ; *P. praetextata*, 64, 156–157 ; *P. rufescens*, 12, 117, 157, 332 ; *P. spuria*, 12, 117, 157 ; *P. spuria* var. *erumpens*, 12
- Pertusaria albescens*, 157 ; *P. amara*, 157, 175 ; *P. amara* var. *amara*, 157 ; *P. amara* var. *flotowiana*, 157 ; *P. corallina*, 157, 165, 177–178 ; *P. dealbata*, 157 ; *P. flavidia*, 157 ; *P. hymenea*, 147, 157 ; *P. lactea*, 158 ;

- P. pertusa*, 57-59, 61, 68, 115, 147, 158, 175, 356 (ascus); *P. pseudocorallina*, 56; *P. reducta*, 305-306
- Petractis clausa*, 158
- Phaeographis* agg., 68-69; *P. dendritica*, 69, 318; *P. inusta*, 69; *P. ramificans*, 186
- Phlyctis argena*, 158, 305-307; *P. argena* f. *erythrosora*, 305; *P. erythrosora*, 305, 307
- Physcia adscendens*, 12, 13, 56, 125, 158, 174, 176; *P. aipolia*, 158; *P. caesia*, 12, 56, 158, 175, 181; *P. ciliaris*, 6; *P. ciliata*, 158; *P. grisea*, 159, 175-176; *P. hispida*, 12, 13; *P. leptalea*, 13, 14, 159; *P. melops*, 159; *P. nigricans*, 175, 186; *P. orbicularis*, 12, 159, 176; *P. orbicularis* var. *virella*, 12; *P. pulverulenta*, 13, 159; *P. stellaris*, 159; *P. tenella*, 12, 13, 56, 125, 159, 175
- Physconia pulverulenta*, 58-59, 61
- Pilophorus*, Key to, 204; *P. acicularis*, 200-204, 206, 208-212; *P. acicularis* f. *halii*, 211; *P. cereolus*, 200-208, 211; *P. cereolus* var. *acicularis*, 208; *P. cereolus* var. *cephalodiferus*, 206; *P. cereolus* f. *cephalodiferus*, 206; *P. cereolus* var. *halii*, 208, 211-212; *P. clavatus*, 200-204, 211-212; *P. distans*, 207-208; *P. fibula*, 201-204, 209-212; *P. halii*, 211-212; *P. robustus*, 200-205, 209, 212; *P. robustus* f. *distans*, 207; *P. robustus* f. *magnus*, 204-205; *P. strumaticus*, 200-204, 206-208, 212
- Placodium atroflavum*, 6; *P. aurantiaca* var. *plicatum*, 172; *P. aurantiacum* var. *flavovirescens*, 6; *P. citrinum*, 6-7; *P. flavescentia*, 7; *P. lobulatum*, 6-7; *P. luteoalbum*, 7; *P. miniatum*, 128; *P. phlegatinum*, 6-7; *P. xanthophana*, 28
- Placynthium nigrum*, 160; *P. tantaleum*, 160
- Platismatia glauca*, 132, 160, 182
- Pleurotheliopsis*, 77-82
- Polyblastia albida*, 160; *P. cupularis*, 160; *P. gelatinosa*, 14; *P. hibernica*, 77-82; *P. schraderi*, 160; *P. tristictula*, 160
- Porina ahlesiana*, 310, 316; *P. chlorotica* var. *carpinea*, 68, 161; *P. chlorotica* var. *chlorotica*, 161, 310, 315, 355 (ascus); *P. chlorotica* f. *tenuifera*, 315; *P. curnowii*, 310, 315; *P. guentheri* var. *curnowii*, 315; *P. lectissima*, 315; *P. leptalea*, 186; *P. nucula*, 316; *P. olivacea*, 161; *P. septemseptata*, 316
- Porocyphus*, 88, 97-98; *P. lichenelloides*, 97
- Protoblastenia immersa*, 161; *P. metzleri*, 161; *P. monticola*, 14, 161; *P. rupestris* var. *calva*, 161; *P. rupestris* var. *rupestris*, 14, 58-59, 61, 161, 181
- **Pseudoevernia furfuracea* var. *ceratea* (Ach.) Hawksworth, n. comb., 162; *P. olivetorum*, 162
- Psoroichia schaeferi*, 111, 162
- Pyrenastrum astroideum*, 77-82; *P. chilense*, 77-82
- Pyrenopsis imbrida*, 52
- Pyrenula biformis*, 228; *P. gemmata*, 223; *P. gemmata* var. *sphaeroides*, 223; *P. laevigata*, 68; *P. nitida*, 68-69, 78, 162; *P. pachycheila*, 77-82; *P. polycarpa*, 228; *P. sphaeroides*, 223
- Racodium*, 181; *R. rupestre*, 141, 162
- Ramalina*, 175, 177; *R. calicaris*, 163; *R. evernioides*, 185; *R. farinacea* var. *farinacea*, 13, 112, 115, 163, 175; **R. farinacea* var. *hypoprotocetraria* (Culb.) Hawksworth n. comb., 163, 175; *R. fastigiata*, 58-59, 61, 163, 175; *R. fraxinea*, 163, 175; *R. fraxinea* var. *ampliata*, 163; *R. siliqueosa*, 55-56, 58-59, 61, 186, chemistry: 273, 275, 277, 281, 286-288
- Rehmia coeruleo-alba*, 73-74
- Rehniopsis*, sect. of *Patellaria*, 73-74
- Rhizocarpon atlanticum*, 13; *R. badio-atrum*, 178; *R. coeruleoalbum*, 73-74; *R. concentricum*, 163; *R. confervoides*, 13; *R. constrictum*, 13; *R. geographicum*, 58, 163, 177-178; *R. hochstetteri*, 13, 163; *R. obscuratum*, 178, 310, 317; *R. obscuratum* var. *reductum*, 13, 163; *R. oederi*, 185; *R. petraeum*, 163; *R. polycarpum*, 13; *R. umbilicatum*, 58-59, 61, 164
- Rinodina atrocinerea*, 164; *R. bischoffii*, 164; *R. demissa*, 13; *R. exigua*, 13, 164; *R. roboris*, 164; *R. subexigua*, 13, 164; *R. umbrinofusca*, 6
- Roccella fuciformis*, 332
- Roccellaceae*, 194
- Roccellales*, 195
- Sagedia biflora*, 228; *S. calcarea*, 221; *S. conoidea*, 221; *S. gemmata*, 223; *Sagedia* sect. *Acrocordia*, 220
- Sarcogyne regularis*, 14, 164; *S. simplex*, 164
- Schismatomma decolorans*, 311
- Sclerophyton circumscriptum*, 309, 314, 317
- Segestrella biflora*, 228; *S. gemmata*, 223; *S. sphaeroides*, 223; *Segestrella*, sect. *Acrocordia*, 220
- Segestria sphaeroides*, 223
- Silaceae*, sect. of *Lecidea*, 74
- Solenopsora candidans*, 164; *S. vulturicensis*, 311
- Solorina saccata*, 111, 165; *S. spongiosa*, 111, 165
- Sphaeriales*, 259

- Sphaerophorus fragilis*, 165 ; *S. globosus*, 165, 273–275 (chemistry), 356 (ascus) ; *S. melanocarpus*, 165
- Squamaria saxicola*, 11 ; *S. crassa*, 144, 165
- Staurothele*, 38 ; *S. caesia*, 166, 181 ; *S. clopima*, 259–267 (culture and synthesis) ; *S. hymenogonia*, 166 ; *S. rugulosa*, 265 ; *S. rupifraga*, 166
- Stenocybe septata*, 186
- Stereocaulon*, 117 (rock climbing and) ; *Stereocaulon*, sect. *Pilophoron*, 199 ; *S. acicularis*, 199 ; *S. botryosum*, 166 ; *S. condensatum*, 166 ; *S. dactylophyllum* var. *dactylophyllum*, 166–167 ; *S. dactylophyllum* var. *flabellatum*, 166 ; *S. evolutum*, 166 ; *S. fastigatum*, 166 ; *S. fibula*, 199 ; 210–211 ; *S. paschale*, 166, Chemistry: 273, 275, 277, 281–282, 284–287 ; *S. pileatum*, 166 ; *S. subcoralloides*, 167 ; *S. vesuvianum*, 167, 187, Chemistry: 273–274, 276, 282, 285–286, 288
- Sticta*, 68 ; *S. fuliginosa*, 290 ; *S. kunthii*, 290
- Stigmatidium dendriticum*, 318
- Teloschistes flavicans*, 167, 291
- Thamnidium willeyi*, 88
- Thamnodium willeyi*, 88
- Thamnolia subuliformis*, 186 ; *T. vermicularis*, 249, 252
- Thelenidia monospora*, 294–296
- Thelidium biforme*, 228 ; *T. conoideum*, 221 ; *T. decipiens*, 167 ; *T. gemmatum*, 223 ; *T. gemmatum* var. *sphaerooides*, 223 ; *T. impressum*, 167 ; *T. incavatum*, 167 ; *T. papulare*, 167 ; *T. Salweii*, 225 ; *T. tersum*, 228
- Thelocarpon epibolum*, 356
- Thelopsis melathelia*, 315
- Thelotrema lepadinum*, 185 ; *T. marginaceum*, 47 ; *T. variolarioides*, 306
- Thrombium epigaeum*, 315
- Tomasellia gelatinosa*, 68–69
- Toninia aromatica*, 167 ; *T. coeruleonigricans*, 3, 167 ; *T. lobulata*, 168
- Umbilicaria*, 117 (rock climbing and) ; *U. cirrhosa*, 273–274 ; *U. cylindrica*, 168, 178, 186 ; *U. hirsuta*, 273–274, 285, 287 ; *U. hyperborea*, 273 ; *U. polyphylla*, 168, 178 ; *U. polyrrhiza*, 168 ; *U. proboscidea*, 186 ; *U. pustulata*, 168, 273–274, 277, 281 ; *U. spodochroa*, 273–274, 277, 281 ; *U. torrefacta*, 112, 168, 178, 187
- Usnea*, 177, 330 ; *U. dasypoga*, 273, 277, 284–287, 291 ; *U. filipendula*, 168 ; *U. florida*, 169 ; *U. florida* var. *hirta*, 13 ; *U. hirta*, 13, 169 ; *U. homescens*, 169 ; *U. intexta*, 169 ; *U. ochroleuca*, 304–305 ; *U. subfloridiana*, 169
- Variolaria exasperata*, 306
- Ventosa*, sect. of *Haematomma*, 248
- Verrucaria acrotella*, 51 ; *V. aethiobola*, 35–36, 38–44, 46, 48–49, 51–52, 169, 310 ; *V. aethiobola* f. *calcarea*, 51 ; *V. aethiobola* var. *cataleptoides*, 41, 169 ; *V. aethiobola* var. *griseocinerea*, 51 ; *V. alba*, 223 ; *V. alpicola*, 47, 51 ; *V. applanata*, 47 ; *V. aquatilis*, 36–40, 47, 50–51, 53 ; *V. biformis*, 228 ; *V. calcarea*, 49, 51 ; *V. calcaria*, 49, 51 ; *V. calciseda*, 169, 180 ; *V. cinereolutescens*, 46 ; *V. coerulea*, 169 ; *V. conformis*, 228 ; *V. conoidea*, 221–222 ; *V. conoidea* var. *subsquamacea*, 221 ; *V. conoidella*, 222 ; *V. conturmutula*, 51 ; *V. degenerascens*, 45 ; *V. devergescens*, 51 ; *V. dufourii*, 169, 181, 186 ; *V. elaeomelaena*, 36–41, 45–49, 52, 170 ; *V. epipolea*, 221 ; *V. filarskyana*, 47 ; *V. gemmata*, 223 ; *V. ger mata* var. *conoidea*, 221 ; *V. gemmata* var. *epipolaea*, 221 ; *V. gemmata* f. *epipolaea*, 221 ; *V. gemmata* var. *glaucia*, 223 ; *V. gemmata* subsp. *Salweii*, 225–226 ; *V. glaurina*, 170 ; *V. griseocinerascens*, 51 ; *V. hibernica*, 51–52, 77–82 ; *V. hochstetteri*, 13, 170 ; *V. hybernica*, 52 ; *V. hydrela*, 36–40, 42, 46–53 ; *V. imbrida*, 52 ; *V. irrigua*, 52 ; *V. kernstockii*, 36–41, 46–47, 50, 53 ; *V. kernstockii* f. *minutipuncta*, 46–47 ; *V. laevata*, 37, 42–43, 52 ; *V. latebrosa*, 36, 38–42, 44–45, 48, 51–52 ; *V. leightonii*, 52 ; *V. macrostoma*, 170 ; *V. margacea*, 36–41, 44, 47–49, 51–52, 126, 170 ; *V. maura*, 7, 13, 55–56 ; *V. maurula*, 50 ; *V. melaleuca*, 223 ; *V. microspora*, 13 ; *V. minutipuncta*, 46 ; *V. mucosa*, 39, 46, 53, 56 ; *V. muralis*, 13, 53, 170 ; *V. murina*, 170 ; *V. nigrescens*, 13, 36, 170, 181 ; *V. papillosa*, 52 ; *V. papillosa* var. *papillulosoides*, 53 ; *V. polycarpa*, 228 ; *V. praetermissa*, 36, 38–43 ; *V. pyrenuloides*, 77–82 ; *V. pyrenuloides* var. *hibernica*, 77–82 ; *V. reecta*, 53 ; *V. rheithrophila*, 37, 46, 53 ; *V. rhodosticta*, 52–53 ; *V. Salweii*, 225 ; *V. sphaerooides*, 223–224 ; *V. sphinctrina*, 170–171, 180–181, 186 ; *V. striatula*, 51 ; *V. striatula* var. *acrotella*, 51 ; *V. submersa*, 37, 53 ; *V. submersa* b. *litorea*, 35 ; *V. trabilis*, 35 ; *V. viridula*, 13, 53, 171 ; *V. vitricola*, 50 ; *Verrucaria* subgenus *Acrocordia*, 220
- Verrucariaceae*, 259

- Xanthoria aureola*, 14, 171, 181, 330, 332 ;
X. candelaria, 14, 115, 171, 175–176 ;
X. elegans, 171 ; *X. parietina*, 7, 14,
56, 58–59, 61–62, 64, 74–75, 171,
174–176, 187, chemistry: 271, 273,
277–279, 281–282, 284, cold resist-
ance: 290, nitrogen dioxide and, 339,
asci: 358 ; *X. parietina* var. *aureola*,
14 ; *X. parietina* var. *ectanea*, 14, 171 ;
X. polycarpa, 14, 115, 172, 175, 340
Xanthothallia, subgenus of *Acarospora*,
16–31
-

DATES OF PUBLICATION

- Vol. 4, Part 1: 27 June 1968.
Vol. 4, Part 2: 2 May 1969.
Vol. 4, Part 3: 16 April 1970.
Vol. 4, Part 4: 20 December 1970.

NOTICE TO AUTHORS

Papers and short notes from contributors anywhere in the world are welcome, and may be on any aspect of lichenology. They should be sent to the Editor. They must be typewritten in double or treble spacing, and a summary should generally be included. Illustrations should be in black ink on white paper, preferably Bristol board.

In taxonomic papers dealing with a series of species it is convenient either to present them alphabetically or to number them serially in their genera.

Localities in Britain should, unless there is good reason to the contrary, follow the spelling given on the latest editions of the maps published by the Ordnance Survey. Localities abroad should follow the spelling in *The Times Atlas of the World*.

The citation of references should be based on the *International Code of Botanical Nomenclature* 1966: 73-75.

Titles of periodicals should be abbreviated as in the latest edition of the *World List of Scientific Periodicals*. Titles of books should be given in full at the end of papers but may be judiciously abbreviated in lists of synonyms. Editions other than the first should be indicated.

EXAMPLES:

In text: Watson (1932), (Watson, 1932), or Watson (1932, p. 292).

In the list at end of paper: Watson, W. (1932) The bryophytes and lichens of moorland. *J. Ecol.* 20: 284-313.

Or if a book: Crombie, J. M. (1894) *A Monograph of Lichens Found in Britain*. London.

In list of synonyms: *Candelariella xanthostigma* (Ach.) Lett., *Hedwigia*, 52: 196 (1912).

Or if a book: *Pyrenotea lutea* Leight., *British Species of Angiocarpous Lichens*: 68, tab. 29, fig. 4 (1851).

Twenty-five reprints of original articles are supplied free. More may be purchased, as may reprints of other contributions such as book reviews. Reprints must be ordered when the author returns the final proofs of his article.

MEMBERSHIP

Membership of the British Lichen Society is open to all persons interested in lichenology. Subscriptions, due on joining and thence on 1 January each year, are: ordinary members, £2; junior members, £1; family members, 5s. Details of membership and application forms may be obtained from the Secretary, Mr J. R. Laundon, Department of Botany, British Museum (Natural History), Cromwell Road, London, S.W.7.

Referees

The function of referees, whose names appear inside the front cover, is to assist members in the identification of lichens. Specimens submitted should be adequate in size and well documented, including details of country, county or vice-county, borough or parish, precise locality and grid reference, altitude, date, habitat and ecology, notes, name of collector and collecting number. Identification should be attempted before sending the specimens to a referee and details of spores and chemical reactions should be included where appropriate. Return postage should be enclosed.

THE LICHENOLOGIST

Vol. 4 Part 4

1970

AHMADJIAN, V. and HEIKKILÄ, H.—The culture and synthesis of <i>Endocarpon pusillum</i> and <i>Staurothele clopima</i>	259–267
AHMADJIAN, V. and JACOBS, J. B.—The ultrastructure of lichens. III. <i>Endocarpon pusillum</i>	268–270
SOLBERG, Y. J.—Studies on the chemistry of lichens. VIII. An examination of the free sugars and ninhydrin-positive compounds of several Norwegian lichen species	271–282
SOLBERG, Y. J.—Studies on the chemistry of lichens. IX. Quantitative determination of monosaccharides and amino acids in hydrolysates of several Norwegian lichen species	283–288
KAPPEN, L. and LANGE, O. L.—The cold resistance of phycobionts from macrolichens of various habitats	289–293
TOPHAM, P. B. and SWINSCOW, T. D. V.— <i>Thelenidia monosporella</i> Nyl. in Greenland	294–296
LAUNDON, J. R.—Lichens new to the British flora: 4	297–308
JAMES, P. W.—The lichen flora of shaded acid rock crevices and overhangs in Britain	309–322
SYMPOSIUM	
DOBBS, G.—Introduction. The phycotropic fungi	323–325
LEWIS, D. H.—Physiological aspects of symbiotic associations between fungi and other plants	326–336
SAUNDERS, P. J. W.—Air pollution in relation to lichens and fungi	337–349
RICHARDSON, D. H. S.—Ascus and ascocarp structure in lichens	350–361
HAYNES, F. N. and MORGAN-HUWS, D. I.—The importance of field studies in determining the factors influencing the occurrence and growth of lichens	362–368
Book Review	369–370
Index	371–382

© British Lichen Society 1970
c/o Department of Botany, British Museum (Natural History),
Cromwell Road, London, S.W.7.

Published for the British Lichen Society by
Blackwell Scientific Publications Ltd, 5 Alfred Street, Oxford
Price £2 (\$6.50). Free to members of the British Lichen Society.

Annual subscription £3 (\$10.00), post free.
Published December 1970