

Original Paper

Cite this article: Impouma B *et al* (2021). The first 8 months of COVID-19 pandemic in three West African countries: leveraging lessons learned from responses to the 2014–2016 Ebola virus disease outbreak. *Epidemiology and Infection* **149**, e258, 1–8. <https://doi.org/10.1017/S0950268821002053>

Received: 6 March 2021

Revised: 31 July 2021

Accepted: 29 August 2021

Key words:




Ebola virus disease; COVID-19; SARS-CoV-2; pandemic; World Health Organization; response; infectious diseases; outbreak

Author for correspondence:

Benido Impouma,

E-mail: impoumab@who.int

The first 8 months of COVID-19 pandemic in three West African countries: leveraging lessons learned from responses to the 2014–2016 Ebola virus disease outbreak

Benido Impouma^{1,2} , George Sie Williams¹ , Fleury Moussana¹, Franck Mboussou¹, Bridget Farham¹ , Caitlin M. Wolfe^{1,3}, Charles Okot¹, Katrina Downing¹, Claudia Codeço Torres⁴, Antoine Flahault², Cyril Pervilhac¹, Georges Ki-Zerbo⁵, Peter Clement⁶, Steven Shongwe⁷, Olivia Keiser² and Ibrahima Socé Fall⁸

¹World Health Organization Regional Office for Africa, Brazzaville, Congo; ²Institute of Global Health, University of Geneva, Geneva, Switzerland; ³College of Public Health, University of South Florida, Tampa, FL, USA; ⁴Fundação Oswaldo Cruz, Rio de Janeiro, Brazil; ⁵World Health Organization Guinea Country Office, Conakry, Guinea; ⁶World Health Organization Liberia Country Office, Monrovia, Liberia; ⁷World Health Organization Sierra Leone Country Office, Freetown, Sierra Leone and ⁸World Health Organization Headquarters, Geneva, Switzerland

Abstract

Experience gained from responding to major outbreaks may have influenced the early coronavirus disease-2019 (COVID-19) pandemic response in several countries across Africa. We retrospectively assessed whether Guinea, Liberia and Sierra Leone, the three West African countries at the epicentre of the 2014–2016 Ebola virus disease outbreak, leveraged the lessons learned in responding to COVID-19 following the World Health Organization's (WHO) declaration of a public health emergency of international concern (PHEIC). We found relatively lower incidence rates across the three countries compared to many parts of the globe. Time to case reporting and laboratory confirmation also varied, with Guinea and Liberia reporting significant delays compared to Sierra Leone. Most of the selected readiness measures were instituted before confirmation of the first case and response measures were initiated rapidly after the outbreak confirmation. We conclude that the rapid readiness and response measures instituted by the three countries can be attributed to their lessons learned from the devastating Ebola outbreak, although persistent health systems weaknesses and the unique nature of COVID-19 continue to challenge control efforts.

Introduction

The coronavirus disease-2019 (COVID-19) pandemic continues to develop across Africa since the first case was reported on the continent in Egypt in February 2020 [1, 2]. As of 1 March 2021, over 3.9 million cases with 104 039 deaths have been reported from all 54 countries [3]. This represents a small fraction of the over 113 million cases and 2.5 million deaths reported globally [4]. However, despite the unique challenges presented by the COVID-19 pandemic, African countries are experienced in responding to outbreaks of emerging and re-emerging infectious diseases. The COVID-19 pandemic in many African countries comes in the context of multiple concurrent infectious disease outbreaks, for example Ebola virus disease (EVD), measles and Lassa fever [5] which require robust public health responses to mitigate the adverse impact of these events.

The major outbreak of EVD in West Africa from 2014 to 2016, during which 28 652 infections and 11 325 deaths were recorded, was unprecedented [6]. Three countries, Liberia, Guinea and Sierra Leone were at the epicentre of this outbreak. At the onset of this EVD outbreak, key pillars required for an effective response, such as coordination, surveillance, case management, infection prevention and control and community engagement, were inadequate or fractured [7, 8]. The inability to rapidly detect, isolate and treat cases to break the chain of transmission was a direct derivative of these weaknesses, resulting in the large-scale transmission of EVD and the associated high mortality rates.

However, with support from the international community, these countries overcame these challenges and the EVD outbreak was ended. At the same time, the countries gained experience and built improved mechanisms for responding to infectious disease outbreaks [9–11]. A key factor in improving capacity to detect and respond to outbreaks was the revitalisation of the Integrated Disease Surveillance and Response (IDSR) strategy [12–14]. The affected

© WHO Regional Office for Africa, Cite du Djoue Brazzaville Congo. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<http://creativecommons.org/licenses/by-nc-sa/4.0>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is used to distribute the re-used or adapted article and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use.

countries reported and successfully responded to several outbreaks of other infectious diseases in the aftermath of EVD [15–17]. In addition, joint assessments of the countries' post-EVD capacities to prevent, detect and respond to public health risks using a multisectoral approach have shown improvements compared to the pre-EVD period, but at the same time have highlighted that health system challenges remain [18–20].

Although EVD and COVID-19 differ in their mode of transmission and pathogenesis, many facets of preparedness and response for outbreaks of these two diseases overlap, and therefore key lessons learned from the response to EVD outbreak are applicable to the response to COVID-19 [21]. An effective response to COVID-19 also requires the capacity for robust surveillance, rapid case detection, disease confirmation, isolation of suspected and confirmed cases, treatment for moderate and severe cases, and a range of other public health measures to limit or prevent onward transmission.

In this study, we assessed whether the capacities and experience gained after the EVD outbreak in the three affected countries have had an impact on the response to COVID-19. In particular, we analysed the timeliness of case reporting and laboratory confirmation, and how readiness and public health response measures have affected the incidence of COVID-19 during the first 8 months of the pandemic. The findings from these analyses offer valuable lessons on the relevance of sustainably building capacities and strengthening health systems to meet the challenges posed by emerging disease outbreaks.

Methods

Study design and setting

We conducted a retrospective observational cross-sectional study to determine if the COVID-19 readiness and response measures implemented in Guinea, Liberia and Sierra Leone, spanning the period 1 February to 30 September 2020, were derived from the health system strengthening measures put in place as a result of lessons learned from the response to EVD outbreaks. We identified and assessed predictors for timely reporting and laboratory confirmation of cases. All three countries are located in West Africa and together have an estimated population of over 26 million [22].

Key readiness capacities and response measures

We selected seven key readiness capacities covering core thematic areas based on lessons learned from the response to the EVD outbreak as well as World Health Organization (WHO) recommendations to countries. An additional seven response measures implemented by the respective countries were also identified. [Table 1](#) outlines these capacities and measures, their relevance or rationale, and the definitions of their key milestone dates. The time to each milestone for readiness capacities was based on the duration (days) from WHO declaration of COVID-19 as a public health emergency of international concern (PHEIC) to the earliest date on which the readiness capacity was achieved or the measure instituted. For the response measures, the time to each milestone measured the duration in days from confirmation of the first COVID-19 case in the respective countries to the earliest date on which response measures were implemented.

Data sources and measurements

Key milestone dates were obtained from official reports and assessments shared by the respective countries with the WHO Regional Office for Africa (AFRO). In instances where the milestone date was missing, we extracted this information from the official government websites of the respective countries. All milestone dates were cross-checked with the official authorities of the respective countries. The population estimates for the respective countries were obtained from the World Bank database [22].

A line list of confirmed COVID-19 cases reported during the studied period was obtained from each country. We consolidated the line lists from all three countries and selected key variables for our analysis on the timeliness of case reporting and laboratory confirmation. The variables selected were country, date of report, age, sex, date of onset, date of confirmation, place of case detection and outcome (alive or dead). The date of report refers to the earliest date on which a person was notified to the health authorities as being suspected of having COVID-19. The date of onset refers to the earliest known date on which the case was reported to have begun experiencing signs and symptoms associated with COVID-19. The date of confirmation refers to the date on which the earliest laboratory results were released confirming the diagnosis of COVID-19 in each case. We defined the reporting timeliness as the time interval in days from onset of symptoms to case report, while the laboratory confirmation timeliness was defined as the duration from onset of symptoms to laboratory confirmation.

Only confirmed cases were included in our analysis on time to report and confirmation. A confirmed case of COVID-19 was defined as 'a person with a positive Nucleic Acid Amplification Test (NAAT) or a person with a positive SARS-CoV-2 Antigen-rapid diagnostic test (Ag-RDT) *and* meeting either the probable case definition or suspected criteria as per the WHO guideline, or an asymptomatic person with a positive SARS-CoV-2 Ag-RDT *and* who was a contact of a probable or confirmed case' [23]. We excluded asymptomatic cases, that is, confirmed cases who were reported as not experiencing any sign or symptoms despite being confirmed positive for COVID-19. This was intended to reduce bias because estimating timely reporting among asymptomatic cases may be more complex, without access to data on their likely dates of exposure to their source case and when that case became infectious.

Missing data

Three (0.1%) values were missing for time to report due to missing dates of report, while 133 (3.2%) were missing for time to confirmation due to missing date of confirmation. We performed multiple imputations for the missing data using the Multivariate Imputation by Chained Equations (MICE) package in R [24]. We performed five iterations using the country, age, sex, outcome and place of detection to impute the missing data.

Data analysis

We plotted a timeline of the key readiness and response milestone dates for the respective countries to show when these capacities or measures were first attained or implemented. Using 30 January 2020 as the date of the WHO declaration of PHEIC, we computed the duration in days to each milestone date for readiness. The number of days to each response measure milestone date was

Table 1. Selected COVID-19 readiness capacities and response measures, their rationale and definition of key milestone date on which capacities were attained on response measures first implemented

Category	Capacities/measures	Relevance/rationale	Definition of milestone date
Readiness	National strategic preparedness and response plan	A strategic preparedness and response plan (SPRP) outlines the public health measures that a country stands ready to implement to prepare for and respond to outbreaks. For a novel disease outbreak, an SPRP is crucial to guide policymakers and public health responders in critical decision-making processes.	Earliest date on which SPRP was finalised and adopted
	Surveillance capacity at health facility level	The ability of health workers to use case definitions to diagnose and report suspected cases of COVID-19 for early case detection.	Earliest date on which 90% of health facilities were covered with staff trained to detect and report suspected cases of COVID-19
	Diagnostic capacity at national level	The ability of the country to test and confirm a case of COVID-19 which is essential to trigger early response actions	Earliest date on which country gained capacity for conducting RT-PCR test for COVID-19
	Capacity for screening at points-of-entry	Useful for early detection of cases given the threat posed by importation of confirmed cases from other parts of the world. The international airports were considered as the place of greatest risk of importation during the early phase of the pandemic.	Earliest date on which screening of travellers began at the international airports
	Capacity for rapid response	A multi-disciplinary team is needed in the early phase of an epidemic for rapid investigation of alerts, quarantine or isolation of cases in order to find cases timely and implement quick public health measures to prevent mortality or interrupt transmission.	Earliest date on which trained national rapid response team was operationalised
	Capacity for case management	Prompt and optimal care necessary for preventing or reducing mortality.	Earliest date on which a functional treatment unit was made operational for the management of COVID-19 cases
	Capacity for risk communication and community engagements	Communicating the risk of the disease early, promoting preventive measures, engaging with the population through influencers to enhance preventive measures and respond early to rumours and misbeliefs	Earliest date on which the country initiated mass communication of COVID-19 messaging
Response	Activation of incidence management system	Critical to improve communication and information flow as well as to coordinate the public health response	Earliest date on which incidence management system was activated
	Suspension of commercial flights	Implemented to reduce the risk of international importation of cases while the countries study the situation and prepared better to manage international travellers	Earliest date on which international airport was closed
	Closure of schools	Implemented to reduce the risk of COVID-19 transmission among students	Earliest date on which all schools were closed
	Restriction on internal movements	Implemented to reduce the risk of spread of COVID-19 between different parts of the country	Earliest date on which restriction on internal movement started
	Restriction on mass gatherings	Implemented to reduce the risk of transmission of COVID-19 among people in mass gatherings such as concerts, sporting events, religious places, funerals etc.	Earliest date on which restrictions were implemented
	Mandatory wearing of face mask in public	Implemented to reduce the risk of human-to-human transmission of COVID-19	Earliest date on which wearing of face mask in public became mandatory
	Mandatory testing of travellers	Implemented after the resumption of commercial flights to prevent the importation of COVID-19 while also preventing international spread from these countries	Earliest date on which mandatory testing of travellers commenced

the duration from the first confirmed case in the respective countries. We computed the average number of days to each milestone across all three countries.

We also present key COVID-19 epidemiological parameters such as the number of cases and deaths, the case fatality ratio (CFR), which was defined as the proportion of deaths among all confirmed cases, the cumulative incidence per 100 000 population, and the mortality rates per one million population in the respective countries. We analysed the burden of infection

among health workers, which was calculated as the proportion of health worker cases among the total health workforce in the respective countries. Data on testing are presented as the cumulative number of tests performed per 10 000 population.

We computed the median number of days for case reporting and laboratory confirmation for symptomatic cases in the respective countries. The effect of selected covariates on the timeliness of reporting and confirmation were quantified by using two separate multivariable negative binomial regression models. The selected

covariates were country, age group, sex and place of detection. Our initial choice of these covariates was based on trends observed in the surveillance reports, a priori knowledge, literature available on this topic, and the extent to which data were available. Based on the age-associated mortality observed in the surveillance data, we categorised the cases into two age groups, <50 years and ≥ 50 years. Given the urban nature of the pandemic and the likelihood of access to health services to be skewed towards the capital cities in these countries, we also categorised the place of detection into two groups, those detected in the capital cities and those detected outside the capital cities. The results are presented as an incidence rate ratio (IRR) with a 95% confidence interval (CI). IRR greater than 1 indicates a longer time to detection or confirmation, while less than 1 indicates a shorter time.

Results

The timeline of key milestones for readiness capacities and response measures are presented in [Figure 1](#). The adoption and finalisation of a COVID-19 national strategic preparedness and response plan was the earliest readiness measure achieved across the three countries, taking an average of 5 days (range 3 days in Liberia to 7 days in Sierra Leone) ([Table 2](#)). The screening of all travellers at the respective international airports and the launch of a mass public communication campaign were the last readiness measures implemented, taking an average of 37 days each. It took an average of 29 days for the countries to obtain capacity for performing COVID-19 laboratory testing (range 25 days in Sierra Leone to 35 days in Liberia). [Table 2](#) further shows that the activation of a national incidence management system was the earliest public health response measure implemented, which predated the confirmation of the first case by 15 days on average (range 47 days before the first case in Sierra Leone to 2 days after the first case in Guinea). Results for the timeliness of various readiness and response measures in the respective countries are further shown in [Table 2](#).

A total of 14 227 cases with 220 deaths (CFR 1.5%) was reported from the three countries during the studied period. Guinea was the most affected with the highest case numbers ($n = 10\,652$) and highest cumulative incidence of 82 per 100 000 population ([Table 3](#)). Guinea also reported the highest number of health worker cases ($n = 513$), accounting for an estimated 5.0% of infection among the total health workforce in the country. Liberia reported the highest numbers of deaths ($n = 82$), CFR (6.1%), and mortality of 16 per million population. Other epidemiological results for the respective countries are shown in [Table 3](#).

Of 2406 symptomatic cases included in the study, the overall median reporting time was 3 days (interquartile range (IQR) (1–6)) while the laboratory confirmation time was 5 days (IQR (2–8)). When stratified by the respective countries, Liberia had the longest median reporting time of 5 days, while Sierra Leone had the shortest median reporting time of 1 day. The median laboratory confirmation time was also shortest in Sierra Leone ([Table 4](#)). Results of the negative binomial regression model showed that country (Liberia and Guinea) and age group (≥ 50 years) predictors were significantly ($P < 0.001$) associated with longer reporting and confirmation times compared to Sierra Leone and those below 50 years old, respectively. The place of detection (outside a capital city) was found to be significantly associated with shorter reporting time compared to those within a capital city, although with no influence on the laboratory

confirmation time. [Table 4](#) further shows that sex had no influence on the reporting or laboratory confirmation timeliness.

Discussion

Our results show that all selected readiness measures were instituted across the three countries within the first 2 months of declaration of the PHEIC by WHO. We also found very early response measures implemented across the three countries, with at least one response measure predating the confirmation of the first case. While the incidence of COVID-19 remained relatively low compared to most affected countries globally, our study showed a high CFR, particularly in Liberia and Sierra Leone. Other findings pointed to a high burden of infection among health workers and definite low testing rates compared to many countries globally. The overall median time of 3 days to report and 5 days for laboratory confirmation of symptomatic cases, although with variability among the countries, indicated that this delay could potentially be a key contributor to community transmission of the disease among the local population in the respective countries, despite the response measures instituted.

The relatively rapid readiness and response measures implemented could be attributed in part to the experience gained by these countries and the international community from the devastating EVD outbreak of 2014–2016 [25]. It took several months for these countries and the international community to recognise the potential threat and scale of the EVD outbreak, thereby delaying the institution of cogent control measures at national level, as well as slow mobilisation of international assistance [26]. Exposure to a previous major epidemic is associated with faster response, an indication that these countries have learned to move into action early [27]. Investments made in strengthening IDSR including laboratory capacities helped these countries to quickly adapt their structures in readiness and response to COVID-19. For example, of the four indicators for real-time surveillance assessed during the joint external evaluation on a scale of 1–5 (1 being no capacity and 5 being sustainable capacity), Liberia and Sierra Leone scored 4 (demonstrated capacity) and Guinea scored 3 (developed capacity) on three of the indicators [18–20]. National laboratory systems were also found to improve with specimen referral systems scoring 3 (developed capacity) across all three countries and testing capacity for priority diseases scoring 4 (demonstrated capacity) in Sierra Leone, 3 (developed capacity) in Guinea, and 2 (limited capacity) in Liberia although capacities in Liberia improved in the aftermath of the evaluation. These existing capacities were leveraged in response to the COVID-19 pandemic, however, the fact that all three countries scored 2 (limited capacity) for an interoperable, interconnected, electronic real-time reporting system as well as 2 (limited capacity) for an effective modern point-of-care and laboratory-based diagnostic indicated major weaknesses in these areas and the need for additional investments or support to effectively respond to a pandemic such as COVID-19. Additionally, the rapid support from WHO and other partner institutions to African countries including Guinea, Liberia and Sierra Leone helped bolster readiness capacities for COVID-19 in key areas, such as in the provision of laboratory reagents and other supplies, technical guidance, training etc [28, 29].

In spite of the rapidity with which measures were taken, it is clear that health systems in these countries remain fragile and underfunded [30], and these constraints can crucially affect health outcomes in a pandemic of this nature. The high CFR in Liberia




Fig. 1. Timeline of selected COVID-19 readiness and response measures in Sierra Leone, Guinea and Liberia, 1 February to 30 September 2020.

(6.1%) and Sierra-Leone (3.2%) vs. 0.6% in Guinea and the high proportion of health worker infections in Liberia (2%), Sierra Leone (3.5%) and Guinea (5%) in the early period of the pandemic could in part be attributed to the fragility of the health systems in these countries. In Liberia and Sierra Leone, the implementation of a policy to test all dead bodies for COVID-19 to improve detection, case investigation and contact tracing resulted in the identification of a high number of COVID-19 community deaths, indicating limited use of, access to and quality of health care services in these countries. The

relatively lower testing rates across these three countries with an average cumulative test of 71 tests per 10 000 population (45 tests per 10 000 population in Liberia, 52 tests per 10 000 population in Sierra Leone and 93 tests per 10 000 population in Guinea) compared to wealthy countries such as Denmark (6741 tests per 10 000 population), United States of America (3595 tests per 10 000 population) and the United Kingdom (3310 tests per 10 000 population) [31] could also be viewed in the context of resource constraints in low- and middle-income countries to offer mass-based testing to a larger per cent of their population.

Table 2. Duration to attainment or implementation of selected COVID-19 readiness capacities and response measures in Guinea, Liberia and Sierra Leone, 1 February–30 September 2020

Categories	Measures/capacities	Number of days to measures			
		Guinea	Liberia	Sierra Leone	Mean
Readiness measures reference date: 30 January 2020	National preparedness and response plan finalised and adopted	4	3	7	5
	90% of health facilities with staff trained in COVID-19 surveillance	N/A	21	27	–
	Testing capacity available (RT-PCR)	27	35	25	29
	Screening of travellers commenced at international airports	37	22	51	37
	National rapid response team activated	N/A	6	39	–
	Functional treatment unit ready for case management	14	25	53	31
	Mass public communication campaign commenced	31	31	50	37
Response measures reference dates Liberia: 16 March 2020; Guinea: 13 March 2020; Sierra Leone: 31 March 2020	National incidence management system activated	2	1	–47	–15
	Suspension of all commercial flights	8	32	–9	10
	School closure	14	5	0	6
	Restrictions on internal movements commenced	8	23	5	12
	Mandatory wearing of face mask commenced	36	42	22	33
	Mandatory testing of all travellers commenced	22	129	174	88
	Restrictions on mass gatherings commenced	14	25	–15	8

N/A means the date on which the milestone was first achieved was not available or could not be determined to compute the number of days to achievement after the declaration of PHEIC by WHO.

Table 3. Key epidemiological features of COVID-19 in Guinea, Liberia, and Sierra Leone, 1 February–30 September 2020

Category	Variable	Guinea	Liberia	Sierra Leone	Total
Population	Estimated population ^a	13 133 000	5 058 000	7 977 000	26 168 000
Cases	Cases	10 652	1344	2231	14 227
	Cumulative incidence per 100 000 pop	82	27	28	54
	Health worker cases	513	214	230	957
	Cases among total health workers (%)	5.0	2.0	3.5	3.5
Deaths	Deaths	66	82	72	220
	CFR (%)	0.6	6.1	3.2	1.5
	Deaths per million population	5	16	9	0.8
Tests	Number of RT-PCR tests ^b	121 487	22 499	41 128	185 114
	Test per 10 000 population	93	45	52	71
	Positivity rate (%) ^c	8.8	6.0	5.4	7.7

^aPopulation estimates for 2020 sourced from World Bank data.

^bNumber of tests based on number of persons tested.

^cPositivity rate is the proportion of all COVID-19 RT-PCR that tested positive for SARS-CoV-2 infection.

The reporting and confirmation timeliness were significantly longer in Guinea and Liberia compared to Sierra Leone. While asymptomatic COVID-19 cases may hold similar transmission

potential as symptomatic cases [32], the delay in detection (reporting and confirmation) of cases after the onset of symptoms particularly in the early phase of the pandemic in these countries

Table 4. Results of negative binomial regression model for reporting and confirmation timeliness of symptomatic COVID-19 cases in Guinea, Liberia and Sierra Leone, 1 February to 30 September 2020

Variables	Modalities	Symptomatic cases (n)	Time to report			Time to laboratory confirmation		
			Median (IQR)	³ IRR (95% CI)	P-value	Median (IQR)	IRR (95% CI)	P-value
	Total	2406	3 (1–6)			5 (2–8)		
Country	Sierra Leone	451	1 (0–4)	Reference		3 (2–6)	Reference	
	Liberia	309	5 (2–8)	1.81 (1.56–2.09)	<0.001	5 (3–8)	1.42 (1.28–1.58)	<0.001
	Guinea	1646	3 (1–5)	1.16 (1.03–1.30)	0.003	5 (3–8)	1.43 (1.31–1.55)	<0.001
Age group	<50 years	1720	2 (1–5)	Reference		5 (2–8)	Reference	
	≥50 years	686	3 (1–6)	1.16 (1.06–1.27)	<0.001	5 (3–9)	1.11 (1.05–1.19)	<0.0001
Sex	Females	838	2 (1–5)	Reference		5 (2–7)	Reference	
	Males	1568	3 (1–6)	1.09 (1.00–1.18)	0.05	5 (2–8)	1.05 (0.98–1.12)	0.23
Place of detection	Capital city	1771	3 (1–6)	Reference		5 (3–8)	Reference	
	Outside capital city	635	1 (0–4)	0.68 (0.62–0.76)	<0.001	4 (2–7)	0.96 (0.89–1.03)	0.53

³IRR, incidence rate ratio.

is an incontestable contributor to the transmission dynamics of the disease among the population. Kieran *et al.* found that the viral load of severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) is highest at or around symptom onset [33]. A systematic review and meta-analysis involving 79 studies also showed that people are likely to be highly infectious with the SARS-CoV-2 virus in the first week after the onset of symptoms [34]. These findings mean that delay in the initiation of intervention measures (such as isolation and contact tracing) as a result of the delay in reporting and confirming cases would provide ample time for symptomatic cases to continue transmitting the infection.

Delays in reporting and confirmation among cases ≥50 years may have led to delay in initiation of treatment, one of several factors which could have resulted in the high CFR in this age group. This finding is consistent with several studies that have shown that people in older age group are at higher risk of complications and deaths from COVID-19, especially when life-saving interventions are delayed [35]. Also, we found it interesting to note that COVID-19 cases were reported earlier among people outside the capital cities compared to those living within the capital cities. Disease surveillance systems are likely to be quickly overwhelmed in high-density population areas as the rate of transmission increases [36]. This could be a result of high population densities in the capital cities of the various countries making surveillance of COVID-19 more challenging. Intense transmission of EVD in the capital cities of the three countries complicated response efforts in the 2014–2016 outbreak [37].

There are a few limitations to our study. First, although we showed how early these readiness and response measures were implemented, we did not assess the efficiency of their implementation. The early implementation of the restriction measures may have resulted in fatigue among the population, who probably also suffered adverse socio-economic effects. Hence, strict adherence to measures such as wearing a face mask would likely wane over time, potentially resulting in resurgence of cases. Second,

most COVID-19 cases are asymptomatic; therefore, the analysis of symptomatic cases may not provide a full picture of the speed with which cases are reported and confirmed in the countries. However, given that mass population-based testing had not been implemented and that most asymptomatic cases were only identified after testing contacts of symptomatic cases, analysis of how rapidly symptomatic cases were identified can provide an estimation of the timeliness of case detection. Third, the statistical power, though not effect, of our analysis, may have been lowered due to omissions or recording errors in onset dates in the line list, leading to the exclusion of potential symptomatic cases. Lastly, we did not account for the effect of contact tracing on reporting and confirmation timeliness because such data were not available. High levels of contacts tracing are more likely to lead to early reporting and confirmation due to the regular monitoring [38].

Despite these limitations, our study has shown that these countries took actions early in the form of readiness and response to avert the negative consequences that they had experienced during the Ebola outbreak of 2014–2016. Using lessons learned from the EVD outbreak as well as capacities gained in its aftermath, the countries were able to take key readiness and response measures early which may have contributed to the low incidence of COVID-19 in these countries, despite the unique challenges posed by COVID-19 given the role of asymptomatic transmission in the context of low level of COVID-19 testing. Strong technical and operational support from WHO and partners has also helped these countries to continue to respond to the pandemic. However, eventual control of the pandemic will require continued implementation of public health measures, along with vaccination campaigns. It would be interesting for future studies to consider assessing the effectiveness of the various response measures implemented and their impact on the COVID-19 pandemic in these countries.

Acknowledgements. We wish to thank all those who contributed to this study. We are particularly grateful to Kwaukuan Yealue of WHO Liberia

Country Office, Robert Musoke of WHO Sierra Leone Country Office, and Mamadou Balde of WHO Guinea Country Office for validating the data on COVID-19 response and readiness milestone dates for the respective countries.

Data availability statements. The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

1. Medhat MA and Kassas ME (2020) COVID-19 in Egypt: uncovered figures or a different situation? *Journal of Global Health* **10**, 010368.
2. Rice BL *et al.* (2021) Variation in SARS-CoV-2 outbreaks across sub-Saharan Africa. *Nature Medicine* **27**, 447–453. doi: <https://doi.org/10.1038/s41591-021-01234-8>.
3. Africa CDC. (2020) Coronavirus Disease 2019 (COVID-19) – Africa CDC. Africa CDC Dashboard. <https://africacdc.org/covid-19/> (accessed 5 March 2021).
4. World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard | <https://covid19.who.int> (accessed 5 March 2021).
5. WHO Regional Office for Africa. (2020) Weekly Bulletin on Outbreaks and Emergencies. (August):1–20. <https://apps.who.int/iris/bitstream/handle/10665/331023/OEW07-1016022020.pdf> (accessed 5 March 2021).
6. Centers for Disease Control and Prevention. 2014–2016 Ebola Outbreak in West Africa | <https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/index.html> (accessed 5 March 2021).
7. Coltart CEM *et al.* (2017) The Ebola outbreak, 2013–2016: old lessons for new epidemics. *Philosophical Transactions Royal Society B Biological Science* **372**, 2013–2016.
8. McNamara LA *et al.* (2016) Ebola surveillance – Guinea, Liberia, and Sierra Leone. *MMWR Supplement* **65**(Suppl-3), 35–43. DOI: <http://dx.doi.org/10.15585/mmwr.su6503a6external-1>.
9. Vetter P *et al.* (2016) The 2014–2015 Ebola outbreak in West Africa: hands On. *Antimicrobial Resistance and Infection Control* **5**, 1–17. <http://dx.doi.org/10.1186/s13756-016-0112-9>.
10. Mobula LM *et al.* (2018) A humanitarian response to the West African Ebola virus disease outbreak. *Journal of International Humanitarian Action* **3**, 10. doi: <https://doi.org/10.1186/s41018-018-0039-2>.
11. Marston BJ *et al.* (2017) Ebola response impact on public health programs. *Emerging Infectious Diseases* **23**, S25–S32.
12. Nagbe T *et al.* (2019) The implementation of integrated disease surveillance and response in Liberia after Ebola virus disease outbreak 2015–2017. *Pan African Medical Journal* **33**, 3.
13. Njuguna C *et al.* (2019) Revitalization of integrated disease surveillance and response in Sierra Leone post Ebola virus disease outbreak. *BMC Public Health* **19**, 364.
14. Hemingway-Foday J *et al.* (2019) Improving integrated disease surveillance and response capacity in Guinea, 2015–2018. *Online Journal of Public Health Informatics* **11**, e364.
15. Rude JM *et al.* (2019) Rapid response to meningococcal disease cluster in Foya district, Lofa County, Liberia January to February 2018. *Pan African Medical Journal* **33**, 6.
16. Suk JE *et al.* (2016) Post-Ebola measles outbreak in Lola, Guinea, January–June 2015. *Emerging Infectious Diseases* **22**, 1106–1108.
17. World Health Organization. Measles outbreak confirmed in northern Sierra Leone _ WHO _ Regional Office for Africa. <https://www.afro.who.int/news/measles-outbreak-confirmed-northern-sierra-leone> (accessed 5 March 2021).
18. World Health Organization. Évaluation externe conjointe des principales capacités RSI de la République de Guinée. 2017; 23–8. <https://apps.who.int/iris/handle/10665/258726> (Accessed 5 March 2021).
19. World Health Organization. Joint External Evaluation of IHR Core Capacities of the Republic of Liberia. Mission report. September 2016 <https://apps.who.int/iris/bitstream/handle/10665/255268/WHO-WHE-CPI-2017.23-eng.pdf?sequence=1&isAllowed=y> (Accessed 5 March 2021).
20. World Health Organization. Joint External Evaluation of IHR Core Capacities of the Republic of Sierra Leone. November 2017. <https://apps.who.int/iris/bitstream/handle/10665/254790/WHO-WHE-CPI-2017.16-eng.pdf?sequence=1> (Accessed 5 March 2021).
21. Mobula LM *et al.* (2020) Recommendations for the COVID-19 response at the national level based on lessons learned from the Ebola virus disease outbreak in the Democratic Republic of the Congo. *American Journal of Tropical Medicine and Hygiene* **103**, 12–17.
22. The World Bank. (2020) Population estimates and projections | DataBank. The World Bank Data Bank. <https://databank.worldbank.org/source/population-estimates-and-projections#> (accessed 5 March 2021).
23. World Health Organization. Public Health Surveillance for COVID-19, WHO Interim Guidance. 1 August 2020. <https://apps.who.int/iris/handle/10665/333752> (accessed 5 March 2021).
24. van Buuren S and Groothuis-Oudshoorn K (2011) Mice: multivariate imputation by chained equations in R. *Journal of Statistical Software* **45**, 1–67.
25. Maxmen A (2020) Ebola prepared these countries for coronavirus – but now even they are floundering. *Nature* **583**, 667–668.
26. Moon S *et al.* (2015) Will Ebola change the game? Ten essential reforms before the next pandemic. the report of the Harvard-LSHTM independent panel on the global response to Ebola. *Lancet* **386**, 2204–2221.
27. Tsuei SH-T (2020) How previous epidemics enable timelier COVID-19 responses: an empirical study using organisational memory theory. *BMJ Global Health* **5**, e003228.
28. World Health Organization. COVID-19 in Africa: marking six months of response | WHO | Regional Office for Africa. <https://www.afro.who.int/covid-19-africa-marking-six-months-response> (Accessed 5 March 2021).
29. World Bank. (2020) World Bank's Response to COVID-19 (Coronavirus) in Africa. Factsheet. <https://www.worldbank.org/en/news/factsheet/2020/06/02/world-banks-response-to-covid-19-coronavirus-in-africa> (accessed 5 March 2021).
30. World Bank. (2020) Health Nutrition and Population Statistics | DataBank. World Bank. <https://databank.worldbank.org/source/health-nutrition-and-population-statistics#> (accessed 5 March 2021).
31. Ra SH *et al.* (2020) Upper respiratory viral load in asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection. *Thorax* **76**, 61–63.
32. Our world in data. <https://ourworldindata.org/grapher/full-list-cumulative-total-tests-per-thousand-map?time=2020-09-30>.
33. Walsh KA *et al.* (2020) SARS-CoV-2 detection, viral load and infectivity over the course of an infection. *Journal of Infection* **81**, 357–371.
34. Cevik M *et al.* (2020) SARS-CoV-2, SARS-CoV-1 and MERS-CoV viral load dynamics, duration of viral shedding and infectiousness: a living systematic review and meta-analysis. *SSRN Electronic Journal* **5247**, 1–10.
35. Starke KR *et al.* (2020) The age-related risk of severe outcomes due to covid-19 infection: a rapid review, meta-analysis, and meta-regression. *International Journal of Environmental Research and Public Health* **17**, 1–24.
36. Lee VJ *et al.* (2020) Epidemic preparedness in urban settings: new challenges and opportunities. *Lancet Infectious Diseases* **20**, 527–529.
37. Campbell L, Adan C and Morgado M. Ebola Response in cities: responding in the context of quarantine. <https://reliefweb.int/sites/reliefweb.int/files/resources/alnap-urban-2017-ebola-response-in-cities-learning-for-future-public-health-crises.pdf> (Accessed 5 March 2021).
38. Keeling MJ, Hollingsworth TD and Read JM (2020) Efficacy of contact tracing for the containment of the 2019 novel coronavirus (COVID-19). *Journal of Epidemiology and Community Health* **74**, 861–866.