

assess the success of these interventions in improving best medical practice.

Results. Of the 23 service users, 20 patients had their VTE risk assessment completed on admission and there was a delay of over a month with the remaining 3 patients. Of significance is that of all initial VTEs, 6 out of 23 contained inaccurate details, such as omission of comorbidities or a subjective assessment of mobility, indicating the need to use a standardized tool which allows for comparison across time. The mean admission duration for all 23 inpatients, as of February 2022 was calculated to be 16.2 months, with a range of 2 and 59 months. 15 patients did not have their VTE risk assessment repeated during admission, and of these 2 did have a change in their risk profile, indicating non-adherence with NICE guidelines.

Conclusion. This study has identified significant areas for improvement, specifically the need for clear timing for repeated VTE assessments, consistent sources of patient's medical history and documentation of mobility status. The project has highlighted the need for a more robust VTE assessment protocol which is currently being developed, to improve patient mortality and outcomes.

Pre-Menstrual Dysphoric Disorder (PMDD) in Young People: What We Know About It, the Role of CBT as a Treatment Option, and the Development of High-Quality Psychoeducation Material for Clinical Use

Dr Judy King* and Miss Hannah Wilkinson

Berkshire Healthcare NHS Foundation Trust, Berkshire, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.314

Aims. We aim to explain more about PMDD in young people and explore the evidence looking at the role of CBT as an intervention. We aim to develop high-quality psychoeducation material with the involvement of young people, to be able to offer clinically relevant information and empower young people with PMDD.

Methods. A literature search was conducted in 2019 and updated in February 2022 using the Cochrane Library, Psych-info, MEDLINE, Cinahl, EMBASE and Google Scholar, to look at the evidence available for CBT in young people with PMDD or PMS. The search included PMS as well as PMDD due to the heterogeneity in definitions used in studies.

Focus groups with young people are underway to develop high quality written psychoeducation material about PMDD.

Results. There were no specific studies looking at CBT as an intervention in young women under the age of 18 with PMS or PMDD. There was one intervention study with a treatment arm of psychoeducation in 62 young people under 18 with PMS versus a no treatment group (Taghizadeh 2013), with improvement in symptoms after 3 months from baseline reported.

There were more studies available in women over the age of 18. The search identified 3 meta-analyses in 2009 and 2012 (Busse 2009, Lusty, 2009; Kleinstauber 2012) and a more recent systematic review (Landolt 2020). Kleinstauber et al included 22 RCTs, with a median age of 39, and broadly showed CBT to have a small to medium positive effect size, although any conclusions were limited due to the small numbers involved in the individual trials and methodological flaws. Landolt 2020 looked at CBT or elements of CBT as an intervention arm over the last 30 years or so in woman of all ages. Variations of CBT including virtual, group, couple and psychoeducation alone were included, and all were reported to have favourable outcomes.

Conclusion. There is a huge gap in research looking at PMDD in the adolescent population. Translating research from adults is not ideal. Increasing awareness and developing psychoeducation is a step in the right direction.

An Evaluation of Admissions to an Old Age Psychiatry Ward: A Quality Improvement Project

Dr Michael Cooper, Dr Clare Langan*, Dr Christopher Haxton, Dr Philip Andrew and Dr Mohd Yazid

NHS Greater Glasgow and Clyde, Glasgow, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.315

Aims. It has been well publicised the pressures on inpatient bed capacity within mental health services in recent years. The RCPsych have stated in their publication 'Exploring mental health inpatient capacity' that bed occupancy has risen above their recommended 85% occupancy target in most areas. Waiting lists for beds have also grown. The aim of this project was to identify whether there could have been any extra community resources in place that could have prevented admissions to hospital within an older adult CMHT. This in turn reducing the demand on inpatient beds.

Methods. Inclusion criteria: All patients from the older adult CMHT admitted to either an organic or functional mental health inpatient bed between the 1st September and 31st December 2021.

All patients who met the inclusion criteria were discussed as part of a panel consisting of members of the MDT who were involved in the patients' ongoing care. The panel discussed each patient and individually scored each admission on a scale of 1-5 (where 1 was deemed to be very avoidable and 5 completely unavoidable). Where an admission did not score a 5 we considered whether anything could have been in place to have prevented the admission.

Results. Our search identified 21 patients who had been admitted to the respective old age psychiatry ward during our period of interest. The predominant diagnosis of these patients was vascular dementia (n=5), followed by Alzheimer's disease (n=3). Following our consensus panel discussion, we identified that the most common reason for admission to hospital was for management of behavioural and psychological symptoms of dementia (n=10), followed by increasing patient vulnerability (n=4) in the community. Carer stress was a theme in 2 admissions. Following panel discussion regarding potential avoidability of admission, we identified that 14 out of the total of 21 admissions scored a 5, 1 scored 4, 1 scored 3, and 5 scored 2.

Conclusion. Behavioural and psychological symptoms of dementia continues to remain a significant clinical challenge and was the most common reason for admission in our patient cohort. The majority of admissions to hospital in our cohort were deemed unavoidable as a result. However, we identified that carer stress was a significant theme in 2 out of our 21 admissions, suggesting potential scope to implement services which may reduce carer stress and ultimately, prevent hospital admission.

The HDAT Helper - Developing an Online Tool to Improve the Safety and Accuracy of Antipsychotic Prescribing

Dr Christopher Lawrence*

Southern Health NHS Foundation Trust, Southampton, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2022.316