

# ALMA spectrum of the extreme OH/IR star OH 26.5+0.6

K. Justtanont<sup>1</sup>, S. Muller<sup>1</sup>, M. J. Barlow<sup>2</sup>, D. Engels<sup>3</sup>,  
D. A. García-Hernández<sup>4,5</sup>, M. A. T. Groenewegen<sup>6</sup>, M. Matsuura<sup>7</sup>,  
H. Olofsson<sup>1</sup>, D. Teyssier<sup>8</sup>, I. Marti-Vidal<sup>1</sup>, T. Khouri<sup>1</sup>,  
M. Van de Sande<sup>9</sup>, W. Homan<sup>9</sup>, T. Danilovich<sup>9</sup>, A. de Koter<sup>10</sup>,  
L. Decin<sup>9</sup>, L. B. F. M. Waters<sup>11,10</sup>, R. Stancliffe<sup>12</sup>, W. Vlemmings<sup>1</sup>,  
P. Royer<sup>9</sup>, F. Kerschbaum<sup>13</sup>, C. Paladini<sup>14</sup>, J. Blommaert<sup>9</sup>  
and R. de Nutte<sup>9</sup>

<sup>1</sup>Chalmers University of Technology, Onsala Space Observatory, S-439 92 Onsala, Sweden

<sup>2</sup>Univ. College London, Dept. of Physics & Astronomy, Gower Street, London, UK

<sup>3</sup>Hamburger Sternwarte, Gojenbergsweg 112, D-21029 Hamburg, Germany

<sup>4</sup>Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

<sup>5</sup>Departamento de Astrofísica, Univ. de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain

<sup>6</sup>Koninklijke Sterrenwacht van België, Ringlaan 3, 1180 Brussels, Belgium

<sup>7</sup>School of Physics & Astronomy, Cardiff University, The Parade, Cardiff, UK

<sup>8</sup>Telespazio Vega UK Ltd for ESA/ESAC, Camino bajo del Castillo, s/n, Urbanización Villafranca del Castillo, Villanueva de la Cañada, E-28692 Madrid, Spain

<sup>9</sup>Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium

<sup>10</sup>Sterrenkundig Instituut “Anton Pannekoek”, Science Park 904, 1098 XH Amsterdam, The Netherlands

<sup>11</sup>SRON Netherlands Institute for Space Research, PO Box 800, 9700 AV Groningen, The Netherlands

<sup>12</sup>Argelander-Inst. fr Astronomie, Univ. of Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

<sup>13</sup>University of Vienna, Dept. of Astrophysics, Türkenschanzstrasse 17, 1180 Wien, Austria

<sup>14</sup>European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile


**Abstract.** We present ALMA band 7 data of the extreme OH/IR star, OH 26.5+0.6. In addition to lines of CO and its isotopologues, the circumstellar envelope also exhibits a number of emission lines due to metal-containing molecules, e.g., NaCl and KCl. A lack of C<sup>18</sup>O is expected, but a non-detection of C<sup>17</sup>O is puzzling given the strengths of H<sub>2</sub><sup>17</sup>O in Herschel spectra of the star. However, a line associated with Si<sup>17</sup>O is detected. We also report a tentative detection of a gas-phase emission line of MgS. The ALMA spectrum of this object reveals intriguing features which may be used to investigate chemical processes and dust formation during a high mass-loss phase.

**Keywords.** stars: AGB and post-AGB, circumstellar matter, stars: individual (OH 26.5+0.6), stars: late-type, stars: abundances

---

## 1. Introduction

A number of intermediate-mass ( $\sim 4\text{--}8M_{\odot}$ ) that evolve on the AGB are known to be undergoing hot-bottom burning (HBB) from observations of enhancement of <sup>7</sup>Li and other s-process elements (e.g., Garcia *et al.* 2013). The CNO cycle operates during this



**Figure 1.** ALMA spectrum of CO  $J=3-2$  and its isotopologues (bottom),  $\text{Si}^{17}\text{O}$  and a tentative detection of MgS. The vertical line denotes the LSR velocity of OH 26.5+0.6.

evolutionary phase and drives the  $^{12}\text{C}/^{13}\text{C}$  towards the equilibrium value of  $\sim 4$ . The process shuts down when the envelope mass is reduced to  $1 M_{\odot}$  (Karakas & Lattanzio 2014). The Herschel spectrum OH 26.5+0.6 shows a lack of  $\text{H}_2^{18}\text{O}$  while  $\text{H}_2^{16}\text{O}$  and  $\text{H}_2^{17}\text{O}$  are readily detected (Justanont *et al.* 2013). HBB preferentially destroys  $^{18}\text{O}$  (Karakas & Lattanzio 2014) thereby confirming that the progenitor of OH 26.5+0.6 is an intermediate-mass star. We subsequently observed the object with ALMA in band 7 in 2016 with spectral windows centered on the transition  $J=3-2$  of CO,  $^{13}\text{CO}$ ,  $\text{C}^{17}\text{O}$  and  $\text{C}^{18}\text{O}$  (Justanont *et al.* 2018, ADS/JAO.ALMA#2015.1.00054.S).

## 2. The ALMA spectrum

A total of about 60 emission lines have been detected in our ALMA observations. Fig. 1 shows the spectrum of CO isotopologues. The  $\text{C}^{17}\text{O}$   $J=3-2$  is not detected above the noise which is unexpected considering that strong  $\text{H}_2^{17}\text{O}$  lines have been detected in the Herschel spectrum of the star. However, we detected a line which can be attributed to  $\text{Si}^{17}\text{O}$   $J=8-7$  at 334.3015 GHz. The resolution of this line is  $14 \text{ km s}^{-1}$  as it falls in a spectral window assigned to a continuum measurement. The ALMA spectrum indicates a possible chemical pathway of molecular formation of oxygen in a high density environment:  $^{17}\text{O}$  is locked up in  $\text{H}_2^{17}\text{O}$  and  $\text{Si}^{17}\text{O}$  rather than  $\text{C}^{17}\text{O}$ .

A line at 335.9845 GHz may be assigned to a new circumstellar molecule. It corresponds to the MgS  $J=21-20$  transition. Previously, a broad dust emission feature at  $30 \mu\text{m}$  has been attributed to MgS dust, but this has been observed only towards C-rich circumstellar environments. A number of lines in the spectrum are due to the lines of SO and metal-containing molecules like NaCl, KCl and their isotopologues. Unlike the low-mass AGB stars, no  $\text{SO}_2$  lines are detected within the spectral range covered by our observations.

## References

- García-Hernández, D. A., Zamora, O., Yagüe, A., *et al.* 2013, *A&A*, 555, L3  
 Justanont, K., Muller, S., Barlow, M. J., *et al.* 2018, *A&A*, submitted  
 Justanont, K., Teyssier, D., Barlow, M. J., *et al.* 2013, *A&A*, 556, A101  
 Karakas, A. I. & Lattanzio, J. C. 2014, *PASA*, 31, 30