

and non-debt group (37% and 44% respectively,  $\chi^2=0.03$ , d.f.=1, NS). The comparable proportions for the control group were 43% and 2% (Fisher exact,  $P=0.006$ ).

This absence of a difference in depression between those in and not in debt in the DSH group is at variance with the Hatcher's report. Methodological differences, a difference in the proportion of males, the inclusion of those other than self-poisoners and bias introduced by nursing staff who did not give questionnaires to all DSH patients may explain this. The similarities in ages and the proportions in debt are however striking.

This study found that a brief questionnaire was acceptable for both DSH and this control group. The proportion of questions answered about debt was high and was representative of the other questions. This method may enable large control groups to be studied. In addition this study supports the hypothesis that debt is more common in the DSH population. To draw further conclusions from these numbers is hazardous although the comparative levels of depression associated with debt in both DSH and control populations warrants further investigation.

HAMER, D., SANJEEV, D., BUTTERWORTH, E., *et al* (1991) Using the Hospital Anxiety and Depression Scale to screen for psychiatric disorders in people presenting with deliberate self-harm. *British Journal of Psychiatry*, **158**, 782-784.

SIMON J. TAYLOR

*Chesterfield Royal Hospital  
Chesterfield S44 5BL*

#### Risk of HIV for women who inject drugs

SIR: Gossop *et al* (*BJP*, January 1994, **164**, 101-104) suggest that women may be at higher risk of relapse to drug use than men because of their closer social attachment to other drug users. Preliminary results from our survey of opiate users presenting for treatment in south London are broadly in line with those of Gossop *et al*. However, we found that women who inject drugs may also be at greater risk of HIV infection than men.

Women comprised 24% of a sample of 97 opiate-dependent drug users questioned about sexual and injecting behaviours. They were more likely to be living with a drug user (61% compared with 31% of men;  $\chi^2=6.6$ ,  $P<0.05$ ) and to have a sexual partner who was an injecting drug user (70% compared with 22% of men;  $\chi^2=18.2$ ,  $P<0.0001$ ). There were no significant differences between the proportions of men and women who had recently injected drugs, but women were more likely to have injected

themselves with equipment that had been used by someone else (30% of women compared with 11% of men;  $\chi^2=5.2$ ,  $P<0.05$ ). For both sexes, the majority of equipment sharing episodes involved their regular sexual partner, with whom condoms were used by only 20% of male and 25% of female clients.

In drug-using couples, women may have less opportunity to negotiate safer sexual and injecting practices, because men play a more active role in procuring drugs and injecting equipment (Klee, 1993). This particular aspect of relationships between male and female drug users has implications for HIV prevention strategies.

KLEE, H. (1993) HIV risks for women drug injectors: heroin and amphetamine users compared. *Addiction*, **88**, 1055-1062.

SALLY PORTER  
JAMES ELANDER  
SUE STEPHENS  
ANDREW JOHNS  
JAMES EDEH

*Division of Psychiatry of Addictive Behaviour  
St George's Hospital Medical School  
London SW17 0RE*

#### Dysphagia in the neuroleptic malignant syndrome

SIR: The neuroleptic malignant syndrome (NMS) is a recognised complication of psychotropic drug use with a variable outcome (Bristow, 1993). We describe two cases where dysphagia with loss of a gag reflex was a major presenting feature.

*Case 1:* A 25-year-old male presented acutely psychotic and received zuclopenthixol, chlorpromazine and droperidol. He developed NMS with dysphagia, loss of a gag reflex followed by right lower zone pneumonia which required ventilation for 7 days, during which time he received dantrolene, bromocriptine and lorazepam (Gratz *et al*, 1992). His fever finally settled on imipenem. No organisms were grown. He was noted to have absent bowel sounds which returned after 9 days following the administration of cisapride. However his swallowing was unaffected by this and returned on day 25. An MRI scan of the brain during the dysphagia showed no evidence of brainstem damage.

*Case 2:* A 32-year-old male presented with dysphagia and fever and was noted to have a poor gag reflex. He had been receiving chlorpromazine and flupenthixol for 7 weeks. A diagnosis of NMS was made and he received dantrolene, bromocriptine and lorazepam. He developed signs of right lower zone pneumonia which responded to ciprofloxacin and erythromycin. By day 11 he could speak but still was unable to swallow; MRI scanning showed no brainstem abnormalities. He required a tracheostomy and feeding gastrostomy. He had a fatal cardiorespiratory arrest on day 16 due to bilateral pulmonary emboli.