

# The Fast Evolution of SN 2010bh associated with GRB 100316D

Felipe Olivares E.<sup>1</sup>, Jochen Greiner<sup>1</sup>, Patricia Schady<sup>1</sup>, Arne Rau<sup>1</sup>,  
Sylvio Klose<sup>2</sup>, and Thomas Krühler<sup>3</sup> for the GROND team

<sup>1</sup>Max-Planck-Institute für extraterrestrische Physik,

Giessenbachstraße 1, 85748, Garching, Germany, email: foe@mpe.mpg.de

<sup>2</sup>Thüringer Landessternwarte Tautenburg, Sternwarte 5, 07778, Tautenburg, Germany

<sup>3</sup>Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen,  
Juliane Maries Vej 30, 2100, Copenhagen, Denmark

**Abstract.** We report on the type-Ic SN 2010bh associated with XRF 100316D at  $z = 0.059$ , which is among the latest spectroscopically confirmed GRB-SNe (Bufano *et al.* 2012). This supernova proves to be the most rapidly evolving GRB-SN to date.

**Keywords.** supernovae: individual (SN 2010bh), gamma-rays: burts

## 1. Introduction

So far only a handful of supernovae (SNe) associated with gamma-ray bursts (GRBs) have been spectroscopically confirmed; see Woosley & Bloom (2006) and Hjorth & Bloom (2011) for reviews. Out of those only SN 2006aj associated with XRF 060218 has shown signatures of the cooling of the shock breakout (Campana *et al.* 2006).


For the case of the association SN 2010bh/XRF 100316D, GROND provided data from 0.5 to 80 days after the burst covering a wavelength range from 350 to 1800 nm, significantly expanding the pre-existing data set for this event (e.g., Cano *et al.* 2011).

## 2. Results

**Broad Band Spectral Energy Distribution:** Detections at 50 ks in  $g'r'i'z'J$  are combined with a UVOT  $uvw1$ -band detection at 33 ks and an interpolated XRT spectrum. We model the SED (left panel of Fig. 1) with a blackbody associated with the cooling of the shock breakout and a power-law representing the afterglow. We obtain a host-galaxy extinction of  $A_{V,\text{host}} = 1.2 \pm 0.1$  mag and a metal absorption equivalent to a hydrogen column density of  $N_{H,\text{host}} = (4.4 \pm 0.4) \times 10^{22} \text{ cm}^{-2}$ .

**Evolution of the Thermal Component:** Early X-ray measurements from Starling *et al.* (2011) complement our results for the temperature and radius of the blackbody component (Fig. 1, right panel). Assuming a linear growth, we obtain  $v = 8000$  km/s. The best model is a power law, which yields an initial apparent radius of  $R_0 = (7.0 \pm 0.9) \times 10^{11}$  cm. As seen in the upper right panel of Fig. 1, adiabatic expansion fails to reproduce  $T_{BB}(t)$ .

**Pseudo Bolometric Light Curve:** The flux integrated from the  $g'$  to the  $H$  bands was fitted using the two component model from Maeda *et al.* (2003), which consists of a dense inner core and an outer core with lower opacity (e.g., Valenti *et al.* 2008). The transition from optically thick to thin occurs at  $\approx 33$  days after the burst for SN 2010bh. The physical parameters of the explosion are  $M_{\text{Ni}} = (0.21 \pm 0.03)M_{\odot}$ ,  $M_{\text{ej}} = (2.60 \pm 0.23)M_{\odot}$ , and  $E_k = (2.4 \pm 0.7) \times 10^{52}$  erg.



**Figure 1.** *Left:* Broad-band spectral energy distribution. The afterglow spectral slope is  $\Gamma = 1.8 \pm 0.1$ . *Right:* Temporal evolution of the blackbody component. Temperature is shown in the upper panel and blackbody radius in the lower panel.

### 3. Conclusions

Combining GROND and *Swift* data, the early broad-band SED is modeled with a blackbody and power-law component attenuated by ISM absorption in the host galaxy. The evolution of the thermal component reveals a cooling envelope at an apparent initial radius, which is compatible with a dense wind surrounding a WR star. Moreover, the early-time expansion velocities are compatible with the SN nature. Multicolor templates of SN 1998bw show that SN 2010bh is on average 70% as bright as SN 1998bw (see Olivares E. *et al.* 2012). Reaching maximum brightness at 8–9 days after the burst in the blue bands, SN 2010bh is the most rapidly evolving GRB-SNe to date. A two-component parametrized model fitted to the pseudo bolometric light curve delivered physical parameters of the explosion. The kinetic energy makes this SN the second most energetic GRB-SN after SN 1998bw. SN 2010bh also shows one of the earliest peaks ever recorded and it fades more rapidly than any other GRB-SN or type-Ic SN. Further analysis can be found in Olivares E. *et al.* (2012).

### Acknowledgements

Part of the funding for GROND (both hardware as well as personnel) was generously granted from the Leibniz-Prize to Prof. G. Hasinger (DFG grant HA 1850/28-1). This work made use of data supplied by the UK *Swift* Science Data Centre at the University of Leicester.

### References

- Bufano, F., Pian, E., Sollerman, J., *et al.* 2012, *ApJ* submitted, arXiv:1111.4527
- Campana, S., Mangano, V., Blustin, A. J., *et al.* 2006, *Nature*, 442, 1008
- Cano, Z., Bersier, D., Guidorzi, C., *et al.* 2011, *ApJ*, 740, 41
- Hjorth, J. & Bloom, J. S. 2011, in *Gamma-Ray Bursts*, ed. C. Kouveliotou, R. A. M. J. Wijers, & S. E. Woosley (Cambridge: Cambridge Univ. Press), arXiv:1104.2274
- Maeda, K., Mazzali, P. A., Deng, J., *et al.* 2003, *ApJ*, 593, 931
- Olivares E., F., Greiner, J., Schady, P., *et al.* 2012, *A&A*, 539, A76
- Starling, R. L. C., Wiersema, K., Levan, A. J., *et al.* 2011, *MNRAS*, 411, 2792
- Valenti, S., Benetti, S., Cappellaro, E., *et al.* 2008, *MNRAS*, 383, 1485
- Woosley, S. E. & Bloom, J. S. 2006, *ARAA*, 44, 507