

From the limited details given it seems that there are four new cases per year in their area. If we assume that the oldest fifth of the population is at risk and a duration of illness of 10 years, we can estimate the prevalence. Doing this reveals that our failure to find a single case in a sample of 1070 is not surprising ($P \approx 0.12$). One would have to know more about the population at risk to make more detailed calculations.

In our own study, taking the year 3 follow-up (the only one done by psychiatrists) we can confirm that no diagnosis of conversion pseudodementia was made by the psychiatrists. The AGE-CAT computerised diagnostic system does not make a diagnosis of hysteria, the nearest equivalent is the diagnosis of hypochondriacal neurosis (Larkin *et al*, 1992), and there were six cases with syndrome case levels. Of these, two were AGE-CAT organic cases, both of these had been organic cases at year 0. The remaining four had Mini-Mental State scores of 24, 26, 29 and 30. One had received an AGE-CAT diagnosis of depression at year 0, at year 3, and was depressed at year 6 follow-up. The remaining three were cases of hypochondriacal neurosis at year 3. One was depressed at year 0 and was again depressed at year 6. One was well at year 0, and a subcase of anxiety at year 6. One was a subcase of depression at year 0 and was not seen at year 6.

The whole area remains a difficult one, we note that the ECA study found an increase in cognitive impairment in people with somatisation disorder (prevalence ratio 5.94), but that this was not significant (Swartz *et al*, 1991). It would be interesting to know whether the cases of pseudodementia in the study of Liberini *et al* recovered at follow-up, perhaps the ultimate test of pseudodementia; we look forward to eventual publication of their study.

LARKIN, B. A., COPELAND, J. R. M., DEWEY, M. E., *et al* (1992) The natural history of neurotic disorder in an elderly urban population: findings from the Liverpool longitudinal study of continuing health in the community. *British Journal of Psychiatry*, **160**, 681–686.

SWARTZ, M., LANDERMAN, R., GEORGE, L. K., *et al* (1991) Somatization disorder. In *Psychiatric Disorders in America* (eds L. N. Robins & D. A. Regier), pp. 220–257. New York: The Free Press.

MICHAEL E. DEWEY
JOHN R. M. COPELAND
PAUL A. SAUNDERS

University Department of Psychiatry
University of Liverpool
PO Box 147
Liverpool L69 3BX

Multiple personality

SIR: Merskey (*Journal*, March 1992, **160**, 327–340) reviews historical literature on cases of multiple personality. He reaches the conclusion that they can be accounted for as an iatrogenic creation based on (a) misdiagnosis of organic or bipolar illness, (b) the conscious development of fantasies, (c) hysterical amnesia and retraining, and (d) explicit or implicit hypnotic suggestion or other forms of therapist persuasion. The article is more like one side of a debate than a scientific review of the literature. In every case in which there is insufficient information, a common problem in reviewing historical case material, the author concludes that the possibility of contamination by therapist suggestion is indeed a probability. The author seems to take particular delight in highlighting the 'embarrassing failure' of American psychiatrists to make a proper diagnosis, a deficiency remedied when the hapless patient is fortunate enough to return to Great Britain.

Dr Merskey acknowledges that many of the patients in these studies have various dissociative symptoms such as dissociative amnesia. If they do in fact have real dissociative symptoms, then the disagreement is really about phenomenology, i.e. a failure of integration of memory, rather than a more pervasive failure of integration of aspects of identity.

While such scepticism can help to strengthen diagnostic procedures, this article constitutes an unusual attack on the diagnostic abilities of psychiatrists who make the diagnosis of MPD. It is true that patients prone to dissociative disorders are generally highly hypnotisable and therefore unusually vulnerable to direct or indirect suggestion. Nonetheless, to argue that all of this form of psychopathology can be accounted for by misdiagnosis, and accidental or deliberate suggestion, is at variance with a wealth of data which suggests otherwise. The author concedes in his diatribe the existence of genuine dissociative psychopathology independent of iatrogenesis.

We intend to include a tightened version of the diagnosis of multiple personality disorder in the fourth edition of the *Diagnostic and Statistical Manual* of the American Psychiatric Association. In addition to some changes in language, we are re-introducing the requirement that the dissociative amnesia be a part of the syndrome in order to make the diagnosis. We strongly encourage further research in this complex but important area of psychopathology.

DAVID SPIEGEL
Stanford University School of Medicine
Stanford
California 94305
USA