

The Journal of Laryngology and Otology

EDITED BY
A. LOGAN TURNER AND J. S. FRASER

Contents

- THE SURGICAL TREATMENT OF SUPPURATIVE AND CERTAIN NON-SUPPURATIVE AFFECTIONS OF THE LABYRINTH SIR WILLIAM MILLIGAN
- SEPTIC OTITIC THROMBOSIS OF THE CRANIAL BLOOD SINUSES AND JUGULAR BULB J. S. FRASER
- THE ESTABLISHMENT OF A SPECIAL DEPARTMENT J. B. CAVENAGH
- CLINICAL RECORD:—
MUCOCELE OF THE FRONTAL SINUS CAUSED BY A LARGE OSTEOMA WALTER G. HOWARTH
- SOCIETIES' PROCEEDINGS:—
ROYAL SOCIETY OF MEDICINE—SECTION OF OTOTOLOGY.
- ABSTRACTS:—
THE EAR. THE NOSE. THE LARYNX. LOCAL ANÆSTHETICS.
- A DUAL-PURPOSE WOOL CARRIER B. SEYMOUR JONES
- REVIEWS OF BOOKS. LETTERS TO THE EDITORS. GENERAL NOTES.

Edinburgh: Oliver and Boyd
London: 33 Paternoster Row

Monthly, 4/- net

Annual Subscription, 40/- net

"We have experimentally proved that Listerine is a powerful antiseptic, preventing the development of bacteria and decomposition of vegetable infusions."—*British Medical Journal*, May 3, 1890.

LISTERINE

*A safe, non-poisonous, unirritating
antiseptic solution*

LISTERINE is particularly useful in the treatment of abnormal conditions of the mucosa, and admirably suited for a wash, gargle, or douche, in catarrhal conditions of the nose and throat.

LISTERINE in proper dilution may be freely and continuously used without prejudicial effect, either by injection or spray, in all the natural cavities of the body.

LISTERINE embodies a two-fold antiseptic effect, in that after the evaporation of its volatile constituents—thyme, eucalyptus, mentha, gaultheria, and ethyl alcohol—a film of boracic and benzoic acids remains upon the surface to which Listerine has been applied, affording more prolonged antiseptic protection.

LISTERINE is a trustworthy surgical dressing; it has no injurious effect upon the tissues in which the healing process is going on.

LISTERINE in teaspoonful doses will often afford relief in fermentative dyspepsia, and is largely prescribed, with excellent results, in the various forms of diarrhœa, occurring in infants and adults.

LISTERINE literature, including special pamphlets upon *Respiratory Diseases*, may be had, by physicians, upon application to

LAMBERT PHARMACAL COMPANY

2101 Locust Street

St Louis, Mo., U.S.A.

British Agents:

S. Maw, Son & Sons, 7 Aldersgate St., London, E.C.

The Journal of Laryngology and Otology

(Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

A. LOGAN TURNER AND J. S. FRASER

ASSISTED BY

DOUGLAS GUTHRIE AND IRWIN MOORE

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked and no additional matter should be added.

4. The order for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied, on which the price of the reprints is stated.

5. Authors of Original Communications on Oto-Laryngology in other Journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

6. Editorial Communications should be addressed to "EDITORS, *Journal of Laryngology*, c/o Messrs OLIVER AND BOYD, Tweeddale Court, Edinburgh."

7. The Annual Subscription is Forty Shillings, post free, and is payable in advance.

8. Single copies of back numbers, both of the present and the previous series, are on sale at Four Shillings each.

9. An Annual Subscription can be commenced with any number of the Journal, and the preceding numbers of the current year can be purchased at Four Shillings each.

UNITED STATES OF AMERICA

Orders for this Journal, for single parts or yearly subscriptions, can be sent either through a local bookseller, or to Messrs G. E. STECHERT & CO., 31-33 East 10th Street, New York.

CONTENTS

	PAGE
THE SURGICAL TREATMENT OF SUPPURATIVE AND CERTAIN NON-SUPPURATIVE AFFECTIONS OF THE LABYRINTH. Sir William Milligan, M.D.	245
SEPTIC OTIC THROMBOSIS OF THE CRANIAL BLOOD SINUSES AND JUGULAR BULB. J. S. Fraser, M.B., F.R.C.S.	253
THE ESTABLISHMENT OF A SPECIAL DEPARTMENT. J. B. Cavenagh, M.B.Oxon., Worcester	262
CLINICAL RECORD :—	
MUCOCELE OF THE FRONTAL SINUS CAUSED BY A LARGE OSTEOMA. Walter G. Howarth, M.A., M.B.Camb., F.R.C.S.Eng.	265
SOCIETIES' PROCEEDINGS :—	
Royal Society of Medicine—Section of Otology	267
ABSTRACTS :—	
The Ear	279
The Nose	282
The Larynx	284
Local Anæsthetics	289
A DUAL-PURPOSE WOOL CARRIER. B. Seymour Jones, F.R.C.S.	291
REVIEWS OF BOOKS	292
LETTERS TO THE EDITORS	294
GENERAL NOTES	296

HEMOPLASTIN

(HÆMOSTATIC SERUM)

Remarkably efficacious in all forms of hæmorrhage

HEMOPLASTIN is a sterile fluid derived from normal serum of horses and cattle. It is composed principally of prothrombin and thrombokinase, together with a third substance known as anti-antithrombin. These constituents are in physiologically balanced solution ready for immediate use. It retains its properties for about three years.

HEMOPLASTIN is indicated in the treatment of all types of hæmorrhage, particularly those depending upon imperfect coagulation of the blood. It has proved of great value in pulmonary hæmorrhage, and in hæmorrhage incidental to surgical procedures, tonsillectomy, etc. Its administration to hæmophilic subjects about to undergo operations is a valuable prophylactic measure.

HEMOPLASTIN may be administered intravenously or subcutaneously. The dose is 1 to 2 c.c. every 4 to 6 hours until the hæmorrhage is controlled. The prophylactic dose is 2 c.c. one hour before the operation. Hemoplastin may also be applied on sterile gauze to bleeding surfaces.

Hemoplastin (P., D. & Co.) is supplied in bulbs containing 2 c.c.

PARKE, DAVIS & Co., 50 Beak St., **LONDON, W.1**
Regent St.,