



Research Article

Residue analysis suggests ritual use of tobacco at the ancient Mesoamerican city of Cotzumalhuapa, Guatemala

Adam Negrin^{1,2,*}, Oswaldo Chinchilla Mazariegos^{3,*}, Cameron L. McNeil²,
W. Jeffrey Hurst^{4,5} & Edward J. Kennelly¹

¹ Department of Biological Sciences, Lehman College, City University of New York, USA

² Department of Anthropology, Lehman College, City University of New York, USA

³ Department of Anthropology, Yale University, New Haven, USA

⁴ The Hershey Center for Health and Nutrition, Hershey, USA

⁵ Gretna Scientific LLC, Mount Gretna, USA

* Authors for correspondence ✉ oswaldo.chinchilla@yale.edu & adam.negrin@lehman.cuny.edu



The widespread significance of tobacco in Mesoamerica is documented in historical and ethnographic sources, yet recovery of the organic remains of this plant from archaeological contexts is rare. Here, the authors present evidence for the ritual use of tobacco at Cotzumalhuapa, Guatemala, during the Late Classic period (AD 650–950). Detection of nicotine in residue analysis of three cylindrical ceramic vases recovered from cache deposits near the El Baúl acropolis suggests that these vessels contained tobacco infusions or other liquid preparations. These results suggest an ancient ritual practice involving tobacco for which there was previously no physical evidence in Mesoamerica.

Keywords: Guatemala, Cotzumalhuapa, El Baúl, Late Classic period, LC-MS, nicotine, *Nicotiana*

Introduction

Tobacco (*Nicotiana* sp., Solanaceae) is one of the most significant and ubiquitous ritual plants of the Americas (Linton 1924; Mason 1924; Wilbert 1987, 1979; Winter 2000). Early colonial accounts and modern ethnographic sources attest to the widespread use of tobacco for religious and medicinal purposes in Mesoamerica (Sahagún 1829; Thompson 1946, 1970; Robicsek 1978; Durán 1994). In all probability, these practices are ancient but direct evidence

Received: 17 August 2022; Revised: 5 April 2023; Accepted: 25 May 2023

© The Author(s), 2024. Published by Cambridge University Press on behalf of Antiquity Publications Ltd. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

concerning the use of tobacco in archaeological contexts is elusive due to the poor preservation of organic material in much of Mesoamerica, and carbonised tobacco seeds are rarely recovered for macrobotanical analysis due to their minute size (Oyuela-Caycedo & Kawa 2015). Artistic representations and ethnographic sources indicate that the preferred method of use throughout the region is, and was, the smoking of dried tobacco leaves in cigars (Thompson 1970; Robicsek 1978), though evidence of such practices rarely survives (Domenici 2014).

In this study, we present evidence of ritual tobacco use in southern Mesoamerica. Residue analysis using liquid chromatography-mass spectrometry (LC-MS) detected nicotine in three ceramic vessels recovered from cache deposits near the El Baúl acropolis at Cotzumalhuapa, Guatemala, dating to the Late Classic Pantaleón phase (AD 650–950). The residues in these vessels raise important questions about the modes of consumption and the ritual uses of tobacco in ancient Mesoamerica.

Residue analysis of tobacco

The genus *Nicotiana* consists of more than 80 herbaceous plant species with a centre of origin in South America, though endemic species have also been identified in Africa, Australia and the South Pacific (Goodspeed 1954; Knapp *et al.* 2004). Tobacco, broadly represented by *N. tabacum* L., *N. rustica* L. and several wild species containing nicotine and other alkaloids, is found throughout the Americas and has been utilised by indigenous cultures for thousands of years (Winter 2000). *Nicotiana tabacum* and *N. rustica* are the only species known to be widely cultivated throughout North and South America (Mangelsdorf *et al.* 1964).

Residue analysis of archaeological samples can identify chemical constituents of biological material associated with artefacts. The detection of plant metabolites provides evidence about the use and contents of excavated artefacts, complementing inferences based on their shape and context. Nicotine, a pyridine alkaloid present in *Nicotiana* species, has been used as a biomarker to identify the presence of tobacco in archaeological samples. Nicotine has been detected in pipe residues from the North American Eastern Woodlands (Rafferty 2002, 2006; Freimuth *et al.* 2012), the Southeast Woodlands (Carmody *et al.* 2018), the Northwest Coast (Tushingam *et al.* 2013, 2018) and central California (Eerkens *et al.* 2012), in artefacts used for smoking and grinding tobacco from Central Chile (Echeverría *et al.* 2014), in mineralised dental plaque from central California (Eerkens *et al.* 2018), in mummified human hair from San Pedro de Atacama in Northern Chile (Echeverría & Niemeyer 2013; Niemeyer *et al.* 2018) and in a miniature vessel from the Maya Lowlands (Zagorevski & Loughmiller-Newman 2012).

Ritual deposits at Cotzumalhuapa

Cotzumalhuapa was one of the largest Late Classic cities in southern Mesoamerica (Chinchilla Mazariegos 2011, 2012). The distinctive Cotzumalhuapa sculptural style spread across the Pacific coast and central Guatemalan highlands, probably reflecting the extent of the city's cultural and political influence (Thompson 1948; Parsons 1969; Chinchilla Mazariegos 1996). Excavations conducted from 2006 to 2007 focused on an architectural group located 200m north-west of the El Baúl acropolis, dating to the Pantaleón phase (Figure 1). Labelled




Figure 1. Map of Cotzumalhuapa, showing the location of Operation EB9. The dotted line marks the estimated extent of the city in the Late Classic period (figure by authors).

as Operation EB9, these excavations were motivated by the proximity of a large deposit of obsidian debitage and were designed to investigate the functions of the buildings in this group and whether they were related to the obsidian industry.

The excavations revealed parts of two patios surrounded by buildings with stone bases, built on an artificial landfill delimited by a large retaining wall (Figure 2). The discovery of two sweat baths and 21 cache deposits under the clay floors of the buildings raised the probability that this architectural group served important ritual functions (Figure 3). The number of cache deposits in Operation EB9 is far larger than those recovered elsewhere at Cotzumalhuapa, attesting to the religious significance of this group. The objects found in these deposits include four ceramic figurines, one mushroom stone, seven small ceramic vessels, and 21 large ceramic vessels, eleven of which were covered with inverted ceramic bowls (Gómez González 2011, 2013). The predominant forms are tall cylindrical vases covered by inverted bowls that protected their contents at the time of deposition. Thirteen cache vessels contained a complete obsidian blade with no visible wear, and one vessel contained two unused blades.

Sculptures in the Cotzumalhuapa style are rich in representations of plants. Tobacco leaves are believed to be sculpted on the headdresses of two royal portraits (El Baúl monument 12 and Pantaleón monument 1) that were originally placed in the Great Precinct of El Baúl (Figure 4) (Chinchilla Mazariegos 2012). The likely presence of tobacco leaves in royal headdresses suggests that the plant was relevant for political legitimation and royal rituals at Cotzumalhuapa.

Materials and methods

Excavation and residue sampling

At the time of excavation, there was no plan to test for nicotine; nevertheless, tobacco use was prohibited during excavations and subsequent handling of the objects. The vessels were taken to the project laboratory in the Museo Popol Vuh, Guatemala City, where they were kept in a tobacco-free environment. The exteriors of the vessels were cleaned and most of the soil from the interiors was extracted. The walls and the bottoms of the vessels were not cleaned or curated to allow sampling for residue analysis. The samples consisted of very fine powder removed directly from the vessel walls. The bulk sediment was not kept and was not tested. All sampling was conducted with metal utensils, except in the case of the miniature bottle (EB9D-I27-05) where a new bamboo pick was used. The metal tools were carefully cleaned with distilled water and Kimwipes™ between each new sample. Samples were placed in four-ounce sterile Whirl-pak™ bags and exported to the United States with permission from the Instituto de Antropología e Historia de Guatemala for testing.

Residue extraction

Samples were processed in the laboratories of the Department of Biological Sciences, Lehman College of The City University of New York. Residue samples (~50–250mg) were extracted twice with 70 per cent aqueous methanol in borosilicate glass vials for 15 minutes in an ultrasonic bath. Extracts were centrifuged for four minutes at 2500 rpm and the resulting supernatants were transferred into pre-weighed glass vials, evaporated to dryness under




Figure 2. Plan of Operation EB9, showing the stone foundations of structures (Str.) and the location of caches (red dots). Red labels identify vessels that were sampled for residue analysis (figure by authors).




Figure 3. Cache vessels from Operation EB9 in situ, originally deposited below clay floors associated with the stone foundations of buildings. Trowels are oriented due north: a) EB9D-E26-04-2 and two additional cache vessels; b) EB9E-G18-05; c) EB9D-F30-03; d) EB9D-Q31-06; e) EB9E-H19-06; f) EB9D-I27-05 with three additional miniature vessels (photographs by Oswaldo Chinchilla Mazariegos).

inert nitrogen gas and stored at -20°C until analysis. Dried extracts were resuspended in 70 per cent mass spectrometry (MS) grade methanol and filtered using a $0.45\mu\text{m}$ syringe filter prior to analysis by LC-MS.



Figure 4. El Batil Monument 12, a royal portrait from Cotzumalhuapa (height 1.85m). The headdress features three ovate leaves whose shape, size and venation are suggestive of tobacco (*Nicotiana* sp., Solanaceae) (photograph by Oswaldo Chinchilla Mazariegos).

Liquid chromatography-mass spectrometry

Two analytical methods using different column chemistry were employed to assess data repeatability: reversed-phase C_{18} (RP-C18) and hydrophilic interaction liquid chromatography (HILIC). The two methods provide complementary evidence, modifying the order of gradient elution and sensitivity of the analyses. Analyses were performed on a Waters Acquity ultra performance liquid chromatography module tandem to a Waters Xevo triple-quadrupole mass spectrometer (UPLC-TQD-MS, Waters Corporation, Milford, MA). Separation by RP-C18 was achieved using a Phenomenex Kinetex UPLC C_{18} column (50×2.1 mm i.d, $1.7 \mu\text{m}$ 100\AA) with a Security Guard ULTRA guard column (Torrance, CA). The mobile phase conditions employed were as follows: MS grade 0.1 per cent formic acid (aqueous) for Solvent A, MS grade acetonitrile for Solvent B, $0.5 \text{ mL} \cdot \text{min}^{-1}$ flow rate, 40°C column temperature. Mass spectrometric parameters: 3.0 kV capillary voltage; 150°C source temperature; 450°C desolvation temperature;

nitrogen desolvation gas at $800 \text{ L} \cdot \text{h}^{-1}$; cone gas at $50 \text{ L} \cdot \text{h}^{-1}$; argon collision gas at $0.15 \text{ mL} \cdot \text{min}^{-1}$. A Waters HILIC Ethylene Bridged Hybrid (BEH) UPLC column (2.1×50 mm, $1.7 \mu\text{m}$) for nicotine detection (Dobrinas *et al.* 2011) used a $0.5 \text{ mL} \cdot \text{min}^{-1}$ flow rate.

Targeted UPLC-TQD-MS analysis employed multiple reaction monitoring (MRM) of two or more ion fragments for metabolite detection. MRMs were optimised for instrument sensitivity using reference standards, plant extracts and ion fragmentation to enable detection of compounds in minute residue samples. Theobromine and caffeine were obtained from Chromadex (Irvine, CA), and (+/-)-nicotine (>99% purity, liquid) and theophylline from Sigma Aldrich (St. Louis, MO). Methanolic extracts of *N. tabacum* and *N. rustica* (nicotine), *Capsicum* sp. (capsaicin, dihydrocapsaicin) and *Bixa orellana* (bixins) were used for MRM optimisation and chromatographic separation. MRMs for capsaicin and dihydrocapsaicin followed published parameters (Powis *et al.* 2013). The analytical method only scanned for compounds which were optimised prior to residue analysis. Other food or psychoactive species were not targeted in the current analysis.

Detection of marker ion peaks employed a 6:1 signal-to-noise ratio using a peak-to-peak algorithm. Samples with peaks detected below a 3:1 signal-to-noise ratio were reanalysed. Limits of detection for the methylxanthines (theobromine, theophylline and caffeine) using reversed-

phase C₁₈ analysis were empirically determined between 1–10 nanograms·mL⁻¹ using reference standards. Analyses of residue sample extracts were performed with blank solvent injections in between residue extract sample injections. Multiple extractions were prepared to retest residues, confirm marker compound signals detected and assess analysis repeatability.

Results

Table 1 summarises the results of LC-MS conducted on samples from seven vessels (Figure 5). Nicotine was detected in residue samples from two cylindrical vases (EB9E-G18-05 and EB9D-G27-04) and a spherical vessel (EB9E-G19-06). All three vessels contained an obsidian blade. Nicotine levels were significantly higher in the residue sample from vessel EB9E-G18-05 as compared with the two other samples (Figure 6). Nicotine was not detected in residue samples in two other cylindrical vases. There were no meaningful distinctions related to the context or mode of recovery of the three vessels that yielded positive results, except that vessels EB9E-G18-05 and EB9E-G19-06 were found in proximity to each other (1.26m apart) on the west side of Structure E, during the 2007 excavations. Vessel EB9D-G27-04 was found near Structure A in 2006.

Nicotine was not detected in residues from a miniature bottle (EB9D-I27-05). Such miniature vessels from the Maya region, called flasks or ‘poison bottles’, are believed to have contained materials used in small amounts such as perfumes, medicines, spices, poisons or pigments (Loughmiller-Cardinal & Zagorevski 2016) and are sometimes labelled as ‘tobacco houses’, suggesting that they contained tobacco snuffs (Houston *et al.* 2006; Boot 2019).

Residue analysis was optimised for the chemical detection of diagnostic compounds from several important regional crops. Three methylxanthines (theobromine, theophylline and caffeine) from *Theobroma cacao*, bixins from achiote (*Bixa orellana*) and capsaicins (capsaicin and dihydrocapsaicin) from chili peppers (*Capsicum* sp.) were not detected in residues from any of the seven sampled vessels.

Indigenous uses of tobacco

Indigenous peoples of the Americas have utilised tobacco for recreational, medicinal and religious purposes: historical and archaeological records show that smoking was the preferred method of consumption throughout the Americas before and after the arrival of Europeans (Wilbert 1987; Oyuela-Caicedo & Kawa 2015). Other forms of administration include chewing, sucking, snuffing, licking and drinking tobacco preparations (Mason 1924; Elferink 1964). Throughout North America (Winter 2000), for the Maya, and in South America (Rosengren 2006), it was believed that gods desired and fed upon tobacco in various forms. Tobacco offerings may be blown as smoke, wafted onto ritual objects (Thompson 1970), provided as burning cigars (Domenici 2014) or thrown in a formal manner within religious activity (Kroeber 1941).

Sixteenth-century reports show that tobacco was integrated into social life as part of meetings and ceremonies, used as a digestive aid after meals and considered as the proper conclusion to feasts (Durán 1994). Tobacco was smoked to increase success in hunting and travel on land and water, to mediate interactions with spirits, to engage in hospitality and friendship

Table 1. Results of residue analysis using LC-MS. N1, N2 and N3 refer to parent-daughter ion transition peaks employed for nicotine detection by multiple reaction monitoring using reversed-phase C₁₈ (RP-C18) and HILIC separation summed across multiple analyses. (+) = peak detected above a 6:1 signal-to-noise ratio; (•) = peak detected above a 3:1 signal-to-noise ratio; (-) = no peak detected.

Sample	Description	RP-C18		HILIC		
		N1	N2	N1	N2	N3
EB9D-E26-04-2	Cylindrical vessel containing obsidian blade, with bowl, found below sealed floor	+	•	-	-	-
EB9D-F30-03	Cylindrical vessel containing obsidian blade, found with bowl	•	•	-	-	-
EB9D-G27-04	Cylindrical vessel containing obsidian blade, found in exterior structure	+	+	+	+	+
EB9D-I27-05	Miniature vessel, with bowl and pitcher, found in exterior structure	-	-	-	-	-
EB9D-Q31-06	Cylindrical vessel with waterfowl design, found below sealed floor	-	-	-	-	-
EB9E-G18-05	Cylindrical vessel containing obsidian blade	+	+	+	+	+
EB9E-H19-06	Spherical vessel with bowl and obsidian blade, found below sealed floor	+	+	+	+	+

with neighbours, to eliminate fatigue as a stimulant and to combat diseases and maintain general health (McGuire 1899; Breedlove & Laughlin 1993a). Tobacco was also perceived as dangerous, requiring attention to its presence in and around homes during use, storage and preparation and in social relations. Tobacco was a talisman to protect oneself, one's property or objects, used to suppress the work of witches, thwart evil, and to imbue 'heat' and potency to ritual objects (Breedlove & Laughlin 1993b; Groark 2019). It was applied to the skin or lips for spiritual protection and power (Thompson 1970).

Pipes were not widespread in Mesoamerica (Thompson 1946, 1970; Robicsek 1978), except in Postclassic West Mexico (Lister & Howard 1955; Cabrero García 1993). The modern Ch'orti' use wooden pipes, but the antiquity of this practice is unknown (Hull 2019). More common are cigars made entirely of tobacco or wrapped with leaves of other species such as sapidilla (*Manilkara zapota*), Barbados cherry/acerola (*Malpighia glabra*), allspice pepper (*Pimenta officinalis*) and common guava (*Psidium guajava*) or smoking through maize bracts and husks or the hollow stems of reeds (Benzoni 1857; Thompson 1970). Smoking tobacco is important in religious rituals and ancient Maya deities were sometimes portrayed smoking cigars (Tozzer 1907; Robicsek 1978; Tedlock 1996; Flores & Kantun Balam 1997). Tobacco is also commonly ground for consumption as a snuff (Starr 1904). The Mexica blended ground tobacco with calcium hydroxide as slaked lime to produce a snuff known as *picietl* (Thompson 1946, 1970). The Tzeltal still carry tobacco gourds for stimulant use and to alleviate hunger and fatigue (Groark 2010, 2019). Mazatec and Maya travellers carry tobacco snuffs to protect themselves against, or to cause, witchcraft (Starr 1904; Houston *et al.* 2006).



Figure 5. Archaeological vessels sampled for residue analysis from El Baúl, Cotzumalhuapa, Guatemala. From left to right, top row: EB9E-G18-05, EB9D-G27-04, EB9E-H19-06, EB9D-I27-05; bottom row: EB9D-Q31-06, EB9D-E26-04-2, EB9D-F30-03 (photographs by Oswaldo Chinchilla Mazariegos).




Figure 6. Chromatograms of (A) (+/-)-nicotine reference standard and (B) (+/-)-nicotine detected in sample EB9E-G18-05 using reversed-phase C_{18} chromatography UPLC-TQD-MS analysis. Chromatograms of (C) nicotine standard, (D) blank solvent injection (95% MS-grade acetonitrile) and (E) nicotine detected in sample EB9E-G18-05 using HILIC chromatography and multiple reaction monitoring of three parent-daughter ion transitions (figure by authors).

While the consumption of tobacco in liquid form or as concentrated syrups is not the most common method of use throughout the Americas, there is some documentation for these practices in the Amazon and the Guianas (Wilbert 1987). Because oral intake of high quantities of nicotine is toxic and potentially lethal, such utilisation highlights tobacco use by ritual practitioners as narcotics to induce deep sleep, visions and divinatory trances (Monardes 1580; Elferink 1983). The use of tobacco as a psychoactive by the Maya was not documented by early explorers (Thompson 1970; Elferink 1983), but there are ethnographic testimonies of its use in historical and modern Maya rituals and healing practices. The Ch'orti' Maya of Guatemala apply tobacco juices to communicate with spirits and diagnose disease based on bodily responses to their inquiries (Wisdom 1940). Sixteenth-century Nahuatl sources describe *teotlaqualli* (food of god) as a black ointment preparation of venomous animals, ground tobacco and psychoactive ololiuqui seeds (*Rivea corymbosa*, Convolvulaceae), used in ceremonial offerings and priestly functions to communicate with spirits and achieve fearlessness during sacrifices (Elferink 1999). Bernardino de Sahagún mentioned *yiaqualli*, made of tobacco, soot and a plant similar to henbane (*Hyoscyamus niger*, Solanaceae) possibly used as a psychoactive (Elferink 1999). In Tlaxcala, large vases of *picietl* were placed among temple altar offerings and monitored for the appearance of an animal or eagle footprint in the ground tobacco powder as divination by priests (Muñoz Camargo 1984).

The few documented preparations of tobacco in liquid form in Mesoamerica are medicinal (Monardes 1580; Sahagún 1829; Benzoni 1857; Breedlove & Laughlin 1993a). Drinking of tobacco juices in various forms (red, white and black) is mentioned as a cure for asthma in *The Ritual of the Bacabs* (Roys 1965; Thompson 1970) and for gastrointestinal treatments in modern ethnography (Berlin *et al.* 1990). Modern healing practices among highland Tzeltal-Tzotzil Maya of Chiapas, Mexico (Breedlove & Laughlin 1993b; Groark 2010), include preparations of *Nicotiana* spp. (*moy*, *moytik*, *bankilal*, *moy pox* or *yanal moy*) rubbed topically on the body, consumed in solid form or as teas prepared with water or cane liquor mixed with ash or slaked lime, in bathing, as plasters applied to the body for ritual cleansing (Wasson 1963), and for skin parasite removal (Houston *et al.* 2006). Tobacco extracted in cold or warm water may be blended with dried chili peppers (*Capsicum* spp.), garlic or other components rendering them ‘hot’. Vessel imagery suggests that the Maya utilised aqueous tobacco extracts with possible plant admixtures as ritualistic enemas, psychoactive substances or medicine (Robicsek 1978; de Smet & Hellmuth 1986). Beer and wine made from palm, jocote fruit and honey were occasionally prepared with tobacco in the Maya lowlands (Dahlin & Litzinger 1986) and the West Indies (Gage 1677), and used as yeast inocula for fermentation, deterring unwanted bacterial/fungal growth (Litzinger 1983).

Discussion

The function of ceramic vessels is normally inferred from their shape, size and archaeological context (Lesure 1998; Rice 2015). Mesoamerican cylindrical vases are commonly believed to have held liquids that were consumed directly from the vases or else poured into smaller vessels for drinking. In the Maya Lowlands, many cylindrical vases have inscriptions that identify various kinds of cacao or maize beverages as their intended contents, though they could also have served for other drinks (Hall *et al.* 1990; Reents-Budet 1994; Stuart 2009; Beliaev *et al.* 2010). The spherical vase (EB9E-G19-06), a variation of this shape, could have been used to serve and drink beverages. The detection of nicotine within residue analyses of three cylindrical vases from Cotzumalhuapa was therefore unexpected, suggesting the use of tobacco in liquid form and within vessels larger than miniature flasks, or ‘tobacco houses’, that are more usually associated with tobacco use (Houston *et al.* 2006; Boot 2019).

Vessels recovered from cache deposits served a terminal function as containers for offerings, forming part of ritual activities that concluded with their deposition and burial (Becker 1993; Lucero 2010). The ritual use of tobacco has been documented among the ancient and modern Mesoamerican peoples alongside cacao, flowers, esteemed objects (Sahagún 1829; Thompson 1970; Robicsek 1978), food, drink and other substances as offerings for the earth, for specific deities and even for the buildings within which the offering were cached, as buildings were conceived as animate beings that could cause harm to their occupants if not properly appeased (Vogt 1976). The cylindrical vases from Cotzumalhuapa may therefore have contained tobacco-infused offerings.

The proximity of sweat baths in the same architectural group as the cache deposits containing cylindrical vases at Cotzumalhuapa suggests that the tobacco infusions deposited in these vessels may have been employed in curing and purification rituals. The EB9 sweat baths consist of small spaces with stone floors laid out as concave surfaces (in one case shaped

as a bathtub), with traces of fireplaces and burning. The drainage canals that are common in Mesoamerican sweat baths are absent, but the size, shape and layout of the spaces suggests that they functioned as sweat baths (Chinchilla Mazariegos 2011).

The sweat bath is especially important in therapeutic and ritual procedures related to childbirth and is associated with deities related to midwifery (Alcina Franch 2000; Chinchilla Mazariegos 2017). Archaeological and ethnographic evidence attest to the use of sweat baths in Mexico and Guatemala for spiritual and physical purification (Cosminsky 2001). In the Maya highlands, steam bathing is used for many health conditions. In Oxchuc, ritual steam bathing and bloodletting are used to cure diseases, while tobacco-wound coverings protect against the entrance of “pathogenic wind” (Groark 1997: 58). Uncured tobacco powders are orally consumed. The *temazcal* (sweat lodge) is used in Tzotzil apprenticeships, bonesetting and midwifery. Midwives employ steam bathing for cleansing, massage, pathogen removal and elimination, infection prevention, removal of bad spirits from the body and purification of the baby (Cosminsky 2001). Among other possibilities, one potential use of the obsidian blades found in the Cotzumalhuapa vessels is for cutting the umbilical cord, as observed among the Huichol and Tarahumara until the mid-twentieth century (Huber & Sandstrom 2001).

Conclusion

Previous work on the chemical detection of tobacco has focused on miniature flasks. Our study shows that nicotine can also be found in larger-volume vessels, providing novel contextual data suggesting ritual tobacco use at Cotzumalhuapa during the Late Classic period. Such chemical markers detected in residues could also reflect substances placed in the vessels prior to their terminal ritualistic use. Nevertheless, the presence of nicotine in three of the vessels that were examined suggests a pattern in the ritual activities that led to their deposition, rather than a casual utilisation of vessels that had previously contained tobacco. The presence of nicotine in these cache vessels is significant considering the ritual importance of tobacco in ancient and modern Mesoamerica and the rarity of its recovery in excavated contexts. This study highlights important questions about the uses of tobacco in religious rituals, while the proximity of the deposits to sweat baths documented in the same architectural group at Cotzumalhuapa suggests associations with curative and purification practices, including maternal care and childbirth. These implications warrant further research at Cotzumalhuapa and other Mesoamerican sites.

Acknowledgements

Institutional and logistical support was provided by the Museo Popol Vuh, Universidad Francisco Marroquín, Yale University, Lehman College, The City University of New York and the agroindustrial company Pantaleón, S.A.

Funding statement

Excavations at Cotzumalhuapa were possible thanks to funding provided by the Wenner-Gren Foundation for Anthropological Research (Grant 6787), the National

Geographic Society (Grant 7958-05) and the Panamerican Institute of Geography and History (Project 2.1.3.4.21).

References

- ALCINA FRANCH, J. 2000. *Temazcalli: higiene, terapéutica, obstetricia y ritual en el Nuevo Mundo*. Sevilla: Escuela de Estudios Hispano-Americanos de Sevilla.
- BECKER, M.J. 1993. Earth offerings among the Classic Period Lowland Maya: burials and caches as ritual deposits, in E.C. Danien & R.J. Sharer (ed.) *New theories on the Ancient Maya*: 185–96. Philadelphia: University Museum, University of Pennsylvania.
- BELIAEV, D., A. DAVLETSHIN & A. TOKOVININE. 2010. Sweet cacao and sour atole: mixed drinks on Classic Maya ceramic vases, in J.E. Staller & M. Carrasco (ed.) *Pre-Columbian foodways*: 257–72. New York: Springer.
- BENZONI, G. 1857 (trans 2010). *History of the New World*. Translated by W.H. Smyth. Cambridge: Cambridge University Press. <https://doi.org/10.1017/CBO9780511694615>
- BERLIN, B., E. BERLIN, D. BREEDLOVE, T. DUNCAN, V. JARA, R. LAUGHLIN & T. VELASEO. 1990. *La herbolaria médica Tzeltal-Tzotzil en los altos de Chiapas*. Chiapas: Gobierno del Estado de Chiapas, Consejo Estatal de Fomento a la Investigación y Difusión de la Cultura, DIF-Chiapas, and Instituto Chiapaneco de Cultura, Tuxtla Gutiérrez.
- BOOT, E. 2019. Tobacco as mentioned in hieroglyphic texts on Classic Maya pottery flasks and in the Late Postclassic codices: an essay on the hieroglyphic signs used, in J.A. Loughmiller-Cardinal & K. Eppich (ed.) *Breath and smoke: tobacco use among the Maya*: 157–81. Albuquerque: University of New Mexico Press.
- BREEDLOVE, D.E. & R.M. LAUGHLIN. 1993a. *The flowering of man: a Tzotzil botany of Zinacantán, volume 1* (Smithsonian Contributions to Anthropology). Washington, D.C.: Smithsonian Institution Press.
- 1993b. *The flowering of man: a Tzotzil botany of Zinacantán, volume 2* (Smithsonian Contributions to Anthropology). Washington, D.C.: Smithsonian Institution Press.
- CABRERO GARCÍA, M.T. 1993. Hallazgos recientes en el Cañón de Bolaños, Jalisco. *Anales de Antropología* 30: 47–72.
- CARMODY, S., J. DAVIS, S. TADI, J. SHARP, R. HUNT & J. RUSS. 2018. Evidence of tobacco from a Late Archaic smoking tube recovered from the Flint River site in southeastern North America. *Journal of Archaeological Science: Reports* 21: 904–10. <https://doi.org/10.1016/j.jasrep.2018.05.013>
- CHINCHILLA MAZARIEGOS, O. 1996. Settlement patterns and monumental art at a major Pre-Columbian polity: Cotzumalguapa, Guatemala. Unpublished PhD dissertation, Vanderbilt University.
- 2011. The obsidian workshop of El Baúl, Cotzumalhuapa, in Z.X. Hruby, G. Braswell & O. Chinchilla Mazariegos (ed.) *The technology of Maya Civilization: political economy and beyond in lithic studies*: 102–118. London: Equinox.
- 2012. *Cotzumalguapa: la ciudad arqueológica: El Baúl, Bilbao, El Castillo*. Guatemala: F&G Editores.
- 2017. *Art and myth of the Ancient Maya*. New Haven (CT): Yale University Press.
- COSMINSKY, S. 2001. Maya midwives of southern Mexico and Guatemala, in B.R. Huber & A.R. Sandstrom (ed.) *Mesoamerican healers*: 179–210. Austin: University of Texas Press.
- DAHLIN, B.H. & W.J. LITZINGER. 1986. Old bottle, new wine: the function of Chultuns in the Maya Lowlands. *American Antiquity* 51: 721–36. <https://doi.org/10.2307/280861>
- DE SMET, P.A. & N.M. HELLMUTH. 1986. A multidisciplinary approach to ritual enema scenes on ancient Maya pottery. *Journal of Ethnopharmacology* 16: 213–62. [https://doi.org/10.1016/0378-8741\(86\)90091-7](https://doi.org/10.1016/0378-8741(86)90091-7)
- DOBRINAS, M., E. CHOONG, M. NOETZLI, J. CORNUZ, N. ANSERMOT & C.B. EAP. 2011. Quantification of nicotine, cotinine, trans-3'-hydroxycotinine and varenicline in human plasma by a sensitive and specific UPLC–tandem mass-spectrometry procedure for a clinical study on smoking cessation. *Journal of Chromatography B* 879: 3574–82. <https://doi.org/10.1016/j.jchromb.2011.09.046>
- DOMENICI, D. 2014. Cueva del Lazo: child sacrifice or special funerary treatment? Discussion of a Late Classic context from the Zoque region of western

- Chiapas (Mexico), in G.D. Wrobel (ed.) *The bioarchaeology of space and place*: 39–75. New York: Springer.
https://doi.org/10.1007/978-1-4939-0479-2_3
- DURÁN, D. 1994. *The history of the Indies of New Spain*. Translated by D. Heyden. Norman: University of Oklahoma Press.
- ECHEVERRÍA, J. & H.M. NIEMEYER. 2013. Nicotine in the hair of mummies from San Pedro de Atacama (Northern Chile). *Journal of Archaeological Science* 40: 3561–8.
<https://doi.org/10.1016/j.jas.2013.04.030>
- ECHEVERRÍA, J., M.T. PLANELLA & H.M. NIEMEYER. 2014. Nicotine in residues of smoking pipes and other artifacts of the smoking complex from an Early Ceramic period archaeological site in central Chile. *Journal of Archaeological Science* 44: 55–60.
<https://doi.org/10.1016/j.jas.2014.01.016>
- EERKENS, J., D. ARDURA, O. FIEHN, J. BLAKE, K. LENTZ, S. TUSHINGHAM & M. PALAZOGLU. 2012. GC/MS analysis of residues reveals nicotine in two Late Prehistoric pipes from CA-ALA-554. *Proceedings of the Society for California Archaeology* 26: 212–19.
- EERKENS, J.W. *et al.* 2018. Dental calculus as a source of ancient alkaloids: detection of nicotine by LC-MS in calculus samples from the Americas. *Journal of Archaeological Science: Reports* 18: 509–515.
<https://doi.org/10.1016/j.jasrep.2018.02.004>
- ELFERINK, J. 1964. Some uses of tobacco among the Maya. *Notes on Middle American Archaeology and Ethnology* 3: 1–5.
- 1983. The narcotic and hallucinogenic use of tobacco in Pre-Columbian Central America. *Journal of Ethnopharmacology* 7: 111–122.
[https://doi.org/10.1016/0378-8741\(83\)90084-3](https://doi.org/10.1016/0378-8741(83)90084-3)
- 1999. Teotlaqualli: the psychoactive food of the Aztec gods. *Journal of Psychoactive Drugs* 31: 435–40.
<https://doi.org/10.1080/02791072.1999.10471773>
- FLORES, J.S. & J. KANTUN BALAM. 1997. Importance of plants in the Ch’a Chaak Maya ritual in the Peninsula of Yucatan. *Journal of Ethnobiology* 17: 97–108.
- FREIMUTH, G.A., S.U. WISSEMAN & A.V. ULANOV. 2012. Analysis of pipe residue by gas chromatography/mass spectroscopy (GC/MS). *Illinois Archaeology* 24: 184–92.
- GAGE, T. 1677. *A new survey of the West-Indies: or the English American his travail by sea and land*. London: A. Clark.
- GÓMEZ GONZÁLEZ, E.M. 2011. Las ofrendas de Cotzumalguapa, Guatemala, en el Clásico Tardío (650/700–1000 d.C.). Unpublished MA dissertation, Universidade de Trás-os-Montes e Alto Douro.
- 2013. Las ofrendas de un pequeño conjunto arquitectónico en Cotzumalguapa, in B. Arroyo & L. Méndez Salinas (ed.) *XXVI simposio de investigaciones arqueológicas en Guatemala, 2012*: 899–910. Guatemala: Museo Nacional de Arqueología y Etnología.
- GOODSPEED, T.H. 1954. The genus *Nicotiana*; origins, relationships and evolution of its species in the light of their distribution, morphology and cytogenetics. Waltham (MA): Chronica Botanica.
- GROARK, K.P. 1997. To warm the blood, to warm the flesh: the role of the steambath in Highland Maya (Tzeltal-Tzotzil) ethnomedicine. *Journal of Latin American Lore* 20: 3–96.
- 2010. The angel in the gourd: ritual, therapeutic, and protective uses of tobacco (*Nicotiana tabacum*) among the Tzeltal and Tzotzil Maya of Chiapas, Mexico. *Journal of Ethnobiology* 30: 5–30.
<https://doi.org/10.2993/0278-0771-30.1.5>
- 2019. “Elder brother tobacco”: traditional *Nicotiana* snuff use among the contemporary Tzeltal and Tzotzil Maya of Highland Chiapas, Mexico, in J.A. Loughmiller-Cardinal & K. Eppich (ed.) *Breath and smoke: tobacco use among the Maya*. Albuquerque: University of New Mexico Press.
- HALL, G.D., S.M. TARKA, W.J. HURST, D. STUART & R.E. ADAMS. 1990. Cacao residues in ancient Maya vessels from Rio Azul, Guatemala. *American Antiquity* 55: 138–43.
<https://doi.org/10.2307/281499>
- HOUSTON, S., D. STUART & K. TAUBE. 2006. *The memory of bones: body, being, and experience among the Classic Maya*. Austin: University of Texas Press.
- HUBER, B.R. & A.R. SANDSTROM. 2001. Recruitment, training, and practice of indigenous midwives from the Mexico–United States border to the isthmus of Tehuantepec, in B.R. Huber & A.R. Sandstrom (ed.) *Mesoamerican healers*: 139–78. Austin: University of Texas Press.

- HULL, K. 2019. The smoking of bones Ch'orti' Maya use of tobacco and ritual tobacco substitutes, in J. Loughmiller-Cardinal & K. Eppich (ed.) *Breath and smoke: tobacco use among the Maya*: 126–56. Albuquerque: University of New Mexico Press.
- KNAPP, S., M.W. CHASE & J.J. CLARKSON. 2004. Nomenclatural changes and a new sectional classification in *Nicotiana* (Solanaceae). *Taxon* 53: 73–82. <https://doi.org/10.2307/4135490>
- KROEBER, A.L. 1941. *Salt, dogs, tobacco*. Berkeley: University of California Press.
- LESURE, R.G. 1998. Vessel form and function in an Early Formative ceramic assemblage from coastal Mexico. *Journal of Field Archaeology* 25: 19–36. <https://doi.org/10.1179/jfa.1998.25.1.19>
- LINTON, R. 1924. Use of tobacco among North American Indians. *Anthropology Leaflet* 15: 1–27.
- LISTER, R.H. & A.M. HOWARD. 1955. The Chalchihuites Culture of Northwestern Mexico. *American Antiquity* 21: 122–9. <https://doi.org/10.2307/276854>
- LITZINGER, W.J. 1983. The ethnobiology of alcoholic beverage production by the Lacandon, Tarahumara, and other Aboriginal Mesoamerican peoples. Unpublished PhD dissertation, University of Colorado at Boulder.
- LOUGHMILLER-CARDINAL, J.A. & D. ZAGOREVSKI. 2016. Maya flasks: the “home” of tobacco and godly substances. *Ancient Mesoamerica* 27: 1–11. <https://doi.org/10.1017/S0956536116000079>
- LUCERO, L.J. 2010. Materialized cosmology among Ancient Maya commoners. *Journal of Social Archaeology* 10: 138–67. <https://doi.org/10.1177/1469605309354403>
- MANGELSDORF, P.C., R.S. MACNEISH & G.S. WILLEY. 1964. Origins of agriculture in Middle America, in R.C. West (ed.) *Handbook of Middle American Indians, volume 1*. Austin: University of Texas Press. <https://doi.org/10.7560/732599-013>
- MASON, J.A. 1924. *Use of tobacco in Mexico and South America*. Chicago (IL): Field Museum of Natural History.
- MCGUIRE, J.D. 1899. *Pipes and smoking customs of the American Aborigines, based on material in the US National Museum*. Washington, D.C.: US Government Printing Office.
- MONARDES, N. 1580. *Primera y segunda y tercera partes de la historia medicinal de las cosas que se traen de nuestras Indias Occidentales que sirven en medicina*. Sevilla: En casa de Fernando Diaz.
- MUÑOZ CAMARGO, D. 1984. Descripción de la ciudad y provincia de Tlaxcala de las Indias y del Mar Océano para el buen gobierno y ennoblecimiento dellas, in R. Acuña (ed.) *Relaciones geográficas del Siglo XVI: Tlaxcala, tomo primero*. Mexico City: Universidad Nacional Autónoma de México.
- NIEMEYER, H.M., P. DE SOUZA, C. CAMILO & J. ECHEVERRÍA. 2018. Chemical evidence of prehistoric passive tobacco consumption by a human perinate (early Formative Period, South-Central Andes). *Journal of Archaeological Science* 100: 130–8. <https://doi.org/10.1016/j.jas.2018.10.010>
- OYUELA-CAYCEDO, A. & N.C. KAWA. 2015. A deep history of tobacco in lowland South America, in A. Russel & E. Rahman (ed.) *The master plant: tobacco in lowland South America*: 27–44. London: Bloomsbury Academic.
- PARSONS, L.A. 1969. *Bilbao, Guatemala: an archaeological study of the Pacific Coast Cotzumalhuapa Region, vol. 2*. Milwaukee (WI): Milwaukee Public Museum.
- POWIS, T.G., E.G. MURRIETA, R. LESURE, R.L. BRAVO, L. GRIVETTI, H. KUCERA & N.W. GAIKWAD. 2013. Prehispanic use of chili peppers in Chiapas, Mexico. *PLoS ONE* 8: e79013. <https://doi.org/10.1371/journal.pone.0079013>
- RAFFERTY, S.M. 2002. Identification of nicotine by gas chromatography/mass spectroscopy analysis of smoking pipe residue. *Journal of Archaeological Science* 29: 897–907. <https://doi.org/10.1006/jasc.2001.0747>
- 2006. Evidence of early tobacco in Northeastern North America? *Journal of Archaeological Science* 33: 453–8. <https://doi.org/10.1016/j.jas.2005.08.006>
- REENTS-BUDET, D. 1994. *Painting the Maya universe*. Durham (NC): Duke University Press.
- RICE, P.M. 2015. *Pottery analysis: a sourcebook*. Chicago (IL): University of Chicago Press.
- ROBICSEK, F. 1978. *The smoking gods: tobacco in Maya art, history, and religion*. Norman: University of Oklahoma Press.
- ROSENGREN, D. 2006. Transdimensional relations: on human-spirit interaction in the Amazon.

- Journal of the Royal Anthropological Institute* 12: 803–16.
<https://doi.org/10.1111/j.1467-9655.2006.00364.x>
- ROYS, R.L. 1965. *Ritual of the Bacabs*. Norman: University of Oklahoma Press.
- SAHAGÚN, B.D. 1829. *Historia general de las cosas de Nueva España*. México: Imprenta del Ciudadano Alejandro Valdés.
- STARR, F. 1904. *Notes upon the ethnography of Southern Mexico. Expedition of 1901*. Davenport (IA): Davenport Academy of Sciences.
- STUART, D. 2009. The language of chocolate: references to cacao on Classic Maya drinking vessels, in C. McNeil (ed.) *Chocolate in Mesoamerica: a cultural history of cacao*: 184–201. Gainesville: University Press of Florida.
- TEDLOCK, D. 1996. *Popol Vuh: the Mayan book of the dawn of life*. New York: Simon and Schuster.
- THOMPSON, J.E.S. 1946. Some uses of tobacco among the Maya. *Notes on Middle American Archaeology and Ethnology* 3: 1–5.
- 1948. *An archaeological reconnaissance in the Cotzumalhuapa Region, Escuintla, Guatemala*. (Contributions to American Anthropology and History series 574). Washington, D.C.: Carnegie Institution of Washington.
- 1970. *Maya history and religion*. Norman: University of Oklahoma Press.
- TOZZER, A.M. 1907. *A comparative study of the Mayas and the Lacandonese*. London: The Macmillan Company.
- TUSHINGHAM, S., D. ARDURA, J.W. EERKENS, M. PALAZOGLU, S. SHAHBAZ & O. FIEHN. 2013. Hunter-gatherer tobacco smoking: earliest evidence from the Pacific Northwest Coast of North America. *Journal of Archaeological Science* 40: 1397–407.
<https://doi.org/10.1016/j.jas.2012.09.019>
- TUSHINGHAM, S., C.M. SNYDER, K.J. BROWNSTEIN, W.J. DAMITIO & D.R. GANG. 2018. Biomolecular archaeology reveals ancient origins of indigenous tobacco smoking in North American Plateau. *Proceedings of the National Academy of Sciences USA* 115: 11742–7.
<https://doi.org/10.1073/pnas.1813796115>
- VOGT, E.Z. 1976. *Tortillas for the gods: a symbolic analysis of Zinacanteco rituals*. Cambridge (MA): Harvard University Press.
- WASSON, R.G. 1963. Notes on the present status of ololiuhqui and the other hallucinogens of Mexico. *Botanical Museum Leaflets, Harvard University* 20: 161–93.
- WILBERT, J. 1979. Magico-religious use of tobacco among South American Indians, in D.L. Browman & R.A. Schwarz (ed.) *Spirits, shamans, and stars*: 13–38. The Hague: De Gruyter Mouton.
- 1987. *Tobacco and shamanism in South America*. New Haven (CT): Yale University Press.
- WINTER, J.C. 2000. *Tobacco use by Native North Americans: sacred smoke and silent killer*. (The Civilization of the American Indian series 236). Norman: University of Oklahoma Press.
- WISDOM, C. 1940. *The Chorti Indians of Guatemala*. Chicago (IL): University of Chicago Press.
- ZAGOREVSKI, D.V. & J.A. LOUGHMILLER-NEWMAN. 2012. The detection of nicotine in a Late Mayan period flask by gas chromatography and liquid chromatography mass spectrometry methods. *Rapid Communications in Mass Spectrometry* 26: 403–11. <https://doi.org/10.1002/rcm.5339>