Article: EPA-1025 Topic: P29 - Psychotherapy

INTEGRATIVE PSYCHOTHERAPY MODEL OF ANXIETY DISORDERS

B. Mykhaylov¹, O. Kudinova¹

¹Psychotherapy, Kharkov Medical Academy of Postgraduate Education, Kharkov, Ukraine

At the present time inUkraine the special priority has the problem of anxiety and somatoform disorders. A prevalence of clinical anxiety and phenomena of somatization neurotic disturbances in general and necessity of differential diagnostics with somatic diseases were the precondition for studying this area.

At the same time, in Ukrainethe diagnosis 'Vegetative-vascular dystonia' which is ciphered G 90.8, according to ICD-10 instead of 'Neurotic stress-induced and Somatoform disorders'F40-F48 is used. It leads unreasonable treatment significant contingents of neurotic patients in neurological departments.

The 60 agoraphobia (F 40.0)patients, 40 panic disorder (PD) and 40 somatoform vegetative dysfunction of cardiovascular system integrative (SVDCS) (F45.30) patients were examined. The emotional disturbances of all thease patients were investigated with the clinicand pathopsychological methods. The high level of anxiety was shown. Pathopsychologically the high level of personality anxiety and low level of neurotism and depression were indentified. The system of psychotherapy correction was established, based on integrativeprinciples, with the suggestive, cognitive-behavioral, autogenic trainingtechniques correspondence.

Our experience showed thenecessity of the integrative models of psychotherapy provided, parted on stage. On the first stage – sedative-adapting the receptions of cognitive and suggestive psychotherapy are used. There is groupe therapy on second-main-stage. On the third stage-pesupportive elements of the autogenic training mastered.

Clinical approbationintegrative psychotherapy system shown the 82% high, 12%-middle,6%- lowefficacy.