



Interplay of combat deployment harassment, testosterone concentrations and post-deployment suicide risk in male veterans


Original Article

Cite this article: Sher L, Bierer LM, Flory J, Makotkine I, and Yehuda R. (2024) Interplay of combat deployment harassment, testosterone concentrations and post-deployment suicide risk in male veterans. *Acta Neuropsychiatrica* 1–5. doi: [10.1017/neu.2024.12](https://doi.org/10.1017/neu.2024.12)

Received: 9 January 2024
Revised: 3 March 2024
Accepted: 20 March 2024

Keywords:
Veterans; harassment; suicide; testosterone

Corresponding author:
Leo Sher; Email: Leo.Sher@mssm.edu

Leo Sher^{1,2} , Linda M. Bierer^{1,2}, Janine Flory^{1,2}, Iouri Makotkine^{1,2} and Rachel Yehuda^{1,2}

¹James J. Peters VA Medical Center, New York, USA and ²Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA

Abstract

Objective: Many combat veterans exhibit suicidal ideation and behaviour, but the relationships among experiences occurring during combat deployment and suicidality are still not fully understood. In this study, we tested the hypothesis that harassment during a combat deployment is associated with post-deployment suicidality and testosterone function. *Methods:* Male combat veterans who made post-deployment suicide attempts and demographically matched veterans without a history of suicide attempts were enrolled in the study. Demographic and clinical parameters of study participants were assessed and recorded. Study participants were interviewed by a trained clinician using the Mini-International Neuropsychiatric Interview (MINI), the Deployment Risk and Resilience Inventory (DRRI) – Relationships within unit scale, the Scale for Suicidal Ideation (SSI), and the Brown–Goodwin Aggression Scale. Free testosterone levels were assessed in morning blood samples. *Results:* DRRI harassment scores were higher and free testosterone levels were lower among suicide attempters in comparison with non-attempters. In the whole sample, DRRI harassment scores positively correlated with SSI scores and negatively correlated with free testosterone levels. Free testosterone levels negatively correlated with SSI scores. Aggression scale scores positively correlated with DRRI harassment scores among non-attempters but not among attempters. *Conclusion:* Our observations that harassment scores are associated with suicidality and testosterone levels, and suicidality is associated with testosterone levels may indicate that there is a link between deployment harassment, testosterone function and suicidality.

Significant outcomes

- More severe harassment during a combat deployment was associated with higher post-deployment suicidality.
- More severe harassment during a combat deployment was associated with lower testosterone levels.
- Lower free testosterone levels were associated with higher suicidality.

Limitations

- A modest sample size.
- Only combat veterans who volunteered to participate in the research project were included in the study.

Introduction

Studies suggest that many combat veterans exhibit suicidal ideation and behaviour (Kang and Bullmann, 2008; Pietrzak *et al.*, 2010; Sher and Yehuda, 2011; Cigrang *et al.*, 2015; Denneson *et al.*, 2015; Monteith *et al.*, 2018; Reger *et al.*, 2018; Sher, 2024). A study of about 300 Iraq/Afghanistan war veterans reported that 12.5% had contemplated suicide in the 2 weeks preceding the survey (Pietrzak *et al.*, 2010). One study found a higher suicide rate in service members who were currently deployed or previously deployed compared with those who had never been deployed (Schoenbaum *et al.*, 2014). A more recent study found that there was an increased risk of suicide when all Iraq/Afghanistan war veterans were compared to the US population (Bullman and Schneiderman, 2021). Both male and female veterans had an increased risk of suicide when compared to their gender-specific non-veteran counterparts.



After 15 years of follow-up, Iraq/Afghanistan war veterans continued to have a suicide rate that exceeds that of the US population.

The relationship between combat deployments and suicidality are not fully understood. Studies suggest that the deployment environment may affect post-deployment suicide risk (Lemaire & Graham, 2011; Monteith *et al.*, 2018). For example, it has been observed that experiences of harassment are associated with suicidal ideation in Iraq/Afghanistan war veterans (Lemaire & Graham, 2011). Exposure to harassment, that is, to unwanted negative behaviour that is intended to cause damage and/or is perceived as unpleasant, hostile and damaging is a psychosocial factor which frequently has powerful negative effects on mental health, including suicidality (Hom *et al.*, 2017; Mitchell *et al.*, 2021; Campbell-Sills *et al.*, 2023; Thomas *et al.*, 2023).

Research observations indicate that high levels of endogenous testosterone promote behaviour intended to dominate (Mazur and Booth, 1998; Dreher *et al.*, 2016; Inoue *et al.*, 2023). Research findings link testosterone with both aggressive and non-aggressive status-seeking. A fraction of testosterone circulates as non-protein-bound or free testosterone. Only this free fraction is biologically active. Therefore, it is possible that men with higher free testosterone levels are less likely to be harassed.

In this study, we tested a hypothesis that harassment during a combat deployment is associated with post-deployment suicidality and testosterone function.

Materials and methods

Fifteen male combat veterans who made post-deployment suicide attempts within 5 years preceding the day of initial evaluation and 17 demographically matched veterans without a history of suicide attempts were enrolled in our study. Suicide attempts were defined as a potentially self-injurious behaviour for which there is evidence that the individual probably planned to kill himself/herself. Veterans without a history of suicide attempts were selected from the same group of individuals with psychiatric disorders as suicide attempters at the James J. Peters VA Medical Center (JJP VAMC). At a time deemed clinically appropriate, JJP VAMC inpatient, outpatient or emergency department clinical staff working with the veteran introduced the study. Inclusion criteria included age 18–65 years and being able to provide informed consent. Exclusion criteria included primary psychotic disorder and major medical or neurological illness. All veterans who participated in the study had a physical examination and laboratory tests, including toxicological screenings to exclude medical or neurological abnormalities that could influence their psychiatric state.

Veterans were informed that study participation was completely voluntary and choosing to participate or to decline participation will not affect VA clinical treatment or services. Written informed consent was obtained prior to study enrolment. The research project was approved by the Institutional Review Board at the JJP VAMC.

Study participants were interviewed by a trained clinician using the *Mini-International Neuropsychiatric Interview (MINI)* (Sheehan *et al.*, 1998) to establish DSM-IV diagnoses, the *Deployment Risk and Resilience Inventory (DRRI) – Relationships within unit scale (DRRI)* (King *et al.*, 2006) to examine harassment during a combat deployment, the *Scale for Suicidal Ideation (SSI)* (Beck *et al.*, 1979) to examine suicidal ideation, and the *Brown–Goodwin Aggression Scale* (Brown and Goodwin, 1986) to assess the severity of aggression, a characteristic often associated with suicidal behavior.

Fasting blood samples were drawn between 8:00 and 8:30 am. Free testosterone kits utilise a competitive immunoassay created and validated for the *in vitro* diagnostic determination of testosterone in human blood. The free testosterone assay sensitivity is 0.018 pg/mL. The intra-assay and inter-assay coefficients of variation for the free testosterone assay are 8.1% and 6.9%, respectively. The interval between the interviews and the blood tests did not exceed 2 weeks.

Demographic, clinical and biological data were compared using Student's *t*-test, chi-square test, and their association evaluated using correlations, as appropriate. We used the *t*-test to analyse continuous variables. The chi-square test was employed to test for differences in proportions between the groups. We used correlations to examine the linear relationship between two continuous variables. DRRI harassment scores and morning free testosterone levels were also compared between attempters and non-attempters by means of a general linear model, with 'Group' as fixed factor and 'DRRI harassment scores' or 'testosterone levels' as dependent factor. 'Mood disorders', 'post-traumatic stress disorder (PTSD)', 'substance use disorders' and 'aggression severity' were included in these analyses as covariates. The SPSS 27 program was used for statistical analysis.

Results

There were no differences regarding demographic parameters because the groups were demographically matched (Table 1). SSI scores were higher among veterans with a history of suicide attempts in comparison with non-attempters (Table 1).

DRRI harassment scores were higher among suicide attempters in comparison with non-attempters (Table 1). The difference between attempters and non-attempters in DRRI harassment scores remained significant after the adjustment for mood disorders, PTSD, substance use disorders, and aggression severity ($df = 1,26$, $F = 5.62$, $p = 0.025$). In the whole sample, SSI scores positively correlated with DRRI harassment scores ($n = 32$, $r = 0.40$, $p = 0.02$).

In the whole sample, morning free testosterone levels negatively correlated with DRRI harassment scores ($n = 32$, $r = -0.36$, $p = 0.04$) and with SSI scores ($n = 32$, $r = -0.57$, $p < 0.001$). Morning free testosterone levels were lower in suicide attempters in comparison with non-attempters (Table 1). The difference between attempters and non-attempters in testosterone levels remained significant after the adjustment for mood disorders, PTSD, substance use disorders and aggression severity ($df = 1,26$, $F = 8.43$, $p = 0.007$).

Aggression scale scores positively correlated with DRRI harassment scores among non-attempters ($n = 17$, $r = 0.56$, $p = 0.02$) but not among attempters ($n = 15$, $r = -0.19$, $p = 0.49$).

Discussion

Harassment and suicidality

We have found that higher deployment harassment scores are associated with post-deployment suicide attempts and greater post-deployment suicidal ideation. Our findings are consistent with research reports suggesting that various forms of harassment are associated with suicidality in veteran and non-veteran populations (Lemaire & Graham, 2011; Hom *et al.*, 2017; Griffith, 2019; Thomas *et al.*, 2023). For example, a cross-sectional review of mental health evaluations of about 2,000 Iraq/Afghanistan war veterans showed that general (non-sexual) harassment was associated with suicidal ideation (Lemaire & Graham, 2011). A study of about 13,000 US soldiers found that

Table 1. Demographic and clinical features of combat veterans with or without a history of suicide attempts

Parameter	Attempters		Non-attempters		Analysis		
	Mean or <i>N</i>	SD or %	Mean or <i>N</i>	SD or %	<i>t</i> / χ^2 / <i>F</i>	df	<i>p</i>
Age (years)	38.33	12.02	35.71	10.83	-0.65	30	0.52
Marital status (% married)	4	26.7%	8	47.1%	1.41	1	0.23
% of subjects who completed at least 4-year college	6	40.0%	8	47.1%	0.16	1	0.69
% of subjects with mood disorders	14	93.3%	16	94.1%	0.008	1	0.93
% of subjects with post-traumatic stress disorder	13	86.7%	11	64.7%	0.52	1	0.15
% of subjects with substance use disorder	13	86.7%	11	64.7%	0.52	1	0.15
Beck Suicide Ideation Scale at the time of study entry	21.0 (<i>R</i> = 29)*	7.37	2.00 (<i>R</i> = 23)*	5.78	-8.17	30	<0.001
Brown-Goodwin Aggression Scale	24.53 (<i>R</i> = 25)*	7.80	22.59 (<i>R</i> = 32)*	9.17	-0.64	30	0.53
DRRI - Relationships within unit	30.27 (<i>R</i> = 38)*	10.41	22.06 (<i>R</i> = 22)*	7.18	-2.62	30	0.014
Morning free testosterone levels (pg/mL)	11.09	7.33	18.67	6.34	30	3.14	0.004

**R*, range.

sexual harassment was associated with a fivefold increase for risk of suicide (Griffith, 2019).

A feature of military life is its institutional nature (Griffith, 2019). Life in the military takes place nearly completely in one setting. In comparison with the civilian population, military units are closed systems. This can be an additional stress on targets of harassment because they have limited opportunity to obtain support outside the military unit.

Harassment frequently involves an individual being put in a position where this person has problems defending him/herself. Harassed individuals frequently feel helpless, stressed out, anxious, and hopeless and expect to become subject of further negative treatment. It has been suggested that lower resilience and higher hopelessness may mediate the relationships between harassment and suicidality (Livingston *et al.*, 2022). Stress and anxiety may also increase suicide risk.

Testosterone

We have observed that free testosterone levels negatively correlated with harassment scores. This may indicate that individuals with higher testosterone levels are less likely to be harassed. Indeed, studies suggest that being victorious in competitive circumstances, including situations involving aggression, can result in the heightened testosterone levels, and this can accrue across recurrent victories to raise the testosterone levels of more successful men (Carre and Archer, 2018). Studies and theoretical papers suggest that testosterone mediates diverse types of status-seeking behaviour, augmenting competitive or aggressive behaviour, and stimulating prosocial behaviour to attain and preserve social status or dominance (Mehta and Beer, 2010; Terburg *et al.*, 2016; Carre and Archer, 2018). Repeated social defeat decreases testosterone levels, subsequently decreasing dominance drive. Social anxiety is associated with lower testosterone levels (Hutschemaekers *et al.*, 2020). Harassment can be regarded as a state of social defeat associated with anxiety. Therefore, harassment may reduce testosterone levels. It should be noted that studies indicate that there is a significant individual and situational variability in the testosterone-behaviour connection. For example, studies have

shown that both low and high testosterone levels may be associated with suicidality (Sher, 2023). Variations in the results of these studies may be related to differences in patient populations and the research methodology.

Our observations that harassment scores are associated with suicidality and testosterone levels and suicidality is associated with testosterone levels may indicate that there is a link between deployment harassment, testosterone function and suicidality.

Aggression

We found that that aggression scale scores positively correlated with harassment scores among suicide non-attempters but not among attempters. This indicates differences in the psychobiological regulation between suicide attempters and non-attempters and is consistent with some previous observations (Sachs-Ericsson *et al.*, 2014; Sher *et al.*, 2020; Sher *et al.*, 2021). For example, we have previously observed that plasma neuropeptide Y (NPY) levels positively correlated with aggression scale scores among suicide attempters but not among non-attempters (Sher *et al.*, 2020).

Limitations

A modest sample size is a limitation of this study. Also, only combat veterans who volunteered to participate in the research project were included in the study. This could influence the outcomes of the study. Differences in combat experiences, severity of psychiatric conditions and medication use could also affect the results of the study.

Conclusion

Multiple psychological and biological factors affect suicidality in both civilians and military veterans (Lemaire & Graham, 2011; Rihmer and Gonda, 2012; Elman *et al.*, 2013; van Heeringen and Mann, 2014; Ejdesgaard *et al.*, 2015; Shen *et al.*, 2016; Sher, 2023; Department of Veterans Affairs, 2022). Harassment is one of these factors and an important issue in the military. It can have significant negative effects on psychological health and lead to suicidal behaviour. Prevention of harassment in the military is a critical task.

Commanders can deter harassing behaviour because they observe and control military personnel (Gilberd, 2018; Griffith, 2019). This shows the importance of selections and effective training of military leaders which affects a lot of individuals and processes that make up the Armed Forces. Effective military leaders are expected to value their subordinates and protect them from harassment. If harassment complaints are substantiated, military leadership is expected to take appropriate disciplinary or administrative action against the harasser (Gilberd, 2018).

The results of our study are relevant to clinical practice and research. Our results suggest that including assessments of deployment harassment may have clinical utility for understanding the degree of suicide risk in veterans. Our study indicates that combat veterans with a history of deployment harassment need to be frequently screened for suicidality. The adequate treatment for psychiatric and medical disorders including targeted psychotherapeutic interventions for trauma, depression, and other symptoms, and effective social assistance may mitigate negative long-term effects of deployment harassment. The results of our study may also contribute to the development of biomarkers for suicidal behaviour among military veterans.

Future studies of harassment in the military should investigate possible mediators in the association of harassment with suicidal behaviour such as shame, perceived burdensomeness, and social alienation. Also, we need studies of what psychotherapeutic and psychopharmacological interventions for victims of harassment in the military are the most effective.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/neu.2024.12>.

Author contribution. Authors LS and RY designed the study and wrote the protocol. Author LS managed the literature searches and undertook the statistical analysis. All authors contributed to and have approved the final manuscript.

Financial support. This study was supported by a Lightfighter Trust Foundation Grant to Prof. Leo Sher.

Competing interests. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation.

References

- Beck AT, Kovacs M and Weissman A (1979) Assessment of suicidal intention: the scale for suicide ideation. *Journal of Consulting and Clinical Psychology* 47(2), 343–352.
- Brown GL and Goodwin FK (1986) Human aggression and suicide. *Suicide and Life-Threatening Behavior* 16(2), 223–243.
- Bullman T and Schneiderman A (2021) Risk of suicide among U.S. veterans who deployed as part of operation enduring freedom, operation Iraqi freedom, and operation new dawn. *Injury Epidemiology* 8(1), 40.
- Campbell-Sills L, Sun X, Kessler RC, Ursano RJ, Jain S and Stein MB (2023) Exposure to bullying or hazing during deployment and mental health outcomes among US Army soldiers. *JAMA Network Open* 6(1), e2252109. DOI: 10.1001/jamanetworkopen.2022.52109.
- Carre JM and Archer J (2018) Testosterone and human behavior: the role of individual and contextual variables. *Current Opinion in Psychology* 19, 149–153.
- Cigrang JA, Balderrama-Durbin C, Snyder DK, Talcott GW, Tatum J, Baker M, Cassidy D, Sonnek S, Smith Slep AM and Heyman RE (2015) Predictors of suicidal ideation across deployment: a prospective study. *Journal of Clinical Psychology* 71(9), 828–842.
- Dennison LM, Teo AR, Ganzini L, Helmer DA, Bair MJ and Dobscha SK (2015) Military veterans' experiences with suicidal ideation: implications for intervention and prevention. *Suicide and Life-Threatening Behavior* 45(4), 399–414.
- Department of Veterans Affairs (2022) Office of Mental Health and Suicide Prevention, 2022 National Veteran Suicide Prevention Annual Report. September 2022 Available at https://www.mentalhealth.va.gov/suicide_prevention/data.asp.
- Dreher JC, Dunne S, Pazderska A, Frodl T, Nolan JJ and O'Doherty JP (2016) Testosterone causes both prosocial and antisocial status-enhancing behaviors in human males. *Proceedings of the National Academy of Sciences of the USA* 13(41), 11633–11638. DOI: 10.1073/pnas.1608085113. Epub 2016-09-26.
- Ejdesgaard BA, Zöllner L, Jensen BF, Jørgensen HO and Kähler H (2015) Risk and protective factors for suicidal ideation and suicide attempts among deployed Danish soldiers from 1990 to 2009. *Military Medicine* 180(1), 61–67.
- Elman I, Borsook D and Volkow ND (2013) Pain and suicidality: insights from reward and addiction neuroscience. *Progress in Neurobiology* 109, 1–27. DOI: 10.1016/j.pneurobio.2013.06.003. Epub 2013-07-01.
- Gilberd K (2018) New military policy on harassment. *Military Law Task Force* <https://nlgmtf.org/military-law/2018/new-military-policy-on-harassment/>
- Griffith J (2019) The sexual harassment-suicide connection in the U.S. Military: contextual effects of hostile work environment and trusted unit leaders. *Suicide and Life-Threatening Behavior* 49(1), 41–53.
- Hom MA, Stanley IH, Spencer-Thomas S and Joiner TE (2017) Women firefighters and workplace harassment: associated suicidality and mental health sequelae. *Journal of Nervous and Mental Disease* 205(12), 910–917.
- Hutschemaekers MHM, de Kleine RA, Davis ML, Kampman M, Smits JAJ and Roelofs K (2020) Endogenous testosterone levels are predictive of symptom reduction with exposure therapy in social anxiety disorder. *Psychoneuroendocrinology* 115, 104612.
- Inoue Y, Burriss RP, Hasegawa T and Kiyonari T (2023) Testosterone promotes dominance behaviors in the ultimatum game after players' status increases. *Scientific Reports* 13(1), 18029. DOI: 10.1038/s41598-023-45247-4.
- Kang HK and Bullman TA (2008) Risk of suicide among US veterans after returning from the Iraq or Afghanistan war zones. *JAMA* 300(6), 652–653.
- King LA, King DW, Vogt DS, Knight JA and Samper R (2006) Deployment risk and resilience inventory: a collection of measures for studying deployment-related experiences of military personnel and veterans. *Military Psychology* 18(2), 89–120.
- Lemaire CM and Graham DP (2011) Factors associated with suicidal ideation in OEF/OIF veterans. *Journal of Affective Disorders* 130(1-2), 231–238.
- Livingston WS, Tannahill HS, Meter DJ, Fargo JD and Blais RK (2022) The association of military sexual harassment/assault with suicide ideation, plans, attempts, and mortality among US service members/veterans: A meta-analysis. *Trauma Violence & Abuse* 24(4), 2616–2629.
- Mazur A and Booth A (1998) Testosterone and dominance in men. *Behavioral and Brain Sciences* 21(3), 353–363.
- Mehta PH and Beer J (2010) Neural mechanisms of the testosterone-aggression relation: the role of orbitofrontal cortex. *Journal of Cognitive Neuroscience* 22(10), 2357–2368.
- Mitchell KJ, Jones LM and Turner HA (2021) Past year technology-involved peer harassment victimization and recent depressive symptoms and suicide ideation among a National sample of youth. *Journal of Interpersonal Violence* 36(3–4), NP1165–1179NP. DOI: 10.1177/0886260517748413.
- Monteith LL, Hoffmire CA, Holliday R, Park CL, Mazure CM and Hoff RA (2018) Do unit and post-deployment social support influence the association between deployment sexual trauma and suicidal ideation? *Psychiatry Research* 270, 673–681.
- Pietrzak RH, Goldstein MB, Malley JC, Rivers AJ, Johnson DC and Southwick SM (2010) Risk and protective factors associated with suicidal ideation in veterans of operations enduring freedom and Iraqi freedom. *Journal of Affective Disorders* 123(1-3), 102–107.
- Reger MA, Tucker RP, Carter SP and Ammerman BA (2018) Military deployments and suicide: a critical examination. *Perspectives on Psychological Science* 13(6), 688–699.
- Rihmer Z and Gonda X (2012) Predisposition for self-destruction? Affective temperaments as a suicide risk factor in patients with mood disorders. *Crisis* 33(6), 309–312. DOI: 10.1027/0227-5910/a000192.

- Sachs-Ericsson N, Hames JL, Joiner TE, Corsentino E, Rushing NC, Palmer E, Gotlib IH, Selby EA, Zarit S and Steffens DC** (2014) Differences between suicide attempters and nonattempters in depressed older patients: depression severity, white-matter lesions, and cognitive functioning. *The American Journal of Geriatric Psychiatry* **22**(1), 75–85.
- Schoenbaum M, Kessler RC, Gilman SE, Colpe LJ, Heeringa SG, Stein MB, Ursano RJ, Cox KL and Army STARRS Collaborators** (2014) Predictors of suicide and accident death in the army study to assess risk and resilience in servicemembers (Army STARRS): results from the army study to assess risk and resilience in servicemembers (Army STARRS). *JAMA Psychiatry* **71**(5), 493–503.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R and Dunbar GC** (1998) The mini-international neuropsychiatric interview (M.I.N.I.): development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* **59**(Suppl 20), 22–33.
- Shen YC, Cunha JM and Williams TV** (2016) Time-varying associations of suicide with deployments, mental health conditions, and stressful life events among current and former US military personnel: a retrospective multivariate analysis. *Lancet Psychiatry* **3**(11), 1039–1048.
- Sher L** (2023) Testosterone and suicidal behavior in bipolar disorder. *International Journal of Environmental Research and Public Health* **20**(3), 2502.
- Sher L** (2024) The uncounted casualties of war: suicide in combat veterans. *QJM: An International Journal of Medicine* **117**(3), 163–167. DOI: [10.1093/qjmed/hcad240](https://doi.org/10.1093/qjmed/hcad240). Epub ahead of print.
- Sher L, Bierer LM, Flory J, Makotkine I and Yehuda R** (2020) Neuropeptide Y plasma levels and suicidal behavior in combat veterans. *European Neuropsychopharmacology* **40**, 31–37.
- Sher L, Bierer LM, Makotkine I and Yehuda R** (2021) The effect of oral dexamethasone administration on testosterone levels in combat veterans with or without a history of suicide attempt. *Journal of Psychiatric Research* **143**, 499–503. DOI: [10.1016/j.jpsychires.2020.11.034](https://doi.org/10.1016/j.jpsychires.2020.11.034). Epub 2020-11-20.
- Sher L and Yehuda R** (2011) Preventing suicide among returning combat veterans: a moral imperative. *Military Medicine* **176**(6), 601–602.
- Terburg D, Syal S, Rosenberger LA, Heany SJ, Stein DJ and Honk Jv** (2016) Testosterone abolishes implicit subordination in social anxiety. *Psychoneuroendocrinology* **72**, 205–211.
- Thomas S, Hummel KV, Schäfer J, Wittchen HU and Trautmann S** (2023) Harassment and its association with depressive symptoms and suicidal behavior: the role of perceived stigma and nondisclosure. *Psychological Services* **20**(1), 84–93.
- van Heeringen K and Mann JJ** (2014) The neurobiology of suicide. *Lancet Psychiatry* **1**(1), 63–72. DOI: [10.1016/S2215-0366\(14\)70220-2](https://doi.org/10.1016/S2215-0366(14)70220-2). Epub 2014-06-04.