

This Section of *Epidemiology and Psychiatric Sciences* regularly appears in each issue of the Journal to describe relevant studies investigating the relationship between neurobiology and psychosocial psychiatry in major psychoses. The aim of these Editorials is to provide a better understanding of the neural basis of psychopathology and clinical features of these disorders, in order to raise new perspectives in every-day clinical practice.

Paolo Brambilla, *Section Editor* and Michele Tansella, *Editor EPS*

## Diffusion imaging studies of white matter integrity in bipolar disorder

M. Bellani<sup>1\*</sup> and P. Brambilla<sup>2,3</sup>

<sup>1</sup> *Section of Psychiatry and Clinical Psychology, Department of Public Health and Community Medicine, Inter-University Center for Behavioural Neurosciences (ICBN), University of Verona, Verona, Italy*

<sup>2</sup> *DISM, Inter-University Center for Behavioural Neurosciences (ICBN), University of Udine, Udine, Italy*

<sup>3</sup> *IRCCS "E. Medea" Scientific Institute, Udine, Italy*

Diffusion tensor imaging (DTI) is a neuroimaging technique with a potential to elucidate white matter integrity. Recently, it has been used in the field of psychiatry to further understand the pathophysiology of major diseases, including bipolar disorder (BD). This review sought to focus on existing DTI findings on white matter organization in BD.

*Received 17 December 2010; Revised 28 December 2010; Accepted 28 December 2010*

**Key words:** Bipolar disorder, magnetic resonance imaging (MRI), white matter.

In the last decade, several efforts have been made in order to detect possible biomarkers of bipolar disorder (BD) and recent advances in neuroimaging research have pointed out the putative role of white matter (Arnone *et al.* 2008; Doron & Gazzaniga 2008). In this regard, structural magnetic resonance imaging (MRI) studies described diffuse cortical and callosal white matter pathology in BD patients (Adler *et al.* 2006; Kempton *et al.* 2008; Vita *et al.* 2009), suggesting the presence of altered intra- and inter-hemispheric connectivity (Atmaca *et al.* 2007; Bellani *et al.* 2009a, b). A relatively recent application of MRI is the diffusion tensor imaging (DTI), a non-invasive imaging technique that measures the motion of water molecules in brain tissue and provides information about the microstructural coherence of white matter (Basser & Jones 2002). Water diffusion within the neural tissue

is impeded by a myelin sheath, axonal density and thickness, and cellular structures. The degree of water molecule diffusion can be quantified by the apparent diffusion coefficient (ADC), and the fibre tract directionality can be evaluated indirectly by fractional anisotropy (FA; Beaulieu, 2002). ADC and FA may be considered as good markers of white matter microstructure organization. In particular, low FA values indicate probable damage to axonal membrane, de-/dys-myelination or reduced amount of intra-axonal structures, whereas high ADC measures are found when water diffusion is unimpeded, e.g. in ventricles or demyelinated white matter (Beaulieu & Allen, 1994). Several DTI data on BD have been published so far, showing some white matter abnormalities (Brambilla *et al.* 2009a). Most studies (see Table 1) reported reduced FA and/or elevated ADC values compared to healthy controls involving specific brain regions such as prefrontal, parietal, temporal and occipital lobes, internal capsule, uncinate fasciculus, superior longitudinal fasciculus and corpus callosum (for extensive review see Bellani *et al.* 2009a, b; Heng *et al.* 2010). However, the effect of mood states on

\* Address for correspondence: Dr M. Bellani, Section of Psychiatry and Clinical Psychology, Department of Public Health and Community Medicine, University of Verona, Piazzale L.A. Scuro 10, 37134 Verona, Italy.

(Email: marcella.bellani@univr.it; paolo.brambilla@uniud.it)

**Table 1.** DTI studies in bipolar disorder

Study	Subjects	Age (years)	Findings in BD
Adler <i>et al.</i> (2004)	9 BD 9 HC	32 ± 8 31 ± 7	↓FA in superior prefrontal cortex, but not in the fronto-polar regions
Beyer <i>et al.</i> (2005)	14 BD 21 HC	44.0 ± 17.6 44.6 ± 13.5	↑ADC in orbital frontal white matter bilaterally, with normal FA
Haznedar <i>et al.</i> , (2005)	34 BD 33 HC	43.9 ± 10.0 40.8 ± 11.7	↓FA in posterior limb of the internal capsule
Regenold <i>et al.</i> , (2006)	8 BD 8 HC	58.4 ± 12.9 54.5 ± 12.8	↑ADC in combined white matter ROIs, but not in the four lobes separately
Yurgelun-Todd <i>et al.</i> (2007)	11 BD 10 HC	32.9 ± 10.5 32.4 ± 9.1	↑FA in the genu of corpus callosum
Wang <i>et al.</i> (2008a)	33 BD 40 HC	32 ± 10.1 29.2 ± 9.2	↓FA in anterior cingulum white matter
Wang <i>et al.</i> (2008b)	42 BD 42 HC	32.6 ± 10.1 28.7.2 ± 9.1	↓FA in genu, rostral body, and anterior midbody of corpus callosum
Versace <i>et al.</i> (2008)	31 BD 25 HC	35.9 ± 8.9 29.48 ± 9.43	↑FA in left uncinate fasciculus, left optic radiation, and right anterothalamic radiation ↓FA in right uncinate fasciculus
Bruno <i>et al.</i> (2008)	36 BD 28 HC	37.4 42.8	↓FA in inferior longitudinal fasciculus
Barnea-Goraly <i>et al.</i> (2009)	21 BD 18 HC	16.1 ± 2.7 14.5 ± 2.7	↓FA in fornix, posterior cingulate, corpus callosum and parietal and temporal corona radiate
Chaddock <i>et al.</i> (2009)	19BD 21 first degree Relatives 18 HC	43.3 ± 10.2 42.5 ± 13.6 41.7 ± 12.2	↓FA in genu of corpus callosum, right inferior and left superior longitudinal fasciculus
Mahon <i>et al.</i> (2009)	30 BD 38 HC	33.4 ± 8.7 31.9 ± 8.6	↓FA in the left cerebellum ↑FA in left frontal white matter and thalamic radiation
Sussmann <i>et al.</i> (2009)	42 BD 28 SCZ 38 HC	39.6 ± 10.1 38.0 ± 9.9 37.2 ± 11.9	↓FA in the left anterior limb of the internal capsule, bilateral uncinate fasciculus, anterior/superior thalamic radiation, and corpus callosum
Wessa <i>et al.</i> (2009)	22 BD 21 HC	45.41 ± 12.60 42.95 ± 13.17	↑FA in medial frontal, precentral, inferior parietal, and occipital white matter
Zanetti <i>et al.</i> (2009)	37 BD 26 HC	34.1 ± 9.0 28.8 ± 9.5	↓FA in inferior fronto-occipital, superior and inferior longitudinal fasciculus
Benedetti <i>et al.</i> (2011)	40 BD 21 HC	45.11 ± 9.82 39.86 ± 11.05	↓FA in the genu of corpus callosum and in anterior and right superior-posterior corona radiate

ADC, apparent diffusion coefficient; FA, fractional anisotropy; BD, bipolar disorder; HC, healthy controls.

white matter integrity is often not taken into account (Adler *et al.* 2004; Bruno *et al.* 2008; Mahon *et al.* 2009) or patients with different clinical states are just studied together (Beyer *et al.* 2005; Versace *et al.* 2008; Wang *et al.* 2008a, b; Barnea-Goraly *et al.* 2009). In this regard, patients suffering from different bipolar episode may be characterized by specific DTI 'signature'. Anyway, till date, few studies have been conducted on patients with homogeneous mood state. In euthymia, FA is usually increased in the genu of corpus callosum, internal capsule, anterior thalamic radiation and uncinate fasciculus (Haznedar *et al.* 2005; Yurgelun-Todd *et al.* 2007; Sussmann *et al.* 2009; Wessa *et al.* 2009; Zanetti *et al.* 2009), whereas in bipolar depression lower FA has been shown in the genu of

the corpus callosum and in corona radiata (Regenold *et al.* 2006; Chaddock *et al.* 2009; Benedetti *et al.* 2011). Not surprisingly, in mixed samples higher and lower FA values were found in different brain regions (Beyer *et al.* 2005; Wang *et al.* 2008a, b; Barnea-Goraly *et al.* 2009).

The impact of mood stabilizers on white matter connectivity in BD should also be considered, particularly lithium and quetiapine, which have been suggested to potentially induce myelination processes (Bearden *et al.* 2008; Zhang *et al.* 2008; Brambilla *et al.* 2009b; Tondo & Baldessarini, 2009). However, a recent DTI study did not find any lithium effect on DTI measures in patients suffering from BD (Benedetti *et al.* 2011).

In summary, the DTI literature on BD suggests loss of white matter network connectivity as a possible

phenomenon of the disease, particularly including altered fronto-occipital, superior longitudinal fasciculus and callosal connections (Wang *et al.* 2008a; Barnea-Goraly *et al.* 2009; Chaddock *et al.* 2009). However, diffusion-imaging studies in BD are mostly limited by heterogeneity and relatively small size of the samples. Moreover, the impact of some clinical variables on white matter coherence such as mood states and mood stabilizer administration still needs to be fully elucidated. In this perspective, future DTI studies are expected to further investigate whether abnormal white matter may represent a trait or a mood state biomarker of BD, potentially being preserved by psychotropic drugs such as lithium.

## References

- Adler CM, Del Bello MP, Strakowski SM (2006). Brain network dysfunction in bipolar disorder. *CNS Spectrum* **11**, 312–320.
- Adler CM, Holland SK, Schmithorst V, Wilke M, Weiss KL, Pan H, Strakowski SM (2004). Abnormal frontal white matter tracts in bipolar disorder, a diffusion tensor imaging study. *Bipolar Disorders* **6**, 197–203.
- Arnone D, McIntosh AM, Chandra P, Ebmeier KP (2008). Meta-analysis of magnetic resonance imaging studies of the corpus callosum in bipolar disorder. *Acta Psychiatrica Scandinavica* **118**, 357–362.
- Atmaca M, Ozdemir H, Yildirim H (2007). Corpus callosum areas in first episode patients with bipolar disorder. *Psychological Medicine* **37**, 699–704.
- Barnea-Goraly N, Chang K.D, Karchemskiy A, Howe ME, Reiss AL (2009). Limbic and corpus callosum aberrations in adolescents with bipolar disorder: a tract-based spatial statistics analysis. *Biological Psychiatry* **66**, 238–244.
- Basser PJ, Jones DK (2002). Diffusion-tensor MRI, theory, experimental design and data analysis, a technical review. *NMR Biomedicine* **15**, 456–467.
- Bearden CE, Thompson PM, Dutton RA, Frey BN, Peluso MA, Nicoletti M, Dierschke N, Hayashi KM, Klunder AD, Glahn DC, Brambilla P, Sassi RB, Mallinger AG, Soares JC (2008). Three-dimensional mapping of hippocampal anatomy in unmedicated and lithium-treated patients with bipolar disorder. *Neuropsychopharmacology* **33**, 1229–1238.
- Beaulieu C (2002). The basis of anisotropic water diffusion in the nervous system, a technical review. *NMR Biomedicine* **15**, 4354–4355.
- Beaulieu C, Allen PS (1994). Determinants of anisotropic water diffusion in nerves. *Magnetic Resonance Medicine* **31**, 394–400.
- Bellani M, Marzi CA, Brambilla P (2009a). Interhemispheric communication in schizophrenia. *Epidemiologia e Psichiatria Sociale* **18**, 19–22.
- Bellani M, Yeh PH, Tansella M, Balestrieri M, Soares JC, Brambilla P (2009b). DTI studies of corpus callosum in bipolar disorder. *Biochemical Society Transactions* **37**, 1096–1098.
- Benedetti F, Yeh PH, Bellani M, Radaelli D, Nicoletti MA, Poletti S, Falini A, Dallaspezia S, Colombo C, Scotti G, Smeraldi E, Soares JC, Brambilla P (2011). Disruption of white matter integrity in bipolar depression as a possible structural marker of illness. *Biological Psychiatry* **69**, 299–300.
- Beyer JL, Taylor WD, MacFall JR, Kuchibhatla M, Payne ME, Provenzale JM, Cassidy F, Krishnan KR (2005). Cortical white matter microstructural abnormalities in bipolar disorder. *Neuropsychopharmacology* **30**, 2225–2229.
- Brambilla P, Bellani M, Yeh PH, Soares JC (2009b). Myelination in bipolar patients and the effects of mood stabilizers on brain anatomy. *Current Pharmacological Design* **15**, 2632–2636.
- Brambilla P, Bellani M, Yeh PH, Soares JC, Tansella M (2009a). White matter connectivity in bipolar disorder. *International Review Psychiatry* **21**, 380–386.
- Bruno S, Cercignani M, Ron MA (2008). White matter abnormalities in bipolar disorder: a voxel-based diffusion tensor imaging study. *Bipolar Disorders* **10**, 460–468.
- Chaddock CA, Barker GJ, Marshall N, Schulze K, Hall MH, Fern A, Walshe M, Bramon E, Chitnis XA, Murray R, McDonald C (2009). White matter microstructural impairments and genetic liability to familial bipolar disorder. *British Journal of Psychiatry* **194**, 527–534.
- Doron KW, Gazzaniga MS (2008). Neuroimaging techniques offer new perspectives on callosal transfer and interhemispheric communication. *Cortex* **44**, 1023–1029.
- Haznedar MM, Roversi F, Pallanti S, Baldini-Rossi N, Schnur DB, Licalzi EM, Tang C, Hof PR, Hollander E, Buchsbaum MS (2005). Fronto-thalamo-striatal gray and white matter volumes and anisotropy of their connections in bipolar spectrum illnesses. *Biological Psychiatry* **57**, 733–742.
- Heng S, Song AW, Sim K (2010). White matter abnormalities in bipolar disorders: insights from diffusion tensor imaging studies. *Journal of Neural Transmission* **117**, 639–654.
- Kempton MJ, Geddes JR, Ettinger U, Williams SC, Grasby PM (2008). Meta-analysis, database, and meta-regression of 98 structural imaging studies in bipolar disorder. *Archives of General Psychiatry* **65**, 1017–1032.
- Mahon K, Wu J, Malhotra AK, Burdick KE, De Rosse P, Ardekani BA, Szeszko PR (2009). A voxel-based diffusion tensor imaging study of white matter in bipolar disorder. *Neuropsychopharmacology* **34**, 1590–1600.
- Regenold WT, D'Agostino CA, Ramesh N, Hasnain M, Roys S, Gullapalli RP (2006). Diffusion-weighted magnetic resonance imaging of white matter in bipolar disorder, a pilot study. *Bipolar Disorders* **8**, 188–195.
- Sussmann JE, Lymer GK, McKirdy J, Moorhead TW, Maniega SM, Job D, Hall J, Bastin ME, Johnstone EC, Lawrie SM, McIntosh AM (2009). White matter abnormalities in bipolar disorder and schizophrenia detected using diffusion tensor magnetic resonance imaging. *Bipolar Disorders* **11**, 11–18.
- Tondo L, Baldessarini RJ (2009). Long-term lithium treatment in the prevention of suicidal behavior in bipolar disorder patients. *Epidemiologia e Psichiatria Sociale* **18**, 179–183.

- Versace A, Almeida JR, Hassel S, Walsh ND, Novelli M, Klein CR, Kupfer DJ, Phillips ML** (2008). Elevated left and reduced right orbitomedial prefrontal fractional anisotropy in adults with bipolar disorder revealed by tract-based spatial statistics. *Archives of General Psychiatry* **65**, 1041–1052.
- Vita A, De Peri L, Sacchetti E** (2009). Gray matter, white matter, brain, and intracranial volumes in first-episode bipolar disorder, a meta-analysis of magnetic resonance imaging studies. *Bipolar Disorders* **11**, 807–814.
- Wang F, Jackowski M, Kalmar JH, Chepenik LG, Tie K, Qiu M, Gong G, Pittman BP, Jones MM, Shah MP, Spencer L, Papademetris X, Constable RT, Blumberg HP** (2008*b*). Abnormal anterior cingulum integrity in bipolar disorder determined through diffusion tensor imaging. *British Journal of Psychiatry* **193**, 126–129.
- Wang F, Kalmar JH, Edmiston E, Chepenik LG, Bhagwagar Z, Spencer L, Pittman B, Jackowski M, Papademetris X, Constable RT, Blumberg HP** (2008*a*). Abnormal corpus callosum integrity in bipolar disorder: a diffusion tensor imaging study. *Biological Psychiatry* **64**, 730–733.
- Wessa M, Houenou J, Leboyer M, Chanraud S, Poupon C, Martinot JL, Paillère-Martinot ML** (2009). Microstructural white matter changes in euthymic bipolar patients: a whole-brain diffusion tensor imaging study. *Bipolar Disorders* **11**, 504–514.
- Yurgelun-Todd DA, Silveri MM, Gruber SA, Rohan ML, Pimentel PJ** (2007). White matter abnormalities observed in bipolar disorder, a diffusion tensor imaging study. *Bipolar Disorders* **9**, 504–512.
- Zanetti MV, Jackowski MP, Versace A, Almeida JR, Hassel S, Duran FL, Busatto GF, Kupfer DJ, Phillips ML** (2009). State-dependent microstructural white matter changes in bipolar I depression. *European Archives of Psychiatry and Clinical Neuroscience* **259**, 316–328.
- Zhang Y, Xu H, Jiang W, Xiao L, Yan B, He J, Wang Y, Bi X, Li X, Kong J, Li XM** (2008). Quetiapine alleviates the cuprizone-induced white matter pathology in the brain of C57BL/6 mouse. *Schizophrenia Research* **106**, 182–189.