

**full
analysis
in
3 hours!
with the new...**



AMINO ACID ANALYSERS

Write for illustrated brochure



EVANS ELECTROSELENIUM LIMITED
HALSTEAD · ESSEX · ENGLAND

Telephone : Halstead 2461

Telex : 98204

EEL INTERNATIONAL LIMITED
BAYSWATER VICTORIA AUSTRALIA

The Annals of Applied Biology

The Annals of Applied Biology publishes results of original experimental work on production and preservation of temperate and tropical agricultural and horticultural crops. It covers aspects of plant physiology and genetics, together with the epidemiology and control of damaging insects, nematodes, fungi, bacteria and viruses. The effects of environment, and the interactions between these various factors are included. A special interest of the journal is in the mode of action of fungicides, insecticides, herbicides and growth regulators.

Contents of Volume 64 Number 3 December 1969

- G. H. F. LUPTON Estimation of yield in wheat from measurements of photosynthesis and translocation in the field
- E. C. HUMPHRIES AND W. BOND Experiments with CCC on wheat: effects spacing, nitrogen and irrigation
- E. C. HUMPHRIES AND S. A. W. FRENCH Effect of seedling treatment on growth and yield of sugar beet in the field
- D. B. HARPER AND R. L. WAIN Studies on plant growth-regulating substances. XXX. The plant growth-regulating activity of substituted phenols
- J. F. DUNACHIE AND W. W. FLETCHER An investigation of the toxicity of insecticides to bird's eggs using the egg-injection technique
- F. P. ROWE AND R. REDFERN Aggressive behaviour in related and unrelated wild house mice (*Mus musculus* L.)
- N. E. A. SCOPES The potential of *Chrysopa carnea* as a biological control agent of *Myzus persicae* on glasshouse chrysanthemums
- I. J. WYATT A laboratory method for studying the biology and control of pests in mushroom compost
- F. T. LAST, M. H. EBBEN, R. C. HOARE, E. A. TURNER AND A. R. CARTER Build-up of tomato brown rootrot caused by *Pyrenochaeta lycopersici* Schneider and Gerlach
- J. F. MALCOLMSON Factors involved in resistance to blight (*Phytophthora infestans* (Mont.) de Bary) in potatoes and assessment of resistance using detached leaves
- J. A. C. HARRISON AND I. ISAAC Host/parasite relations up to the time of tuber initiation in potato plants infected with *Verticillium* spp.
- P. W. TALBOYS AND M. BENNETT Growth and wilt (*Verticillium dahliae*) development in strawberry cultivars over a transition between two soil series
- S. W. BURRAGE Dew and the growth of the uredospore germ tube of *Puccinia graminis* on the wheat leaf
- D. HORNBY Methods of investigating populations of the take-all fungus (*Ophiobolus graminis*) in soil
- J. N. GIBBS Inoculum sources for coffee berry disease
- E. GRIFFITHS AND J. N. GIBBS Early-season sprays for the control of coffee berry disease
- Laws of the Association of Applied Biologists

Subscription £5/\$16.50 per volume (2 vols. a year)

Single parts 45s./\$7 plus packing

Further information from

CAMBRIDGE UNIVERSITY PRESS

P.O. BOX 92, LONDON N.W. 1

32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Gardona

the spray-safe insecticide

Gardona is a new insecticide from Shell. And it's a spray-safe insecticide because it can be sprayed without the use of special protective clothing. What's new about Gardona is its low toxicity to man and animals.

Gardona is especially effective against caterpillars, flies and beetles. It protects apples and pears from Codling moth, citrus fruits from fruit flies, and vegetables from caterpillars and flea-beetles.

And Gardona's selective activity against these pests means most beneficial insects and organisms survive the spraying.

Although Gardona will control pests for up to fourteen days, depending on the dosage, its low-mammalian toxicity means crops may be safely harvested a very short time after treatment. In most cases the following day.

Spray-safe Gardona is also showing promising results in the control

of major pests in a variety of other crops such as cotton, rice and maize.

Gardona is another product of Shell's continuing research into the development of effective and safe insecticides. For further information or supplies of Gardona contact your Shell company or Shell Chemicals distributor.

Shell Chemicals



Continued from back page

	PAGE
DRAYCOTT, A. P. and HOLLIDAY, R. Comparisons of liquid and solid fertilizers and anhydrous ammonia for sugar beet. (With 1 text-figure)	139
DRAYCOTT, A. P. and LAST, P. J. Effect of previous cropping and manuring on the nitrogen fertilizer needed by sugar beet. (With 1 text-figure)	147
COLLIER, P. A., HOOD, A. E. M., POWELL, R. A. and TEMPLEMAN, W. G. The agronomic properties of urea-containing compound fertilizers. I. High-nitrogen type. (With 2 text-figures)	153
MANSTON, R. and VAGG, M. J. Urinary phosphate excretion in the dairy cow. (With 3 text-figures)	161
VAN GYLSWYK, N. O. The effect of supplementing a low-protein hay on the cellulolytic bacteria in the rumen of sheep and on the digestibility of cellulose and hemicellulose. (With 4 text-figures)	169
EDEY, T. N. Nutritional stress and pre-implantation embryonic mortality in Merino sheep, 1965. (With 2 text-figures)	181
EDEY, T. N. Nutritional stress and pre-implantation embryonic mortality in Merino sheep, 1966. (With 2 text-figures)	187
EDEY, T. N. Nutritional stress and pre-implantation embryonic mortality in Merino sheep, 1967. (With 2 text-figures)	193
EDEY, T. N. Nutritional stress and pre-implantation mortality in Merino sheep, 1964-7. General discussion and conclusions	199
MORAN, J. B. Seasonal and diurnal variations in body temperatures of Hereford and Brahman cross cattle in a cool temperate environment (With 2 text-figures)	205

THE JOURNAL OF AGRICULTURAL SCIENCE

CONTENTS

Vol. 74 Part 1 February 1970

	PAGE
GOLDSWORTHY, P. R. and TAYLER, R. S. The effect of plant spacing on grain yield of tall and short sorghum in Nigeria	1
MUDD, A. J. The influence of heavily fertilized grass on mineral metabolism of dairy cows. (With 1 text-figure)	11
ALLINSON, D. W. and OSBOURN, D. F. The cellulose-lignin complex in forages and its relationship to forage nutritive value. (With 4 text-figures)	23
BOYD, D. A., TINKER, P. B. H., DRAYCOTT, A. P. and LAST, P. J. Nitrogen requirement of sugar beet grown on mineral soils. (With 2 text-figures)	37
YOELAO, K., JACKSON, M. G. and SARAN, ISHWAR. The effect of wilting berseem and lucerne herbage on voluntary dry-matter intake by buffalo heifers	47
PATTERSON, H. D. and LOWE, BRIDGET, I. The errors of long-term experiments	53
MCEWEN, J. Fertilizer nitrogen and growth regulators for field beans (<i>Vicia faba</i> L.). I. The effects of seed bed applications of large dressings of fertilizer nitrogen and the residual effects on following winter wheat	61
MCEWEN, J. Fertilizer nitrogen and growth regulators for field beans (<i>Vicia faba</i> L.). II. The effects of large dressings of fertilizer nitrogen, single and split applications, and growth regulators	67
TINKER, P. B. Fertilizer requirements of sugar beet on fen peat soils	73
MATTINGLY, G. E. G. Total phosphorus contents of soils by perchloric acid digestion and sodium carbonate fusion. (With 2 text-figures)	79
LYNE, A. G., JOLLY, M. and HOLLIS, D. E. Effects of experimentally produced local subdermal temperature changes on skin temperature and wool growth in the sheep. (With 8 text-figures)	83
GALLAGHER, J. R. and YEATES, N. T. M. An evaluation of Merino wool quality. I. The incidence of coarse fibres in two Australian Merino flocks (With 1 plate and 2 text-figures)	91
GALLAGHER, J. R. An evaluation of Merino wool quality. II. An estimate of the incidence of coarse fibres in Australian Merino wool. (With 2 text-figures)	99
HALLIDAY, R. Protein concentrations in colostrum from Finnish Landrace × Scottish Blackface ewes during the first week of lactation and in sera from the ewes and their lambs on the third day of lactation. (With 1 text-figure)	103
GASSER, J. K. R. The decomposition of crotonylidene di-urea in soil and the recovery of nitrogen by ryegrass. (With 2 text-figures)	107
SPRATT, E. D. and GASSER, J. K. R. The effect of ammonium sulphate with a nitrification inhibitor, and calcium nitrate, on growth and N-uptake of spring wheat, ryegrass and kale. (With 3 text-figures)	111
ADDISCOTT, T. M. Use of the quantity/potential relationship to provide a scale of the ability of extractants to remove soil potassium. (With 1 text-figure)	119
ADDISCOTT, T. M. The uptake of initially available soil potassium by ryegrass. (With 5 text-figures)	123
ADDISCOTT, T. M. The potassium Q/I relationships of soils given different K manuring. (With 4 text-figures)	131

Continued on inside back cover

SUBSCRIPTIONS. Two volumes of three parts are published annually. The subscription price is £6 net (U.S.A. \$19.50) per volume (post free); single parts are available at £2 14s. net (U.S.A. \$8.00) plus postage. Orders or enquiries may be sent to any bookseller or subscription agent, or to Cambridge University Press, P.O. Box 92, London, N.W. 1, (U.S.A. and Canada, Cambridge University Press American Branch, 32 East 57th Street, New York, N.Y. 10022, U.S.A.)

Printed in Great Britain at the University Printing House, Cambridge