

CORRESPONDENCE

THE LOWEST TEMPERATURES RECORDED IN
ANTARCTICA AND GREENLAND

May I point out with reference to the note on the coldest place on earth, published in the July 1953 number of the *Polar Record* (p. 821-22), that the lowest temperature measured in Antarctica is not that recorded during the winter journey in July 1911, -60.5°C . (-77°F .) but the -61°C . (-78°F .) and possibly -64°C . (-83°F .) measured by R. E. Byrd at Bolling Base (lat. $80^{\circ} 8' \text{S}$., long. $163^{\circ} 55' \text{W}$.) on 21 and 22 July 1934.^{1,2} The mean for 21 July was -60.6°C . (-77.1°F .)¹ A temperature of -60.5°C . (-77°F .) was reached on 20, 21 and 30 August 1934.²

It may interest readers to learn that the lowest temperature recorded on the Greenland ice sheet was -66°C . (-87°F .), on 21 February 1950 at the French central station in lat. $70^{\circ} 54' \text{N}$., long. $40^{\circ} 42' \text{W}$., 2993 m. (9820 ft.).³ On this day the temperature in the free atmosphere did not rise above -48°C . (-54°F .) at any level. This is probably the coldest troposphere temperature ever recorded. The coldest day, 6 December 1949, had a mean of -63.3°C . (-82°F .)

DEPARTMENT OF METEOROLOGY
UNIVERSITY OF MELBOURNE
5 September 1953

F. LOEWE

References

¹ G. GRIMMINGER and W. C. HAINES. Meteorological results of the Byrd antarctic expeditions 1928-30, 1933-35: tables. *United States. Weather Bureau, Monthly Weather Review, Supplement No. 41*, 1939, p. 302-03.

² G. GRIMMINGER. Meteorological results of the Byrd antarctic expeditions 1928-30, 1933-35: summaries of data. *United States. Weather Bureau, Monthly Weather Review, Supplement No. 42*, 1941, p. 72.

³ Recueil des observations météorologiques effectuées par la station française du Groënland entre le 20 Juillet 1949 et le 14 Juillet 1950. *Expéditions Polaires Françaises [Publications Préliminaires No. 12]*, 1950, p. 30, 50.

"ICE STREAMS"

I welcome the opportunity to comment upon C. W. M. Swithinbank's note on "ice streams."* This term aptly describes many areas in Antarctica where glacier ice progresses to lower altitudes at a relatively greater speed than in adjacent parts of the continental glacier. The differential rate of movement causes severe crevasses to occur on the surface of these "ice streams", but in some instances pressure or drifting snow tends to conceal them. Although governed by different forces, such "ice streams" are in some ways comparable to certain ocean currents, such as the Gulf Stream, which move more rapidly than adjacent waters of the surrounding ocean.

I have used the term "channel glacier" for this phenomenon, not only because these features act as channels for massive movements of glacier ice

* See p. 185-86.