

**Aims.** Rampton Hospital is the High Secure Hospital of Nottinghamshire Healthcare NHS Foundation Trust's Forensic Service. It is one of three such hospitals in England, following Security Directions set out by the Department of Health. Patient management occurs through the implementation of strict policies and procedures. Policy requirements highlight the need for MDT post-incident discussion of restrictive interventions, and in particular, of Rapid Tranquilisation (RT). This primary audit aimed to ascertain current practice and if necessary, suggest interventions to ensure that patient-care remains safe, effective, and well-led.

**Methods.** To establish current practice with regards to the discussion of individual cases of RT in MDT settings, specifically in Ward Round, we commenced a retrospective data collection from electronic notes covering all directorates within the High Secure estate between May and June 2022.

From these notes, we tried to ascertain whether the following policy standards were being met:

- A de-brief with the patient should take place as soon after the incident as is practicable and reasonable, ideally within 72 hours.
- The MDT meeting post RT episode should explicitly discuss the episode, and consider medication and any triggers of periods of acutely disturbed behaviour.
- There were 81 data sets to explore.

**Results.** Not all data sets were viable. Out of those analysed, less than 10% were found to have met the aforementioned ideal policy standards of having had a reflective discussion within 72 hours with both the patient and as an MDT, exploring the episode itself and its antecedents.

**Conclusion.** There are several interesting factors to consider from the results obtained. We postulate that the frequency of episodes of RT makes meeting the policy standard problematic; pragmatically, there is a significant time barrier to exploring these incidents in detail and the various teams, operating in dynamic and high-risk environments, may find it difficult to coalesce in order to debrief appropriately.

Furthermore, the reflections may actually be happening, but the burden of documentation mean that these are not being recorded formally in a way that can be measured.

There are limitations to the searches of electronic notes and we did not have access to Incident Reports, often completed at the time of these episodes; further information may have been uncovered if they were available.

Despite this, there is room for interventions that inform staff of this need and to provoke improvements in current practice.

---

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Seclusion on Psychiatric Intensive Care Unit – Is the Trust Medical Review Policy Being Followed?

Dr Hannah Scanlon\*

South West London and St George's Mental Health NHS Trust, London, United Kingdom

\*Corresponding author.

doi: 10.1192/bjo.2023.470

**Aims.** Seclusion is a restrictive intervention used in inpatient settings for the safe management of patients who present with severe behavioural disturbance which is likely to cause harm to others.

Clinical notes were used to establish if the trust policy of medical reviews for patients in seclusion was being followed on the Psychiatric Intensive Care Unit (PICU).

**Methods.** Patients admitted to male PICU at Springfield Hospital, Southwest London, over a 4 month period (February 2022 to May 2022) were included in this audit. Patients who were secluded outside this time period or prior to admission to PICU were excluded from this audit.

The clinical notes computer system (Rio) was searched using the term "seclusion". The timing of initiation and termination of seclusion were noted as well as the timing and grade of medical professionals present for documented reviews.

**Results.** Over this period, 12 discrete episodes of seclusion were identified. The length of seclusion varied from 8 hours 45 minutes to over 5 days, with a mean length of almost 3 days (2 days, 20 hours, 25 minutes).

As the length of seclusion differed so did the required medical reviews in line with trust policy. This involves Senior House Officer (SHO) review at 30 minutes, Registrar review at 8 hours, Consultant review at 24 hours followed by 2 senior reviews (one Registrar and one Consultant) over each subsequent 24 hour period of continuous seclusion.

10 episodes of seclusion lasted over 24 hours in this audit. Of these 40% had the required medical reviews documented in the clinical notes appropriately for the full period of seclusion. 50% of cases had at least 1 missed or not documented Registrar review. There were 2 incidents of missed Consultant medical reviews for a 24 hour period of continuous seclusion.

**Conclusion.** From these results medical reviews were not being correctly carried out, or were not documented correctly, in the majority (60%) of cases of seclusion over 24 hours. This suggests missed opportunities for patient review to terminate seclusion at the earliest safe opportunity in line with national and trust guidance. These results have informed the update of trust guidelines on seclusion to bring it in line with national guidance with a view to improve patient care and will be re-audited.

---

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Clinical Audit Reviewing Compliance With Respective Trust-Based Physical Health Monitoring Guidelines, Amongst Inpatients Prescribed Anti-Psychotic Medication in Two Distinct Secure Care Facilities: A Low Secure Unit and a Prison Personality Disorder Unit

Dr Ashvini Selvaraj\* and Dr Falade Adebowale

Fulbourn Hospital, Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, United Kingdom

\*Corresponding author.

doi: 10.1192/bjo.2023.471

**Aims.**

1. To review the current level of compliance with CPFT (Cambridge & Peterborough Foundation Trust) guidelines by inpatients prescribed anti-psychotics at George Mackenzie House (GMH) low-secure unit and likewise, with NHFT (Northamptonshire Foundation Trust) guidelines by inmates prescribed anti-psychotics at HMP Whitemoor's Fens Unit.
2. To identify any differences or similarities in compliance rates between both sites.