Bull. Austral. Math. Soc. Vol. 46 (1992) [177-178]

## ISOMETRIC IMMERSION OF A COMPACT RIEMANNIAN MANIFOLD INTO A EUCLIDEAN SPACE

## SHARIEF DESHMUKH

We show that an isometric immersion of an n-dimensional compact Riemannian manifold of non-negative Ricci curvature with scalar curvature always less than  $n(n-1)\lambda^{-2}$  into a Euclidean space of dimension n+1 can never be contained in a ball of radius  $\lambda$ .

Jacobowitz [2] proved that an isometric immersion of an n-dimensional compact Riemannian manifold with sectional curvature always less than some constant  $\lambda^{-2}$  into Euclidean space of dimension 2n-1 can never be contained in a ball of radius  $\lambda$ . This result generalised the results of Tompkins [4], Chern and Kuiper [1] and Otsuki [3]. However it is not known whether such a nonimmersibility theorem holds with the condition on sectional curvature replaced by a suitable condition on Ricci curvature or the scalar curvature. In this note we prove the following co-dimension one nonimmersibility theorem.

**THEOREM.** Let E be Euclidean space of dimension n+1 and M be a compact n-dimensional Riemannian manifold of non-negative Ricci curvature whose scalar curvature is less than some constant  $n(n-1)\lambda^{-2}$ . Then no isometric immersion of M into E is contained in a ball of radius  $\lambda$ .

PROOF: Assume that  $\psi \colon M \to E$  is the isometric immersion such that  $\psi(M)$  is contained in a ball of radius  $\lambda$ . Thus we can assume that  $\|\psi\| \leqslant \lambda$ , where  $\|\cdot\|$  is the norm on E with respect to the Euclidean metric  $\langle \cdot, \cdot \rangle$ . Let N be the unit normal vector field to M. Then the support function  $\rho \colon M \to R$  is defined by  $\rho = \langle \psi, N \rangle$ , and we have the Minkowski's formula

(1) 
$$\int_{M} (1 + \rho \alpha) dv = 0,$$

where  $\alpha$  is the mean curvature of M.

Denote by g,  $\nabla$  and A the Riemannian metric, the Riemannian connection and the Weingarten map on M respectively. The position vector field  $\psi$  can be expressed

Received 16 August 1991

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/92 \$A2.00+0.00.

as  $\psi = t + \rho N$ , where t is a vector field on M. Then using Gauss and Weingarten formulae we immediately get

$$\nabla_X t = X + \rho A X,$$

for any vector field X on M.

If  $\eta$  is a 1-form dual to t, then using equation (2), we obtain

(3) 
$$d\eta = 0, \qquad \delta \eta = n(1 + \rho \alpha).$$

Also, we obtain

$$\|\nabla t\|^2 = n + 2\rho n\alpha + \rho^2 \operatorname{tr} A^2.$$

Now using  $||t||^2 + \rho^2 = ||\psi||^2$  and the expression for the scalar curvature S,  $S = n^2 \alpha^2 - \text{tr } A^2$ , we get

(4) 
$$\|\nabla t\|^2 - (\delta \eta)^2 = \|t\|^2 S + n(n-1) - \|\psi\|^2 S - 2n(n-1)(1+\rho\alpha).$$

For a compact Riemannian manifold the following integral formula is known (see [5], p.41)

$$\int_{M} \left\{ \operatorname{Ric}(t, t) - \frac{1}{2} \left\| d\eta \right\|^{2} + \left\| \nabla t \right\|^{2} - \left( \delta \eta \right)^{2} \right\} dv = 0,$$

where Ric is the Ricci tensor of M.

Using (1), (3) and (4) in the above integral formula we get

(5) 
$$\int_{M} \left\{ \operatorname{Ric}(t, t) + \|t\|^{2} S + \left( n(n-1) - \|\psi\|^{2} S \right) \right\} dv = 0.$$

From the hypothesis of the Theorem it follows that  $\text{Ric}(t, t) \ge 0$ ,  $||t||^2 S \ge 0$  and  $||\psi||^2 S \le \lambda^2 S < n(n-1)$  which contradicts (5). This proves the Theorem.

## REFERENCES

- [1] S.S. Chern and N.H. Kuiper, 'Some theorems on the isometric embedding of compact Riemannian manifolds in Euclidean space', *Ann. of Math.* 56 (1952), 442-430.
- [2] H. Jacobowitz, 'Isometric embedding of a compact Riemannian manifold into Euclidean space', Proc. Amer. Math. Soc. 40 (1973), 245-246.
- [3] T. Otsuki, 'On the existence of solutions of a system of quadratic equations and its geometrical application', *Proc. Japan. Acad.* 29 (1953), 99-100.
- [4] C. Tompkins, 'Isometric embedding of flat manifolds in Euclidean space', Duke Math. J. 5 (1939), 58-61.
- [5] K. Yano, Integral formulas in Riemannian geometry (Marcel Dekker, New York, 1970).

Department of Mathematics King Saud University PO Box 2455 Riyadh 11451 Saudi Arabia