



Received 9 February 1980
Final 15 September 1980

Analysis of Multiple Birth Rates in Japan. II. Secular Trend and Effect of Birth Order, Maternal Age, and Gestational Age in Stillbirth Rate of Twins

Yoko Imaizumi,¹ Akio Asaka,² Eiji Inouye²

¹ *Institute of Population Problems, Ministry of Health and Welfare, Tokyo;* ² *Institute of Brain Research, University of Tokyo School of Medicine*

Stillbirth rates of MZ and DZ twins in Japan gradually decreased during the period 1960–1967 and in 1974. The stillbirth rates of MZ and DZ twins were 0.270 and 0.224 in 1960, respectively, whereas the corresponding figures in 1974 were 0.135 and 0.099. The stillbirth rate was higher in MZ than DZ twins, in males than females, and in the second- than in the first-born. In both zygositys, the rates were higher in the first birth order than in the second, and then increased with birth order except in the sixth birth or more for MZ twins and for DZ twins born to mothers aged 35–39 years. The shorter the gestational age, the higher the stillbirth rate in both MZ and DZ twins, as expected. Mean gestational age was slightly shorter in MZ than in DZ twins. The decrease of the stillbirth rates can be explained partly by the reduction of twins with higher birth order, where high stillbirth rates of MZ and DZ twins are seen, and the reduction of older mothers.

Key words: Stillbirth rates of twins, Birth order, Maternal age, Gestational age, Secular trends, Japan

INTRODUCTION

Among previous studies on the stillbirth rates of twins, Eriksson et al [2] reported that the rate in Sweden during the period from 1901 to 1960 remained constant up to 1940 and decreased gradually thereafter. Yerushalmy and Sherar [11] reported stillbirth rates among monozygotic (MZ) and dizygotic (DZ) twins in New York during the period 1936–1937. They found that the second of a pair of twins is exposed to a greater risk of a stillbirth than the first twin. The stillbirth rate of twins is more than three times as high as those of single births, and it was higher in MZ than in DZ twins. Barr and Stevenson [1] reported the stillbirth rate among total deliveries separately for MZ and DZ twins in England and Wales during the period 1949–1950. It was higher in MZ than in DZ twins in each sex, and was higher in male twins than in female twins in this population.

Imaizumi and Inouye [4] reported in the first report of the present series that stillbirth rate in Japan is high in young and old maternal age groups in both MZ and DZ twins. The present study deals with secular change of stillbirth rate of twins. It also deals with the effects of birth order, maternal age and gestational age on stillbirth rate of twins.

MATERIALS AND METHODS

There were two sources of data available for the present analysis: The annual volumes of Vital Statistics of Japan for the years 1951–1968 and 1974 [5] and the “Survey on Socio-Economic Aspects of Vital Events—Plural Births in 1975” [6]. In the first data, sex of the children and maternal age are listed from 1960 to 1967 and in 1974. Second data were derived from two surveys, A and B. Survey A was made on birth or stillbirth certificate records in 1974 on 25,189 children or fetuses born in plural deliveries to mothers of Japanese nationality, where the fetal death is defined as that which occurred after the beginning of 13th week of gestation. The survey B was conducted on a part of children included in survey A, and was not available in the present analysis. Details of the surveys have been reported elsewhere [4]. Items used in the present analysis are sex of children, birth order of twins, live birth or fetal death, maternal age, gestational age, and the numbers of previous pregnancies and live births.

RESULTS

Secular Change of Stillbirth Rate of Twins

Table 1 shows the number of twin deliveries according to survival states of twins during the period from 1951 to 1968 and in 1974. In the following analysis, the stillbirth rate of twins was taken among twin deliveries, where total number of stillborn twin individuals was taken as numerator. The stillbirth rate of twins thus computed is also shown in Table 1. The rate increased significantly with the years up to 1961, and decreased thereafter.

Table 2 shows secular trends of the stillbirth rates of MZ and DZ twins during the period from 1960 to 1967 and in 1974. In computing the stillbirth rate separately for MZ and DZ twins, the numbers of twin individuals (live birth or stillbirth) were obtained after the numbers of MZ and DZ twins were estimated by Weinberg's method for each category of survival state of twins, two live-born, one live-born and one stillborn, and two stillborn.

In 1960, the stillbirth rates of like- and unlike-sexed twins were 0.260 and 0.224, respectively, the difference being significant ($P < 0.001$). Corresponding stillbirth rates of MZ and DZ twins were 0.270 and 0.224 in 1960, respectively. The rates gradually decreased with the years both in MZ and DZ twins, and the figures in 1974 are less than one half of those in 1960. Figure 1 shows secular trends of the stillbirth rates of MZ, DZ, and total twins including unknown sex. Similar decrease of stillbirth rates in the general population is seen from 1961 (Fig. 1), but the stillbirth rate of twins remained more than 2–3 times as high as that of the general population.

Table 3 shows secular trends of the stillbirth rates of twins according to sex during the period from 1960 to 1967 and in 1974. In both sexes the stillbirth rates are gradually decreased with the years in parallel to the decrease of stillbirth rates of MZ and DZ twins. In every year, sex difference of the stillbirth rates is statistically significant.

Stillbirth rates of twins in 1974 were analyzed according to zygosity, sex, and birth order of twins (Table 4). The stillbirth rate was higher in males than females in MZ twins, but this trend was not remarkable in DZ twins. The stillbirth rate was higher in MZ than DZ twins for both sexes. The stillbirth rate was lower for the first twin (0.112) than for the second (0.127), the difference being significant ($P < 0.001$).

Effect of Birth Order, Maternal Age, and Gestational Age on Stillbirth Rate of Twins

In the following analysis, the birth order concerns the numerical order of all children and fetuses (live-born and prenatally dead in the 21st week of gestation or later), where a delivery of twins was counted once. Table 5 shows the number of twin deliveries in 1974 according to survival state, sex, and the birth order, from which was computed the stillbirth rate of MZ and DZ twins according to birth order. Table 6 shows the result, in which the number of fetal deaths and stillbirth rates are shown according to maternal age. In MZ twins the stillbirth rate decreased from the first to the second birth, then increased but again decreased in higher birth orders in most maternal age groups. In DZ twins, the same pattern is seen in the maternal age group of 35–39 years, and in maternal age groups of 25–29 and 30–34 years are seen a decrease from the first to the second and an increase to the fourth.

As to the effect of maternal age in MZ twins, the lowest stillbirth rate is seen in the maternal age group of 25–29 years in the first birth order. As the birth order increases, the lowest rate shifts to the higher maternal age groups. Similar but less clear tendency is also seen in DZ twins.

Table 7 shows the number and frequency of deliveries of twins by zygosity according to survival state and gestational age. Gestational age of MZ twins tended to be shorter than DZ twins in the total of all survival states, which reflects the shorter gestational age of MZ twins than DZ twins in two live-born and one live-born categories. In the two-stillborn category, the frequency is higher in MZ twins than DZ twins in most gestational age groups.

TABLE 1. Number of Twin Deliveries According to Survival States of Twins and Stillbirth Rates Among Twins, 1951–1968 and 1974

| Year | 2 Live births | 1 Live birth, 1 fetal death | 2 Fetal deaths | Total | Stillbirth rate per 100 twin individuals |
|------|---------------|--------------------------------|----------------|--------|---|
| 1951 | 10,600 | 1,888 | 2,655 | 15,143 | 23.77 |
| 1952 | 9,822 | 1,682 | 2,503 | 14,007 | 23.87 |
| 1953 | 8,952 | 1,621 | 2,480 | 13,053 | 25.21 |
| 1954 | 8,727 | 1,415 | 2,513 | 12,655 | 25.45 |
| 1955 | 8,221 | 1,447 | 2,374 | 12,042 | 25.72 |
| 1956 | 8,023 | 1,364 | 2,338 | 11,725 | 25.76 |
| 1957 | 7,762 | 1,331 | 2,314 | 11,407 | 26.12 |
| 1958 | 8,027 | 1,332 | 2,458 | 11,817 | 26.44 |
| 1959 | 7,907 | 1,280 | 2,392 | 11,579 | 26.19 |
| 1960 | 7,701 | 1,149 | 2,309 | 11,159 | 25.84 |
| 1961 | 7,855 | 1,178 | 2,361 | 11,394 | 25.89 |
| 1962 | 8,110 | 1,043 | 2,301 | 11,454 | 24.64 |
| 1963 | 8,304 | 979 | 2,355 | 11,638 | 24.44 |
| 1964 | 8,993 | 1,035 | 2,140 | 12,168 | 21.84 |
| 1965 | 9,338 | 901 | 2,027 | 12,266 | 20.20 |
| 1966 | 7,333 | 691 | 1,824 | 9,848 | 22.03 |
| 1967 | 10,483 | 844 | 1,885 | 13,212 | 17.46 |
| 1968 | 9,931 | 660 | 1,756 | 12,347 | 16.89 |
| . | | | | | |
| . | | | | | |
| . | | | | | |
| 1974 | 10,418 | 645 | 1,187 | 12,250 | 12.32 |

TABLE 2. MZ and DZ Twin Deliveries According to Survival States of Twins and Stillbirth Rates Among Twins, 1960-1967 and 1974

| Year | MZ twin deliveries | | | | DZ twin deliveries | | | | Stillbirth rate per 100 twin individuals |
|------|--------------------|------------|-------|-------|--------------------|------------|------|-------|--|
| | 2 LB | 1 LB, 1 FD | 2 FD | Total | 2 LB | 1 LB, 1 FD | 2 FD | Total | |
| 1960 | 4,865 | 723 | 1,576 | 7,164 | 2,836 | 416 | 670 | 3,922 | 22.39 |
| 1961 | 4,871 | 720 | 1,649 | 7,240 | 2,984 | 448 | 628 | 4,060 | 20.99 |
| 1962 | 5,154 | 677 | 1,673 | 7,504 | 2,956 | 354 | 558 | 3,868 | 19.00 |
| 1963 | 5,288 | 661 | 1,694 | 7,643 | 3,016 | 312 | 580 | 3,908 | 18.83 |
| 1964 | 5,639 | 676 | 1,547 | 7,862 | 3,354 | 352 | 514 | 4,220 | 16.35 |
| 1965 | 5,868 | 618 | 1,469 | 7,955 | 3,470 | 272 | 470 | 4,212 | 14.39 |
| 1966 | 4,803 | 484 | 1,351 | 6,638 | 2,530 | 196 | 392 | 3,118 | 15.72 |
| 1967 | 6,893 | 560 | 1,408 | 8,861 | 3,590 | 268 | 398 | 4,256 | 12.50 |
| 1974 | 6,954 | 459 | 889 | 8,302 | 3,464 | 186 | 298 | 3,948 | 9.90 |

LB = live birth; FD = fetal death.

TABLE 3. Stillbirth Rates of Twins for Males and Females, 1960–1967 and 1974

| Year | Males | Females |
|------|--------|---------|
| 1960 | 0.2777 | 0.2291 |
| 1961 | 0.2755 | 0.2297 |
| 1962 | 0.2643 | 0.2173 |
| 1963 | 0.2609 | 0.2155 |
| 1964 | 0.2299 | 0.1951 |
| 1965 | 0.2137 | 0.1777 |
| 1966 | 0.2383 | 0.1874 |
| 1967 | 0.1859 | 0.1518 |
| . | | |
| . | | |
| 1974 | 0.1349 | 0.1133 |

TABLE 4. Stillbirth Rates of Twins According to Sex, Zygosity, and Birth Order, 1974

| | Male | | | Female | | | Total | |
|-----------------|--------|--------|--------|--------|--------|--------|----------|----------|
| | MZ | DZ | Total | MZ | DZ | Total | 1st-born | 2nd-born |
| Stillbirth rate | 0.1494 | 0.1029 | 0.1349 | 0.1194 | 0.0953 | 0.1133 | 0.1118 | 0.1266 |

TABLE 5. Survival States and Sex of Twins According to Birth Order, 1974

| Survival states of twins | Sex of twins | Birth order | | | | | | Total |
|-----------------------------|--------------|-------------|-------|-----|-----|----|----|-------|
| | | 1 | 2 | 3 | 4 | 5 | 6+ | |
| 2 live births | MM | 1,887 | 1,565 | 571 | 101 | 17 | 17 | 4,158 |
| | FF | 1,970 | 1,518 | 563 | 96 | 21 | 11 | 4,179 |
| | MF | 735 | 628 | 232 | 33 | 13 | 8 | 1,649 |
| 1 live birth, 1 fetal death | MM | 123 | 122 | 43 | 11 | 0 | 1 | 300 |
| | FF | 112 | 84 | 40 | 12 | 1 | 1 | 250 |
| | MF | 44 | 32 | 11 | 6 | 0 | 0 | 93 |
| | Unknown | 9 | 6 | 2 | 0 | 0 | 1 | 18 |
| 2 fetal deaths | MM | 249 | 105 | 135 | 48 | 25 | 6 | 568 |
| | FF | 182 | 104 | 84 | 35 | 15 | 12 | 432 |
| | MF | 56 | 27 | 38 | 13 | 5 | 5 | 144 |
| | Unknown | 52 | 27 | 22 | 8 | 6 | 4 | 119 |

Table 7 also shows mean gestational ages of MZ and DZ twins according to survival state. It is seen that the means are lower in MZ twins than DZ twins in two or one live-born categories as well as in the total of all survival states.

DISCUSSION

The present analysis indicated that stillbirth rate of twins in Japan increased up to 1961, and decreased thereafter. A similar pattern was seen in the stillbirth rate of the general population (Fig. 1). In every year analyzed, the stillbirth rate was higher in MZ than DZ

TABLE 6. Stillbirth Rates of MZ and DZ Twins According to Maternal Age and Birth Order, 1974

| Birth order | Maternal age | | | | | | | | | | | |
|-------------|--------------|--------|--------|--------|---------|--------|----------|--------|--------|--------|---------|--------|
| | MZ | | | | | | DZ | | | | | |
| | Under 25 | 25-29 | 30-34 | 35-39 | Over 40 | Total | Under 25 | 25-29 | 30-34 | 35-39 | Over 40 | Total |
| 1 | 0.1347 | 0.1174 | 0.1343 | 0.2065 | 0.2273 | 0.1276 | 0.0944 | 0.0797 | 0.1264 | 0.1458 | 0.5000 | 0.0934 |
| 2 | 0.1255 | 0.0893 | 0.0850 | 0.0882 | 0.3333 | 0.0957 | 0.0958 | 0.0479 | 0.0710 | 0.0625 | 0.2500 | 0.0626 |
| 3 | 0.4491 | 0.1955 | 0.1172 | 0.1989 | 0.3333 | 0.1879 | 0.2308 | 0.1652 | 0.1208 | 0.1724 | 0.5000 | 0.1548 |
| 4 | 0.8333 | 0.4143 | 0.2325 | 0.1625 | 0.1667 | 0.3127 | 0 | 0.2500 | 0.3333 | 0.4545 | 0 | 0.3077 |
| 5 | 0.5000 | 0.7778 | 0.6111 | 0.4667 | 0.5000 | 0.5820 | - | 0.2000 | 0 | 0.5000 | 1.0000 | 0.2778 |
| 6+ | - | 0.6250 | 0.2857 | 0.2813 | 0.6250 | 0.4000 | 0 | 1.0000 | 0.7500 | 0.3333 | 0 | 0.3846 |

TABLE 7. Frequency of Twin Deliveries of MZ and DZ Twins According to Survival State and Gestational Age

| | Survival state | Gestational age (weeks) | | | | | | | | | | Mean gestational age (weeks) |
|----------|-----------------|-------------------------|--------|--------|--------|--------|---------|--------|------|--|--|------------------------------|
| | | Gestational age (weeks) | | | | | | | | | | |
| | | 12-23 | 24-27 | 28-31 | 32-35 | 36-39 | Over 40 | Total | | | | |
| MZ twins | 2 live births | 0 | 0.0037 | 0.0264 | 0.1476 | 0.6513 | 0.0069 | 0.8359 | 37.0 | | | |
| | 1 live birth, | 0 | 0.0017 | 0.0071 | 0.0130 | 0.0346 | 0.0006 | 0.0570 | 35.8 | | | |
| | 1 fetal death | 0.0519 | 0.0276 | 0.0139 | 0.0056 | 0.0080 | 0 | 0.1070 | 25.0 | | | |
| DZ twins | 2 fetal deaths | 0.0519 | 0.0330 | 0.0474 | 0.1662 | 0.6939 | 0.0075 | 0.9999 | 35.7 | | | |
| | Total | 1.0000 | 0.8640 | 0.3539 | 0.0726 | 0.0363 | 0.0417 | 0.1355 | | | | |
| | Stillbirth rate | 0 | 0.0053 | 0.0265 | 0.1151 | 0.7216 | 0.0058 | 0.8743 | 37.2 | | | |
| DZ twins | 2 live births | 0.0005 | 0.0016 | 0.0058 | 0.0058 | 0.0355 | 0 | 0.0492 | 36.0 | | | |
| | 1 live birth, | 0.0387 | 0.0260 | 0.0080 | 0.0021 | 0.0016 | 0 | 0.0764 | 23.9 | | | |
| | 1 fetal death | 0.0392 | 0.0329 | 0.0403 | 0.1230 | 0.7587 | 0.0058 | 0.9999 | 36.1 | | | |
| DZ twins | 2 fetal deaths | 0.9934 | 0.8203 | 0.2688 | 0.0396 | 0.0243 | 0 | 0.1010 | | | | |
| | Total | 0.9934 | 0.8203 | 0.2688 | 0.0396 | 0.0243 | 0 | 0.1010 | | | | |




Fig. 1. Secular change of stillbirth rates of MZ twins, DZ twins, total twins, and the general population (1951–1968 and 1974).

twins and higher in males than females, except in 1974 where sex difference in DZ twins was not seen. A higher stillbirth rate of twins was seen in the second-born than in the first-born (Table 4). According to Yerushalmy and Sheerar [11], the stillbirth rate of MZ twins decreased, but that of DZ twins increased with birth order. In the present study, ignoring the classes with small numbers of subjects, the stillbirth rates of MZ twins are higher in the first birth order than in the second; and the rates increased with birth order up to the fourth birth order for maternal age groups under 34 years (Table 6). The same pattern is seen in DZ twins for maternal age groups between 25 and 34 years. As to the effect of maternal age, Yerushalmy and Sheerar reported that the stillbirth rates of MZ and DZ twins increased in old maternal age groups, but in the present study, a higher stillbirth rate of twins was seen in young and old maternal age groups both in MZ and DZ twins, as was reported by Imaizumi and Inouye [4] and is seen in Table 8.

According to Karn [7–9], mean gestational age was slightly shorter in same-sex twin pairs than in opposite-sex twin pairs, but Fraccaro [3] obtained the opposite result. In the present study, the mean gestational age was slightly shorter in MZ than in DZ twins (Table 7).

The decline of the stillbirth rate of twins in Japan seems to be partly due to a decline of children with higher birth order and a decline of maternal age, as indicated below. Stillbirth rates of MZ and DZ twins in 1974 are less than a half of those in 1960 (Table 2). Assuming the stillbirth rates of the fourth birth or more in 1974 (0.369 for MZ twins, 0.313 for DZ twins from Table 6) equal to a half of those in 1960, the stillbirth rates of MZ and DZ twins in the fourth birth or more are expected to decrease by 0.06 in MZ twins and 0.05 in DZ twins, since the proportion of the fourth birth or more was 9.8% in 1960 and 3.1% in the general population in 1974 (from vital statistics in 1960 and 1974). Similarly, the

TABLE 8. Stillbirth Rates in MZ and DZ Twins According to Maternal Age, 1960–1967 and 1974

| Year | Maternal age | | | | | | Total | |
|------|--------------|--------|--------|--------|--------|--------|--------|--------|
| | –19 | 24–24 | 25–29 | 30–34 | 35–39 | 40+ | | |
| MZ | 1960 | 0.5079 | 0.2984 | 0.2420 | 0.2599 | 0.2979 | 0.4235 | 0.2704 |
| | 1961 | 0.4252 | 0.2879 | 0.2556 | 0.2706 | 0.3622 | 0.4063 | 0.2775 |
| | 1962 | 0.4565 | 0.2813 | 0.2337 | 0.2759 | 0.3743 | 0.5300 | 0.2681 |
| | 1963 | 0.5042 | 0.2792 | 0.2261 | 0.2636 | 0.4388 | 0.5723 | 0.2649 |
| | 1964 | 0.4350 | 0.2486 | 0.2243 | 0.2259 | 0.3186 | 0.4127 | 0.2398 |
| | 1965 | 0.4286 | 0.2360 | 0.1953 | 0.2271 | 0.2952 | 0.5077 | 0.2235 |
| | 1966 | 0.4145 | 0.2462 | 0.2096 | 0.2442 | 0.3159 | 0.5183 | 0.2400 |
| | 1967 | 0.4201 | 0.1979 | 0.1678 | 0.1855 | 0.2640 | 0.4744 | 0.1905 |
| 1974 | 0.3247 | 0.1497 | 0.1216 | 0.1262 | 0.1738 | 0.2925 | 0.1347 | |
| DZ | 1960 | 0.4333 | 0.2282 | 0.1921 | 0.2361 | 0.2527 | 0.3108 | 0.2239 |
| | 1961 | 0.4643 | 0.2215 | 0.1989 | 0.2024 | 0.2197 | 0.3600 | 0.2099 |
| | 1962 | 0.3529 | 0.2138 | 0.1562 | 0.1984 | 0.2766 | 0.3333 | 0.1900 |
| | 1963 | 0.3889 | 0.2029 | 0.1658 | 0.1887 | 0.2197 | 0.4750 | 0.1883 |
| | 1964 | 0.5000 | 0.1571 | 0.1508 | 0.1502 | 0.2464 | 0.3800 | 0.1635 |
| | 1965 | 0.3810 | 0.1401 | 0.1184 | 0.1602 | 0.2028 | 0.3611 | 0.1439 |
| | 1966 | 0.1250 | 0.1984 | 0.1257 | 0.1462 | 0.2402 | 0.3571 | 0.1572 |
| | 1967 | 0.1667 | 0.1457 | 0.1091 | 0.1187 | 0.1791 | 0.2222 | 0.1250 |
| 1974 | 0.3182 | 0.0943 | 0.0809 | 0.1122 | 0.1792 | 0.3214 | 0.0990 | |

stillbirth rates of male and female twins in 1960 were 0.278 and 0.229, respectively (Table 3), whereas the respective rates in 1974 were 0.135 and 0.113, a decrease to a half during the past 14 years. Assuming the stillbirth rates of the fourth birth or more in 1974 (0.366 for males and 0.334 for females from Table 5) equal to a half of those in 1960, the stillbirth rates for males and females are expected to decrease by 0.06 in male twins and 0.05 in female twins in the fourth birth or more. As to the effect of maternal age, it is seen in Table 8 that the stillbirth rates for MZ and DZ twins decreased with the years in mothers under the age of 35, whereas in mothers over 40 the decline of the rates was least marked. In maternal age groups of 20–34 years, the stillbirth rates of MZ and DZ twins in 1960 are approximately two times as high as those in 1974, but the rates in maternal age groups less than 20 years and 35 years or more in 1960 are between 1.44 and 1.71 times (MZ twins) and between 0.97 and 1.41 times (DZ twins) as high as those in 1974. The proportion of mothers over 35 years of age was 6.7% in 1960 and 4.4% in the general population in 1974 (from vital statistics in 1960 and 1974). A diminished proportion of old mothers over 35 years of age during 14 years was only 2.3%, and the stillbirth rates in the same maternal age group in 1960 are more than 1.69 (0.321/0.190) times for MZ twins and 1.56 (0.305/0.196) times for DZ twins as high as those in 1974 from Table 8. Therefore, expected decreases of the stillbirth rates are only 0.013 for MZ twins and 0.012 for DZ twins in mothers over 35 years, and the cause of the decrease of stillbirth rate of twins should not be entirely attributed to the reductions of older mothers and higher birth orders. Other factors may also be important, including improvement in medical care.

REFERENCES

1. Barr A, Stevenson AC (1961): Stillbirths and infant mortality in twins. *Ann Hum Genet* 25:131–140.
2. Eriksson AW, Eskola M, Fellman J, Forsius H (1973): The value of genealogical data in population studies in Sweden and Finland. In Morton NE (ed): "Genetic Structure of Populations." Honolulu: University of Hawaii, pp 102–118.
3. Fraccaro M (1957): A contribution to the study of birth weight based on an Italian sample twin data. *Ann Hum Genet* 21:224–236.
4. Imaizumi Y, Inouye E (1979): Analysis of multiple birth rates in Japan. I. Secular trend, maternal age effect, and geographical variation in twinning rates. *Acta Genet Med Gemello* 28:107–124.
5. Japan, Ministry of Health and Welfare. Annual, 1951–1968 and 1974: Vital Statistics, Health and Welfare Statistics and Information Department, Ministry of Health and Welfare, Tokyo.
6. Japan, Ministry of Health and Welfare. (1977): Survey on Socio-Economic Aspects of Vital Events—Plural Births, 1975. Health and Welfare Statistics and Information Department, Ministry of Health and Welfare, Tokyo.
7. Karn MN (1952): Birth weight and length of gestation of twins, together with maternal age, parity and survival rate. *Ann Eugen* 16:365–377.
8. Karn MN (1953): Twin data: A further study of birth weight, gestation time, maternal age, order of birth, and survival. *Ann Eugen* 17:233–248.
9. Karn MN (1953): Data of twin births occurring in 2 English metropolitan hospitals. *Acta Genet Med Gemello* 2:152–163.
10. Weinberg W (1901): Beitrage zur Physiologie und Pathologie der Mehrlingsgeburten beim Menschen. *Archiv fuer die gesamte Physiologie des Menschen und der Tiere* 88:346–430.
11. Yerushalmy J, Sheerar SE (1940): Studies on twins. II. On the early mortality of like-sexed and unlike-sexed twins. *Hum Biol* 12:247–263.

Correspondence: Dr. Yoko Imaizumi, Institute of Population Problems, Ministry of Health and Welfare, 2-2 1-Chome, Kasumigaseki, Chiyoda-ku, Tokyo, Japan.