

# Validity of two short screeners for diet quality in time-limited settings

Helmut Schröder<sup>1,2,\*</sup>, Alejandra Benitez Arciniega<sup>1,2</sup>, Cristina Soler<sup>3</sup>,  
 Maria-Isabel Covas<sup>1,2</sup>, José Miguel Baena-Díez<sup>4</sup> and Jaume Marrugat<sup>3</sup>  
 on behalf of the REGICOR and HERMES investigators†

<sup>1</sup>Cardiovascular Risk and Nutrition Research Group (CARIN-ULEC), Program of Research in Inflammatory and Cardiovascular Disorders (RICAD), IMIM – Hospital del Mar, Biomedical Research Park, c/Doctor Aiguader 88, 08003 Barcelona, Spain; <sup>2</sup>Network of Centers for Biomedical Research (Centros de Investigación Biomédica en Red – CIBER), Physiopathology of Obesity and Nutrition (CIBEROBN), Instituto de Salud Carlos III, Madrid, Spain; <sup>3</sup>Cardiovascular Epidemiology and Genetics Research Group (EGEC-ULEC), Program of Research in Inflammatory and Cardiovascular Disorders (RICAD), IMIM, Barcelona, Spain; <sup>4</sup>Centro de Salud La Marina; IDIAP Jordi Gol, Barcelona, Spain

Submitted 18 October 2010; Accepted 14 June 2011; First published online 23 August 2011

## Abstract

**Objective:** An urgent need in dietary assessment is the development of short tools that provide valid assessments of dietary quality for use in time-limited settings. The present study assessed concurrent and construct validity of the short Diet Quality Screener (sDQS) and brief Mediterranean Diet Screener (bMDSC) questionnaires.

**Design:** Relative validity was measured by comparing three dietary quality indices – the Diet Quality Index (DQI), the modified Mediterranean Diet Score (mMDS) and the Antioxidant Score (ANTOX-S) – derived from the two questionnaires with those from multiple 24 h recalls over 12 months. Construct validity was demonstrated by correlations between average nutrient intake recorded on multiple 24 h recalls and the DQI, mMDS and ANTOX-S derived by the short screeners.

**Setting:** Both short questionnaires were administered to 102 participants recruited from a population-based survey in Spain.

**Results:** DQI, mMDS and ANTOX-S correlated ( $P < 0.001$ ) with the corresponding 24 h recall indices ( $r = 0.61, 0.40$  and  $0.45$ , respectively). Limits of agreement lay between 96 and 126%, 59 and 144% and 61 and 118% for the DQI, ANTOX-S and mMDS, respectively. Dietary intakes of fibre, vitamin C, vitamin E, Mg and K reported on the 24 h recalls were positively associated ( $P < 0.04$ ) with the DQI, mMDS and ANTOX-S indices.

**Conclusions:** The sDQS and bMDSC provide reasonable approximations to food-based dietary indices and accurately situate subjects within the indices constructed for the present validation study.

**Keywords**  
 Validation  
 Short screener  
 Construct validity  
 Diet quality  
 Mediterranean diet

Non-communicable diseases such as CVD and cancer are estimated to be responsible for 60% of world deaths, and dietary habits appear to be strong determinants for their development<sup>(1)</sup>. Therefore, analysis of dietary habits in large populations to identify dietary deficiencies is of paramount importance. FFQ are widely used to estimate food intake in large epidemiological settings. However, full-length FFQ are time consuming for participants and thus are unsuitable for routine clinical use and for non-dietary studies incorporating a broad spectrum of measurements. For this reason, several brief screening

tools have been developed to assess intakes of major food groups. Most of these short dietary questionnaires focus on one or two dietary components<sup>(2–7)</sup>. Only a few brief screeners assess a broader range of dietary intake<sup>(8,9)</sup>, an approach that might help to identify subjects at nutritional risk by indicating their level of adherence to diet quality recommendations. To address the need for a brief, effective instrument, we designed short dietary screeners for two different settings. The brief Mediterranean Diet Screener (bMDSC) is intended to assess adherence to the healthy Mediterranean dietary pattern. The short Diet Quality Screener (sDQS) was created to estimate overall diet quality in primary-care settings. We calculated a modified Mediterranean Diet Score (mMDS) and an

† A full roster of REGICOR investigators and collaborators can be found at [www.REGICOR.org/regicor.inv](http://www.REGICOR.org/regicor.inv)

\*Corresponding author: Email [hschroeder@imim.es](mailto:hschroeder@imim.es)

Antioxidant Score (ANTOX-S) from the bMDSC and a Diet Quality Index (DQI) from the sDQS. The rationale for constructing the mMDSC, ANTOX-S and DQI was adherence to the Mediterranean diet, antioxidant capacity of foods and dietary recommendations for the Spanish population, respectively<sup>(10,11)</sup>. The objective of the present study was first to analyse the degree to which the dietary indices derived from the bMDSC and sDQI questionnaires correlate with those obtained by a reference method (concurrent validity) and second to determine the ability of the screeners to measure diet quality (construct validity) in a random subsample of a population-based cohort in Spain.

## Material and methods

### *Study participants*

The validation study consecutively selected 150 men and women from a population-based cross-sectional survey performed in Girona Province between 2004 and 2006. That survey of 6352 randomly selected free-living men and women, aged 3 to 80 years (71.5% response rate), obtained sociodemographic, anthropometric and lifestyle variables including diet from all participants. Of those initially recruited, 133 agreed to participate in the validation study and 102 completed both short dietary screeners and a 24 h food recall questionnaire on at least ten occasions. The study was approved by the institutional ethics committee (Comités Éticos de Investigación Clínica–Instituto Municipal de Asistencia Sanitaria, Barcelona, Spain), participants signed an informed consent, and results of the examination were sent to all participants.

### *Measurement of non-dietary variables*

Information on demographic and socio-economic variables, co-morbidity history, diet and lifestyle factors, including tobacco smoking and alcohol consumption, was obtained through structured standard questionnaires administered by trained personnel<sup>(12)</sup>.

Leisure-time physical activity was measured by the Minnesota leisure-time physical activity questionnaire, also administered by a trained interviewer<sup>(13,14)</sup>.

### *Dietary assessment instruments*

#### *Multiple 24 h recalls (reference method)*

Monthly 24 h recalls were collected by telephone over a 12-month period by a trained interviewer and data from at least ten recalls were required for inclusion in the analysis. Dietary recalls were conducted on non-consecutive days, including at least five weekdays and one weekend day, and participants were not alerted in any way to the date when they would be interviewed. Food intake data recorded during the 24 h recalls were grouped into the food-based dietary components of the bMDSC and sDQS for analysis.

We validated the dietary quality estimates from the bMDSC and sDQS by comparing them with the data gathered from the 24 h recalls.

#### *The brief Mediterranean diet screener*

The bMDSC was originally developed for another study<sup>(15)</sup>. The objective of this screener was to calculate adherence to the Mediterranean diet and to qualitatively estimate dietary antioxidant intake. For this reason we included food items considered characteristic for the Mediterranean diet and food items with high antioxidant capacity. After participants had completed the multiple 24 h recalls they received the bMDSC by postal mail, together with short instructions on completion of the screener. Participants were asked to report their consumption frequency of fifteen selected food items during the preceding year: (i) pulses; (ii) green leafy and cabbage-like vegetables; (iii) other vegetables; (iv) red meat, sausages and cold cuts; (v) white meat; (vi) blue fish; (vii) white fish; (viii) dairy products; (ix) citrus fruits and berries; (x) other fruits; (xi) wholegrain products; (xii) olive oil; (xiii) nuts; (xiv) juice; (xv) red wine. The eight frequency categories range from 'never' to 'more than 4 times a day'. Standardized portion sizes were used to quantify food intake<sup>(16,17)</sup>.

#### *The short diet quality screener*

The sDQS was developed for use in primary-care settings to estimate overall diet quality. Within 1 week of returning the completed bMDSC, the participants received the sDQS, which was mailed to them with brief instructions on completion of the screener. Subjects were asked to base their responses on their usual dietary behaviours over the previous 12 months, reporting their habitual intake of eighteen food items grouped in three food categories. These categories were based on recommended frequencies of food intake<sup>(11)</sup>. Standardized portion sizes were used to quantify frequency of food intake<sup>(16,17)</sup>. The first category includes eight items: (i) bread; (ii) vegetables (cooked and raw); (iii) fruit; (iv) milk and yoghurt; (v) rice and pasta; (vi) vegetable oils (olive and sunflower); (vii) alcoholic beverages; and (viii) cereals (cornflakes, muesli, etc.). The second category includes seven items: (i) meat; (ii) sausages; (iii) cheese; (iv) pastry; (v) animal fat (butter, lard); (vi) other vegetable oils (palm oil, etc.); and (vii) fast food. The third category had just three items: (i) fish; (ii) legumes; and (iii) nuts.

Food frequency consumption was arranged in three frequency response categories: (i) 'less than once a day', 'once a day' and 'more than once a day' for the eight food items in the first category; (ii) 'less than 4 times a week', '4 to 6 times a week' and 'once a day' for the seven food items in the second category; and (iii) 'less than 2 times a week', '2 to 3 times a week' and '4 or more times a week' for the three food items in the third food category.

The aim of the study was to validate the scores by analysis of the corresponding food items. Therefore, food items listed in the 24 h recall that were not part of the scores obtained by the screeners were omitted.

### Calculation of dietary scores

#### *The antioxidant food and modified Mediterranean diet scores derived from the brief Mediterranean diet screener*

The bMDS provides two sub-scores: the antioxidant food score (ANTOX-S) and the modified Mediterranean diet score (mMDS; Table 1). The standardized portion size of each food group was multiplied by the corresponding frequency response category to obtain the amount of intake in g/d. Tertile distribution of food items included in the ANTOX-S and mMDS was calculated. The ANTOX-S was calculated by assigning 1 point for lowest tertile ratings, 2 points for medium tertile ratings and 3 points for highest tertile ratings for the following food items: (i) citrus fruits; (ii) other fruits; (iii) green leafy vegetables; (iv) other vegetables; (v) fruit juice; and (vii) red wine. The mMDS includes ten food items: (i) vegetables (green leafy vegetables and other vegetables); (ii) fruits (citrus fruits and other fruits); (iii) dairy products; (iv) red meat and sausages; (v) wholegrain products; (vi) fish; (vii) legumes; (viii) nuts; (ix) red wine; and (x) olive oil. Tertile ratings for vegetables, fruits, wholegrain products, legumes, nuts and olive oil were scored as mentioned above. Tertile distribution ratings of two food groups (dairy products; red meat and sausages) were coded inversely (first tertile 3 points, second tertile 2 points, and third tertile 1 point) because these foods are not considered part of the Mediterranean diet. Moderate red wine consumption (1–2 glasses/d) was coded as 3 and

included as a favourable component in the score; no red wine consumption and more than 2 glasses/d were both coded as 1. These values for the food items were added together to determine the ANTOX-S and mMDS, which range from 6 (very low adherence) to 18 (optimal adherence), and from 10 (very low adherence) to 30 (optimal adherence), respectively.

#### *The diet quality index derived from the short diet quality screener*

The diet quality index (DQI) includes three food group categories (Table 2): with the exception of alcoholic beverage consumption, daily intake of 1 portion of foods in the first food group category is scored 2; lower and higher intakes are scored 1 and 3, respectively. Daily consumption of one alcoholic drink (1 bottle of beer, 1 glass of wine, or 1 cup of liquor equivalent to approximately 12g of alcohol) is scored 3; lower and higher intakes are scored 1. Consumption of foods considered detrimental in the second food group category is scored 2 if reported as 4–6 times per week; more and less frequent consumption are scored 1 and 3, respectively. High consumption (4 or more times per week) of food items considered beneficial of the third food group category is scored 3. Intakes of 2–3 times and less than twice a week are scored 2 and 1, respectively. All food item scores are added up. The total possible score ranges from 18 to 54.

### Statistical analyses

Differences in continuous variables with normal distribution were compared between participant and non-participant groups using Student's *t* test; the Mann–Whitney *U* test was used otherwise. The  $\chi^2$  test was used for categorical variables. Relative agreement of the dietary quality indices,

**Table 1** Scoring method for the modified Mediterranean Diet Score (mMDS) and the Antioxidant Score (ANTOX-S)

		Scoring by tertile distribution of food		
		1st tertile	2nd tertile	3rd tertile
Pulses (beans, peas and lentils)*	1 serving	1	2	3
Green leafy and cabbage-like vegetables (lettuce, endive, broccoli, etc.)*††	1 serving	1	2	3
Other vegetables*††	1 serving	1	2	3
Red meat, sausages, cold cuts (pork, beef, lamb, salami, bratwurst, etc.)*	1 serving	3	2	1
White meat (poultry and rabbit)	1 serving	3	2	1
Blue fish (tuna, sardine, salmon, etc.)*§	1 serving	1	2	3
Other fish (codfish, sole, flounder, etc.)*§	1 serving	1	2	3
Dairy products (cheese, whole milk, etc.)*	1 serving	3	2	1
Citrus fruits and berries (oranges, lemons, kiwis, strawberries, etc.)*†	1 piece or 1 serving	1	2	3
Other fruits*†	1 piece or 1 serving	1	2	3
Wholegrain products*	1 serving	1	2	3
Olive oil*	1 tablespoon	1	2	3
Nuts (almonds, walnuts, etc.)*	1 serving	1	2	3
Juicet	1 cup	1	2	3
Red wine*†	1 cup	1	2	3

\*Included in the mMDS.

†Included in the ANTOX-S.

‡Combined in the mMDS 'vegetable' food group.

§Combined in the mMDS 'fish' food group.

||Combined in the mMDS 'fruit' group.

**Table 2** Scoring method for the Diet Quality Index (DQI)

1. Daily frequency consumption of the following foods during the last 12 months				
Food	Amount	<1 time/d	1 time/d	≥2 times/d
Bread	1–2 slices	1	2	3
Vegetable/salad	1 serving	1	2	3
Fruit	1 piece or serving	1	2	3
Yoghurt or milk	1 tub/1 glass	1	2	3
Pasta or rice	1 serving	1	2	3
Oil (olive or sunflower)	1 tablespoon	1	2	3
Alcoholic beverages	1 drink	1	3	1
Breakfast flakes	1 bowl	1	2	3
2. Weekly frequency consumption of the following foods during the last 12 months				
Food	Amount	<4 times/week	4–6 times/week	≥7 times/week
Meat	1 serving	3	2	1
Sausages	1–3 slices	3	2	1
Cheese	1 serving	3	2	1
Pastry or sweets	1 piece or serving	3	2	1
Butter or lard	1 teaspoon	3	2	1
Other vegetable oils	1 tablespoon	3	2	1
Fast food	1 serving	3	2	1
3. Weekly frequency consumption of the following foods during the last 12 months				
Food	Amount	<2 times/week	2–3 times/week	≥4 times/week
Fish	1 serving	1	2	3
Legumes	1 serving	1	2	3
Nuts	1 handful	1	2	3

indicating the agreement between test and reference method in the ranking of subjects, was assessed by calculating Pearson product-moment correlation coefficients to compare the 24 h recall scores (reference method) with the subject's scores on the short screeners (test method). In addition, cross-classification models were fitted to test for gross misclassification of rating for the DQI, mMDS and ANTOX-S obtained from the test *v.* the reference method. Gross misclassification was defined as classification in the opposite tertile (lowest and highest) by the test method compared with the reference method. The proportions of correctly (i.e. similarly) categorized participants were also calculated.

Absolute agreement between scores derived from the short screeners and the 24 h recalls was analysed by the intra-class correlation coefficient (ICC). Additionally, we applied the limits of agreement (LOA) method<sup>(18)</sup> expressed as a percentage. A mean agreement (difference of means) of 100% indicates total agreement between the test and reference method. For each score we plotted the difference between methods against the mean of the two. Linear regression analysis of mean differences of the dietary indices, as the dependent variable, and the mean of the corresponding dietary indices obtained by the test and the reference, as the independent variable, was performed to detect proportional biases (i.e. mean difference was significantly changed by the magnitude ratings for dietary indices). We applied the criteria for a reasonable agreement between estimates according to Ambrosini and colleagues<sup>(19)</sup>: (i) LOA of the dietary indices derived

from the test method were between one-half (50%) and twice (200%) their reference method estimate; and (ii) there were no significant changes of mean differences over the range of average ratings for the dietary indices.

To analyse construct validity, we hypothesized a relationship between a higher scoring for the dietary indices obtained by the short screeners and a more favourable nutrient intake profile reported on the 24 h recalls. General linear modelling procedures (GLM) were used to estimate nutrient intakes according to the tertile distribution of the dietary indices (DQI, mMDS and ANTOX-S). Linear trend was tested by including the categorized variable (tertile distribution of the dietary indices as continuous in this model). The polynomial contrast was used to determine *P* for linear trend for continuous variables and a *post hoc* Bonferroni correction for multiple comparisons was carried out. Differences were considered significant if *P* < 0.05. The SPSS for Windows statistical software package version 15 (SPSS, Inc., Chicago, IL, USA) was used to carry out all statistical analyses.

## Results

With the exception of educational level, no statistically significant differences were observed between the main characteristics of participants included in the validation study and the remaining participants of the population-based survey (Table 3). The DQI, mMDS and ANTOX-S were normally distributed.

**Table 3** Characteristics of the validation study participants and the remaining participants of the population-based cross-sectional survey\*

	Validation study participants (n 102)		Non-participants (n 6250)		P value
	Mean or median	sd, 95% CI or P25–P75	Mean or median	sd, 95% CI or P25–P75	
Age (years)	58.6	12.1	56.6	12.6	0.119
Sex (% women)	49.0	39.3, 58.7	52.3	39.3, 58.7	0.487
Leisure-time physical activity (MET × min/d)	263	157–417	231	119–402	0.233
BMI (kg/m <sup>2</sup> )	27.6	4.2	27.4	5.1	0.562
Obesity (%)†	26.7	18.8, 35.1	24.4	23.3, 25.5	0.587
Education higher than primary school (%)	62.7	53.0, 72.4	50.0	48.8, 51.3	0.011
Low energy reporter (%)‡	17.8	10.1, 25.5	19.3	18.3, 20.3	0.707

P25, 25th percentile; P75, 75th percentile; MET, metabolic equivalent task.

\*Categorical variables are presented as relative frequency (95% CI); continuous variables are presented as mean or median (sd or P25–P75).

†BMI  $\geq 30.0$  kg/m<sup>2</sup>.

‡Ratio of energy intake (reported on a validated full-length FFQ) to BMR of  $<1.2$ .

**Table 4** Correlation coefficients and between-method agreement measurements of dietary quality indices derived from the short Diet Quality Screener (sDQS) and the brief Mediterranean Diet Score Screener (bMDSS) with the reference method (24 h recall)

	sDQS		bMDSC			
	DQI		mMDS		ANTOX-S	
	Mean	sd or 95% CI	Mean	sd or 95% CI	Mean	sd or 95% CI
Corresponding screener	39.3	2.8	18.3	2.7	11.0	2.1
24 h recall	35.5	2.8	20.7	3.0	10.9	2.1
Difference of means*	3.82	3.33, 4.31	-2.44	-3.01, -1.82	0.05	-0.49, 0.39
Proportions of means (%)†	111	109, 112	89	86, 92	101	97, 106
LOA (%)‡	96;126		61;118		59;144	
Regression coefficient§	0.040	-0.162, 0.242	-0.200	-0.503, 0.103	0.061	-0.270, 0.392
ICC		0.32		0.30		0.45
r¶ (screener v. 24 h recall)		0.61		0.40		0.45
Same tertile (%)¶ (screener v. 24 h recall)		48.5		44.0		50.0
Opposite tertile (%)¶ (screener v. 24 h recall)		3.9		11.0		9.0

DQI, Dietary Quality Index; ANTOX-S, Antioxidant Score; mMDS, modified Mediterranean Diet Score.

One hundred and two participants recruited from a population-based survey in Spain were administered both short questionnaires and completed at least ten interviewer-administered 24 h recalls over a 12-month period.

\*Calculated as: screener - 24 h recall.

†Calculated as: (screener/24 h recall) × 100.

‡95% limits of agreement indicating lower and upper limits of agreement.

§Regression coefficient ( $\beta$ ) between mean (dependent variables) and mean differences (independent variables) of the dietary indices obtained by the test and the reference method.


||Intra-class correlation coefficient.

¶Pearson correlation coefficient.

The sDQS significantly overestimated the DQI rating, whereas the opposite finding was observed for the bMDSC, which underestimated the mMDS score (Table 4). Mean difference of the ANTOX-S was close to zero. The test methods' estimates for DQI and mMDS were 11% higher and 11% lower than those of the 24 h recall reference method, respectively (Table 4). The range of LOA was highest for the ANTOX-S and lowest for the mMDS. No significant proportional variation of the agreement between the test and reference method was observed across ratings of the dietary indices (Table 4, Fig. 1).

The ICC between scores obtained by the screeners and the corresponding scores retrieved by the reference method was highest for the ANTOX-S followed by the DQI and mMDS.

Furthermore, Table 4 shows the Pearson correlation coefficients between methods, indicating the capacity of the bMDSC and the sDQS to rank subjects in accordance with their DQI, mMDS and ANTOX-S ratings. Additionally, gross misclassification models were fitted. Pearson correlation coefficients between scores (DQI, mMDS and ANTOX-S) retrieved from both screeners (sDQI and bMDSC) and the corresponding scores obtained from the reference method (multiple 24 h recalls) were calculated. All three dietary scores obtained from both short screeners were significantly correlated with the corresponding scores obtained using the reference method. The correlation coefficient was highest and gross misclassification was lowest for the DQI derived from the sDQS; misclassification was somewhat higher for mMDS and ANTOX-S ratings.



**Fig. 1** Bland–Altman plots indicating the mean difference between indices obtained from the 24 h recall (reference method) and the corresponding dietary assessment method plotted v. the mean of the indices obtained from the two methods for: (a) the Diet Quality Index (DQI) derived from the short Dietary Quality Screener (sDQS); (b) the modified Mediterranean Diet Score (mMDS) derived from brief Mediterranean Diet Score Screener (bMDSC); and (c) the Antioxidant Score (ANTOX-S) derived from the sDQS. One hundred and two participants recruited from a population-based survey in Spain were administered both short questionnaires and completed at least ten interviewer-administered 24 h recalls over a 12-month period. ---- indicates mean difference; -.-.- indicate upper and lower 95% limits of agreement. \*Regression coefficient and statistical significance of the slope from linear regression of the mean of the methods v. the difference between methods

Table 5 shows the construct validity of the DQI, mMDS and ANTOX-S. As expected, a direct association existed between dietary fibre, vitamin C, vitamin E, Mg and K and the tertiles of all three dietary scores. The intakes of folic acid and flavonoids increased across tertile distributions of the DQI and ANTOX-S. In contrast, *trans* fatty acids and the ratio of saturated fat to unsaturated fat decreased with higher ratings for the DQI.

## Discussion

The three food-based indices derived from the sDQS and the bMDSC reasonably ranked populations into similar score levels derived by 24 h recalls. Furthermore, the three dietary indices were positively associated with a beneficial nutrient intake profile.

Following a healthy diet is paramount for physical and mental health<sup>(1,20–22)</sup>. The prevention and treatment of non-communicable diseases such as diabetes, obesity and CVD is strongly influenced through diet<sup>(23)</sup>, and therefore identification of individuals at nutritional risk is important to public health policy and initiatives. However, lengthy comprehensive dietary assessment is a time-consuming process that becomes a real challenge in multidisciplinary epidemiological research and primary-care settings. The sDQS and bMDSC were created to rapidly capture diet

quality based on food intake. The DQI, mMDS and ANTOX-S dietary indices derived from these questionnaires showed moderate to good correlations with the corresponding indices obtained by 24 h recalls. The observed range of Pearson correlation (0.40–0.61) and the average level of gross misclassification (7.7%) are comparable to those reported for food and nutrient intakes derived from short diet screeners and full-length FFQ<sup>(24–28)</sup>. Although a slightly higher correlation has been reported between the Diet Quality Index-Revised (DQI-R) derived from a full-length FFQ and two 1-week dietary records<sup>(29)</sup>, our results suggest that the sDQS and bMDSC adequately rank subjects with respect to the dietary indices' ratings.

However, reasonable relative agreement between methods does not necessarily imply good absolute agreement. To address this issue we plotted mean differences between the dietary indices derived from the test and reference methods against the corresponding mean of both methods and calculated the LOA. The sDQS and the bMDSC systematically overestimated and underestimated the DQI and mMDS ratings, respectively. However, no significant proportional bias was observed for all three dietary indices. Furthermore, the widest range of LOA between methods was found for the ANTOX-S, meaning that 95% of all bMDSC estimates were between 59% (underestimating by 41%) and 144% (overestimating by

**Table 5** Energy and nutrient intakes recorded on 24 h recalls according to tertile distribution of dietary indices derived from the short Dietary Quality Screener (sDQS) and brief Mediterranean Diet Score Screener (bMDSC)

	First tertile		Second tertile		Third tertile		<i>P</i> for trend
	Mean	95% CI	Mean	95% CI	Mean	95% CI	
Energy (MJ)							
DQI	7.4	6.9, 7.9	7.2	6.6, 7.8	7.1	6.5, 7.1	0.205
mMDS	7.0	6.5, 7.5	7.6	7.0, 8.2	7.2	6.6, 7.7	0.697
ANTOX-S	7.2	6.7, 7.7	7.2	6.6, 7.7	7.3	6.6, 8.0	0.870
Carbohydrates (% of energy)							
DQI	40.6	38.6, 42.6	41.9	39.6, 44.1	42.5	40.4, 44.7	0.205
MMDS	42.4	40.9, 44.8	39.3	37.0, 41.5	43.0	40.9, 45.1	0.670
ANTOX-S	41.6	39.7, 43.5	41.4	39.3, 43.4	42.4	39.7, 45.2	0.632
Protein (% of energy)							
DQI	16.3	15.6, 17.0	17.1	16.3, 18.0	17.2	16.4, 18.0	0.088
MMDS	17.2	16.5, 17.9	16.8	15.9, 17.6	16.5	15.7, 17.3	0.196
ANTOX-S	17.3	16.6, 18.0	16.5	15.9, 17.3	16.5	15.5, 17.5	0.163
Fat (%)							
DQI	43.8	42.3, 45.5	43.3	41.6, 45.1	41.5	39.7, 43.2	0.047
MMDS	42.9	41.3, 44.5	44.1	42.1, 45.9	41.8	40.1, 44.0	0.378
ANTOX-S	43.1	41.6, 44.7	42.8	41.4, 44.5	42.5	40.3, 44.7	0.664
Ratio of saturated to unsaturated fat							
DQI	0.51	0.48, 0.54	0.47	0.44, 0.50	0.45	0.42, 0.48	0.008
MMDS	0.47	0.44, 0.51	0.50	0.46, 0.53	0.47	0.44, 0.50	0.790
ANTOX-S	0.48	0.45, 0.51	0.47	0.43, 0.51	0.49	0.45, 0.52	0.854
Total cholesterol (mg)							
DQI	290	263, 317	277	247, 308	253	224, 283	0.007
MMDS	283	256, 310	296	265, 327	248	219, 277	0.087
ANTOX-S	290	264, 316	265	237, 293	262	225, 299	0.224
<i>Trans</i> fatty acids (g)							
DQI	1.07	0.82, 1.31	0.73	0.46, 1.0	0.67	0.41, 0.93	0.031
MMDS	0.92	0.67, 1.17	0.87	0.58, 1.16	0.71	0.44, 0.98	0.262
ANTOX-S	1.0	0.78, 1.24	0.83	0.57, 1.08	0.50	0.15, 0.83	0.015
Fibre (g)							
DQI	12.9	11.5, 14.4	15.2	13.6, 16.4	16.9	15.3, 18.5	<0.001
MMDS	13.5	12.0, 15.0	14.0	12.3, 15.7	17.1	15.5, 18.7	<0.001
ANTOX-S	13.6	12.2, 15.0	14.7	13.2, 16.2	17.6	15.6, 19.6	<0.001
Vitamin C (mg)							
DQI	71.6	57.6, 85.6	110.4	94.8, 125.9	107.6	92.5, 122.6	0.001
MMDS	80.4	65.8, 94.9	94.0	77.3, 110.6	109.5	93.9, 125.1	0.008
ANTOX-S	72.4	60.0, 85.0	99.1	85.4, 112.7	129.2	111.2, 147.1	<0.001
Vitamin E (mg)							
DQI	5.4	4.8, 5.9	6.0	5.4, 6.7	5.9	5.3, 6.6	0.163
MMDS	5.2	4.6, 5.8	5.8	5.2, 6.3	6.3	5.7, 6.9	0.012
ANTOX-S	5.4	4.9, 6.0	5.7	5.1, 6.3	6.4	5.7, 7.2	0.032
Folic acid (µg)							
DQI	184	168, 200	225	207, 242	226	209, 243	0.001
MMDS	198	181, 215	208	189, 228	221	203, 239	0.067
ANTOX-S	193	178, 209	213	196, 230	233	211, 255	0.005
Mg (mg)							
DQI	206	190, 221	232	216, 249	248	232, 264	<0.001
mMDS	207	192, 223	228	210, 245	246	230, 263	0.001
ANTOX-S	216	201, 230	227	211, 244	245	225, 267	0.023
K (g)							
DQI	2.3	2.1, 2.5	2.6	2.5, 2.8	2.7	2.5, 2.9	0.001
MMDS	2.4	2.2, 2.5	2.5	2.3, 2.7	2.7	2.6, 2.9	0.001
ANTOX-S	2.4	2.3, 2.6	2.5	2.4, 2.7	2.8	2.6, 3.0	0.008
Flavonoids (g)							
DQI	48.8	37.4, 60.2	70.6	58.0, 83.1	69.0	56.5, 80.9	0.020
MMDS	53.8	42.4, 65.1	62.2	49.2, 75.1	66.9	54.7, 79.0	0.121
ANTOX-S	49.2	39.1, 59.2	60.7	49.7, 71.7	60.7	69.1, 97.9	<0.001

DQI, Dietary Quality Index; mMDS, modified Mediterranean Diet Score; ANTOX-S, Antioxidant Score.

One hundred and two participants recruited from a population-based survey in Spain were administered both short questionnaires and completed at least ten interviewer-administered 24 h recalls over a 12-month period.

44%) of their reference method estimates. These results indicate a reasonable absolute agreement between the dietary indices derived from the sDQS and the bMDSC and those of the reference method.

In addition to estimating the relative validity of both short dietary screeners studied, we sought to analyse their construct validity, i.e. the degree to which these dietary quality assessment tools measure the theoretical construct

(diet quality) they intend to assess. For this purpose, we hypothesized that higher ratings on the dietary indices should be positively correlated with a healthy nutrient intake profile (i.e. higher intakes of vitamin C, vitamin E, dietary fibre, etc.) In most cases, the sDQS and bMDSC dietary indices correlated as anticipated with the average nutrient intakes derived from multiple 24 h recalls. A healthier nutrient intake profile was related to higher ratings of the DQI, mMDS and ANTOX-S, indicating a reasonable construct validity of the short dietary screeners. Construct validity was somewhat stronger for the sDQS-derived DQI compared with the mMDS and ANTOX-S obtained from the bMDSC. The different food composition of both screeners might particularly explain this finding.

A limitation of the present study is the somewhat higher educational level of participants who completed the validation study.

We conclude that the sDQI and bMDSC accurately rate subjects with respect to the dietary indices derived from these screeners without significant proportional variations over the range of average ratings for the dietary indices. Additionally, the LOA variations are within a reasonable range. Furthermore, both screeners show reasonable construct validity, as indicated by the correlations between the dietary indices derived from the sDQS and bMDSC and nutrient intakes reported on the 24 h recalls. Hence, the sDQS and the bMDSC are valid dietary assessment tools for rapid estimation of dietary quality.

## Acknowledgements

This research was supported by a grant 2FD097-0297-CO2-01 from Fondo Europeo de Desarrollo Regional (FEDER) and by parts of grants from Spain's Ministerio de Sanidad y Consumo, Instituto de Salud Carlos III FEDER (PI080439), Red HERACLES RD06/0009, and by a joint contract of the Instituto de Salud Carlos III and the Health Department of the Catalan Government (Generalitat de Catalunya), CP 03/00115. The CIBEROBN is an initiative of the Instituto de Salud Carlos III, Madrid, Spain. The authors declare no conflict of interest. H.S. led data analysis and writing of the manuscript. J.M. and A.B.A. participated in the discussion and interpretation of results, and writing of the manuscript. J.M.B.-D., C.S. and M.-I.C. provided expertise in data analysis and in interpretation and discussion of results. They made substantial suggestions on the manuscript. All authors read and approved the final manuscript. The authors appreciate the English revision made by Elaine Lilly, PhD (Writers First Aid).

## References

1. The World Health Report (2002) *Reducing Risks, Promoting Healthy Life*. Geneva: WHO.
2. Wiens L, Schulzer M, Chen C *et al.* (2010) Reliability and validity of the SmartDiet Canadian version questionnaire. *J Am Diet Assoc* **110**, 101–105.
3. Spencer EH, Elon LK, Hertzberg VS *et al.* (2005) Validation of a brief diet survey instrument among medical students. *J Am Diet Assoc* **105**, 802–806.
4. Cena H, Roggi & Turconi G (2008) Development and validation of a brief food frequency questionnaire for dietary lutein and zeaxanthin intake assessment in Italian women. *Eur J Nutr* **47**, 1–9.
5. Thompson FE, Midthune D, Subar AF *et al.* (2004) Performance of a short tool to assess dietary intakes of fruits and vegetables, percentage energy from fat and fibre. *Public Health Nutr* **7**, 1097–1105.
6. Nelson MC & Lytle LA (2009) Development and evaluation of a brief screener to estimate fast-food and beverage consumption among adolescents. *J Am Diet Assoc* **109**, 730–734.
7. Thompson FE, Midthune D, Subar AF *et al.* (2007) Development and evaluation of a short instrument to estimate usual dietary intake of percentage energy from fat. *J Am Diet Assoc* **107**, 760–767.
8. Mochari H, Gao Q & Mosca L (2008) Validation of the MEDFACTS dietary assessment questionnaire in a diverse population. *J Am Diet Assoc* **108**, 817–822.
9. Rifas-Shiman SL, Willett WC & Lobb R (2001) PrimeScreen, a brief dietary screening tool: reproducibility and comparability with both a longer food frequency questionnaire and biomarkers. *Public Health Nutr* **4**, 249–254.
10. Willett WC, Sacks F, Trichopoulos A *et al.* (1995) Mediterranean diet pyramid: a cultural model for healthy eating. *Am J Clin Nutr* **61**, 6 Suppl, 1402S–1406S.
11. Aranceta J & Serra-Majem L, Working Party for the Development of Food-Based Dietary Guidelines for the Spanish Population (2001) Dietary guidelines for the Spanish population. *Public Health Nutr* **4**, 1403–1408.
12. Baena-Díez JM, Alzamora-Sas MT, Grau M *et al.* (2009) Validity of the MONICA cardiovascular questionnaire compared with clinical records. *Gac Sanit* **23**, 519–525.
13. Elosua R, Garcia M, Aguilar A *et al.* (2000) Validation of the Minnesota Leisure Time Physical Activity Questionnaire in Spanish women. Investigators of the MARATDON Group. *Med Sci Sports Exerc* **32**, 1431–1437.
14. Elosua R, Marrugat J, Molina L *et al.* (1994) Validation of the Minnesota Leisure Time Physical Activity Questionnaire in Spanish men. The MARATHOM Investigators. *Am J Epidemiol* **139**, 1197–1209.
15. Peters A, Schneider A, Greven S *et al.* (2007) Air pollution and inflammatory response in myocardial infarction survivors: gene–environment interactions in a high-risk group. *Inhal Toxicol* **19**, Suppl. 1, 161–175.
16. Serra Majem L (2004) *Guía de Alimentación Saludable*. Madrid: Sociedad Española de Nutrición Comunitaria.
17. Salas Salvado J, Bonada A, Tralero R *et al.* (editors) (2000) *Nutrición y Dietética Clínica*. Barcelona: Editorial Masson.
18. Bland JM & Altman DG (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **1**, 307–310.
19. Ambrosini GL, van Roosbroeck SA, Mackerras D *et al.* (2003) The reliability of ten-year dietary recall: implications for cancer research. *J Nutr* **133**, 2663–2668.
20. Féart C, Samieri C & Barberger-Gateau P (2010) Mediterranean diet and cognitive function in older adults. *Curr Opin Clin Nutr Metab Care* **13**, 14–18.
21. Muñoz MA, Fito M, Marrugat J *et al.* (2009) Adherence to the Mediterranean diet is associated with better mental and physical health. *Br J Nutr* **101**, 1821–1827.
22. Schroder H (2007) Protective mechanisms of the Mediterranean diet in obesity and type 2 diabetes. *J Nutr Biochem* **18**, 149–160.



23. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series no. 916.* Geneva: WHO.
24. Schroder H, Covas MI, Marrugat J *et al.* (2001) Use of a three-day estimated food record, a 72-hour recall and a food-frequency questionnaire for dietary assessment in a Mediterranean Spanish population. *Clin Nutr* **20**, 429–437.
25. Chan SG, Ho SC, Kreiger N *et al.* (2008) Validation of a food frequency questionnaire for assessing dietary soy isoflavone intake among midlife Chinese women in Hong Kong. *J Nutr* **138**, 567–573.
26. Osowski JM, Beare T & Specker B (2007) Validation of a food frequency questionnaire for assessment of calcium and bone-related nutrient intake in rural populations. *J Am Diet Assoc* **107**, 1349–1355.
27. Sevak L, Mangtani P, McCormack V *et al.* (2004) Validation of a food frequency questionnaire to assess macro- and micro-nutrient intake among South Asians in the United Kingdom. *Eur J Nutr* **43**, 160–168.
28. Fernández-Ballart JD, Piñol JL, Zazpe I *et al.* (2010) Relative validity of a semi-quantitative food-frequency questionnaire in an elderly Mediterranean population of Spain. *Br J Nutr* **103**, 1808–1816.
29. Newby PK, Hu FB, Rimm EB *et al.* (2003) Reproducibility and validity of the diet quality index revised as assessed by use of a food-frequency questionnaire. *Am J Clin Nutr* **78**, 941–949.