
ERRATUM

Consensus guideline on parental methadone use in pain and palliative care

LAUREN SHAIOVA, M.D.,¹ ANN BERGER, M.D.,² CRAIG D. BLINDERMAN, M.D., M.A.,³
EDUARDO BRUERA, M.D.,⁴ MELLAR P. DAVIS, M.D., F.C.C.P.,⁵ SUSAN DERBY, N.P.,⁶
CHARLES INTURRISI, PH.D.,⁷ JILL KALMAN, M.D.,⁸ DEVENDRA MEHTA, M.D., PH.D., F.R.C.P.,⁹
MARCO PAPPAGALLO, M.D.,¹⁰ AND EUGENE PERLOV, M.D.¹¹

¹Department of pain Medicine and Palliative Care, Metropolitan Hospital Center, Health and Hospital Corporation of New York City, New York, New York

²Pain and Palliative Care Physician, Bethesda, Maryland

³Palliative Care Service, Department of Medicine, Massachusetts General Hospital and the Harvard Medical School Center for Palliative Care, Boston, Massachusetts

⁴Department of Palliative Care and Rehabilitation Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas

⁵Harry R. Horvitz Center for Palliative Medicine, Cleveland Clinic Foundation, Cleveland, Ohio

⁶Pain and Palliative Care Service, Memorial Sloan-Kettering Cancer Center, New York, New York

⁷Weill Cornell Medical College, New York, New York

⁸Cardiovascular Institute, Mount Sinai Medical Center, New York, New York

⁹Section of Cardiac Electrophysiology, Mount Sinai Hospital and School of Medicine, New York, New York

¹⁰Mount Sinai School of Medicine, New York, New York

¹¹Visiting Nurse Service of New York Hospice Care, New York, New York

(Volume 6, Number 2, 2008, Pages 165–176; doi:10.1017/S1478951508000254)

Due to a production error, the wrong affiliation for Lauren Shaiova was given. The above is the correction affiliation for all authors in this article.