
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSIDE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

Contents

	Pages
Observations on the morphology of <i>Toxocara pteropodis</i> eggs. P. PROCIV	271-277
Evaluation of chemotherapy in experimental toxocarosis by determination of specific immune complexes. C. CUELLAR, S. FENOY, C. AGUILA and J. L. GUILLEN	279-289
Apparent absence of genetic differences among species of <i>Teladorsagia</i> (Nematoda: Trichostrongylidae). R. H. ANDREWS and I. BEVERIDGE	290-294
Possible direct effect of diethylcarbamazine on the infective larvae of <i>Brugia pahangi</i> . Y. FUJIMAKI, M. SHIMADA, Y. MITSUI, E. KIMURA and Y. AOKI	295-301
Analysis of polyunsaturated fatty acid composition of <i>Strongyloides ratti</i> in relation to development. T. MINEMATSU, S. YAMAZAKI, Y. UJI, H. OKABE, M. KORENAGA and I. TADA	303-309
Immunoblot analysis of serum IgG, IgA and IgE responses against larval excretory-secretory antigens of <i>Anisakis simplex</i> in patients with gastric anisakiasis. N. AKAO, T. OHYAMA and K. KONDO	310-318
Prevalence of <i>Cooperia punctata</i> , <i>C. pectinata</i> and <i>C. oncophora</i> infections in dairy calves in Brazil. M. P. GUIMARAES, H. M. A. COSTA and W. S. LIMA (Research Note)	319-322
Variation in the morphology of the dorsal and dorso-lateral tegument of male <i>Schistosoma haematobium</i> from southern Africa. F. J. KRUGER and C. T. WOLMARANS	323-329
Do all human urinary infections with <i>Schistosoma mattheei</i> represent hybridization between <i>S. haematobium</i> and <i>S. mattheei</i> ? F. J. KRUGER and A. C. EVANS (Research Note)	330-332
Frequency and possible consequences of hybridization between <i>Schistosoma haematobium</i> and <i>S. mattheei</i> in the Eastern Transvaal Lowveld. F. J. KRUGER (Research Note)	333-336
Antagonistic effects of <i>Schistosoma mansoni</i> on superimposed <i>Hymenolepis diminuta</i> and <i>H. microstoma</i> infections in mice. J. ANDREASSEN, A. B. ODAIBO and N. Ø. CHRISTENSEN (Research Note)	337-339
The rate of development and longevity of the monogenean skin parasite <i>Entobdella soleae</i> . G. C. KEARN (Research Note)	340-342
An experimental <i>in vitro</i> model for evaluating drugs against protoscolecids of <i>Echinococcus granulosus</i> . R. LORENZINI and A. RUGGIERI	343-348
Announcements	278, 302
List of Referees	349-350
Subject index	351-354
Author index	355-357
Contents of Volume	i-iv