September 1959

Vol. LXXIII No. 9

The Journal of Laryngology and Otology

EDITED BY WALTER HOWARTH

> ASSISTANT EDITOR G. H. BATEMAN

Contents

MASTOIDITIS

THE RECONSTRUCTION OF THE MOUTH OF OESOPHAGUS BY USING THE LARYNX. ONE STAGE OPERA-TION

- -

CAROTID ABNORMALITIES AND ADENOID SURGERY

POSTERIOR CHOANAL ATRESIA

CLINICAL RECORDS— Rhabdomyosarcoma of the Nasopharynx

MONILIASIS OF THE LARYNX AND HYPOPHARYNX.

A Case of Nasal Hæmangioma with Unusual Features

IDIOPATHIC RECURRENT LARYNGEAL PARALYSIS GENERAL NOTES

London Headley Brothers Ltd

Annual Subscription £4/4/0 net, U.S.A. \$13

https://doi.org/10.1017/S0022215100055729 Published online by Cambridge University Press

Monthly, 10/- net post free

TAUNO PALVA and KARI PULKKINEN

PROFESSOR DR. A. ŠERCER


WILLIAM MCKENZIE and CYRIL I. WOOLF

DAVID H. CRAIG and NOREEN M. SIMPSON

H. N. PERKINS and P. D. STEWART J. J. LEES

> M. E. N. SMITH W. M. Mollison

ADVERTISEMENTS



THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$14.00 per year

Canada \$13.00 per year

MAX A. GOLDSTEIN, M.D. Founder THEODORE E. WALSH, M.D. EDITOR

640 SOUTH KINGSHIGHWAY SAINT LOUIS 10, MO.

CONTENTS

	PAGE
MASTOIDITIS. Tauno Palva and Kari Pulkkinen (Turku)	573
THE RECONSTRUCTION OF THE MOUTH OF OESOPHAGUS BY USING THE LARYNX. ONE STAGE OPERATION. Professor Dr. A.	
Šercer (Zagreb, Jugoslavia)	589
CAROTID ABNORMALITIES AND ADENOID SURGERY, William McKenzie	
and Cyril I. Woolf (London)	596
POSTERIOR CHOANAL ATRESIA. David H. Craig and Noreen M.	
Simpson (Belfast)	603
CLINICAL RECORDS	
Rhabdomyosarcoma of the Nasopharynx. H. N. Perkins and	
P. D. Stewart (Dumbarton)	612
Moniliasis of the Larynx and Hypopharynx. J. J. Lees (Glasgow)	619
A Case of Nasal Hæmangioma with Unusual Features. M. E. N.	
Smith (Swansea)	621
Idiopathic Recurrent Laryngeal Paralysis. W. M. Mollison	
(Ringwood)	624
General Notes	627

For Advertisement space in this Journal apply to: HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

The Journal of Laryngology and Otology

(Founded in 1887 by Morell MACKENZIE and Norris Wolfenden)

EDITED BY WALTER HOWARTH

ASSISTANT EDITOR G. H. BATEMAN

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization 1952.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of f_{10} per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., 109 Kingsway London, WC2.

5. Orders for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2. 8. The annual subscription is four guineas sterling (U.S.A. \$13) post free,

and is payable in advance. 9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol.

LXIII may be purchased at 4s. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2.

United States of America

Orders for this Journal may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2, England.

In schools for the DEAFmultitone Telesonic INDUCTION LOOP SYSTEM brings new advantages


- 1 Can be installed in adjoining classrooms without risk of 'spill over'.
- 2 Combines good hearing with complete freedom of movement. Unaffected by distance between pupil and teacher.
- 3 Clear speech at high amplification.
- 4 Any number of pupils may use the same installation.
- 5 The child can use the instrument as a normal hearing aid outside school.



The Kindergarten Class at the London Residential School for Jewish Deaf Children where all classrooms have been installed with the Multitone Telesonic Induction Loop System Write for further details to: **MULTITONE** ELECTRIC COMPANY LTD., 12-20 UNDERWOOD STREET, LONDON, N.I.

Please mention The Journal of Laryngology and Otology when replying to advertisements


CRC M23




INDICATED in the treatment of otitis externa and chronic suppurative otitis media.


EFFECTIVE, in the presence of blood, pus and tissue debris against many organisms including pyogenic cocci, some Gram-negative bacilli and a number of pathogenic fungi.

CONVENIENT, for topical application in the form of drops or as a ribbon gauze dressing, in 15 c.c. dropper bottles and 100 c.c. bottles.



Please mention The Journal of Laryngology and Otology when replying to advertisements





DIAGNOSIS

vi

The excellent optical performance of the Hallpike Ear Microscope gives the user greatly increased confidence in the diagnosis of abnormal conditions; and many disorders may be recognized at an earlier stage than has hitherto been possible with "headlamp and speculum" techniques. Thus slight pulsation of swellings and vascular tumours—of high diagnostic significance and almost impossible to detect by conventional methods—can readily be detected under

this microscope.

Use of the Siegle attachment permits observation of small bubbles present in fluid collections within the tympanum; it is also invaluable for demonstrating minute perforations. Where defects of the tympanic membrane are present, the middle-ear cavity can be scrutinized minutely; the detailed structure of the stapes, stapedius tendon, round window and Eustachian orifice—not normally visible by ordinary methods of examination—are seen with quite remarkable clarity when the Ear Microscope is used.

TREATMENT

Manipulative procedures may be undertaken with greater confidence and precision if the Ear Microscope and its special instruments are used.

The curettes enable adherent wax and debris to be removed with the greatest precision from the walls of the deep meatus or from the tympanic membrane itself. Cholesteatomatous masses or neoplasms may be examined critically and biopsy specimens obtained by means of the curettes. These may also be used as delicate probes for palpation of the incus and head of the stapes in certain cases of otosclerosis with retraction and atrophy of the tympanic membrane; in this way unmistakable evidence may be obtained of bony fixation of the footplate. Or again, it is sometimes possible to identify by palpation defects of the bony tympanic ring or meatal wall which may underlie granulations.


Myringotomy or removal of foreign bodies can be undertaken with surprisingly high precision because the optics of the Ear Microscope and the specially angled myringotome blade eliminate all shadows so that the surgeon can see clearly where and what he is cutting. The outfit is invaluable also for postoperative examinations and treatment of mastoid cavities. Granulations can be cauterized and exposed areas of dura mater carefully palpated.

*J. Laryng. (1953) 67, 108.


39 WIGMORE STREET LONDON W.1



5241 WHITBY AVENUE PHILADELPHIA 43, Pa, U.S.A.



Please mention The Journal of Laryngology and Otology when replying to advertisements



'OTOSPORIN' clears infected ears

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' brand Drops.

'Otosporin' is an aqueous suspension containing both neomycin (5 mgm. per c.c.) and polymyxin B sulphate (10,000 units per c.c.) which are together effective against practically all the bacteria found in ear infections. They are unlikely to induce bacterial resistance or skin sensitisation, and neither of them gives rise to cross-resistance or cross-sensitisation to penicillin and other antibiotics. Their effectiveness is visibly enhanced by the hydrocortisone (5 mgm. per c.c.) in 'Otosporin'; this, by reducing inflammation, not only relieves pain, but provides greater access for the antibiotics.

'OTOSPORIN... drops

issued in bottles of 5 c.c.

