

Scaling relations in early-type galaxies from integral-field stellar kinematics

M. Cappellari,¹ N. Scott,¹ K. Alatalo,² L. Blitz,² M. Bois,³ F. Bournaud,⁴ M. Bureau,¹ R. L. Davies,¹ T. A. Davis,¹ P. T. de Zeeuw,^{5,14} E. Emsellem,^{5,3} J. Falcon-Barroso,⁶ S. Khochfar,⁷ D. Krajnovic,¹ H. Kuntschner,⁵ P.-Y. Lablanche,³ R. M. McDermid,⁸ R. Morganti,⁹ T. Naab,¹⁰ M. Sarzi,¹¹ P. Serra,⁹ R. C. E. van den Bosch,¹² G. van de Ven,¹³ A. Weijmans¹⁴ and L. M. Young¹⁵

¹University of Oxford, UK; ²University of California, Berkeley, USA; ³Université de Lyon, France; ⁴Université Paris Diderot, France; ⁵ESO, Garching, Germany; ⁶IAC, La Laguna, Spain; ⁷MPI for Extraterrestrial Physics, Garching, Germany; ⁸Gemini Observatory, Hilo, USA; ⁹ASTRON, Dwingeloo, The Netherlands; ¹⁰Universitäts-Sternwarte München, Germany; ¹¹University of Hertfordshire, Hatfield, UK; ¹²The University of Texas, Austin, USA; ¹³IAS, Princeton, USA; ¹⁴Leiden University, The Netherlands; ¹⁵New Mexico Tech, Socorro, USA

Early-type galaxies (ETGs) satisfy a now classic scaling relation $R_e \propto \sigma_e^{1.2} I_e^{-0.8}$, the Fundamental Plane (FP; Djorgovski & Davis 1987; Dressler *et al.* 1987), between their size, stellar velocity dispersion and mean surface brightness. A significant effort has been devoted in the past twenty years to try to understand why the coefficients of the relation are not the ones predicted by the virial theorem $R_e \propto \sigma_e^2 I_e^{-1}$.

Recent studies, using independent approaches from either (i) detailed dynamical models or (ii) strong galaxy lensing, point to a genuine variation of the mass-to-light ratio M/L in galaxies as the reason for nearly all the observed ‘tilt’ in the FP (e.g. Cappellari *et al.* 2006; Bolton *et al.* 2008). However these studies are limited by a small and biased sample or are restricted to only the most massive ETGs respectively.

We overcome both limitations by modeling the stellar dynamics, using axisymmetric Jeans anisotropic models (JAM; Cappellari 2008), for the K -band selected, volume-limited ATLAS^{3D} sample of 263 nearby ETGs, spanning a large range of masses and with $60 < \sigma_e < 350$ km s⁻¹. A key for the project is the availability for all galaxies of high-quality integral-field kinematics observed with the SAURON spectrograph and detailed Multi-Gaussian Expansion (Emsellem *et al.* 1994) models of the photometry.

We confirm the genuine M/L variation and construct both the FP and the More FP (MFP; Bolton *et al.* 2007) for the ATLAS^{3D} sample, relating the mean surface density $\Sigma_e \equiv I_e \times (M/L)_{\text{JAM}}$, σ_e and R_e . Our MFP produces a relation as tight as the FP over the full mass range. We compare the global $(M/L)_{\text{JAM}}$ variation among galaxies with predictions from two-SSP stellar population models and find that variations of both dark matter (or IMF) and population are required to explain the observations.

References

- Bolton, A. S., *et al.* 2008, *ApJ*, 684, 248
Bolton, A. S., Burles, S., Treu, T., Koopmans, L. V. E., & Moustakas, L. A. 2007, *ApJ*, 665, L105
Cappellari, M., *et al.* 2006, *MNRAS*, 366, 1126
Cappellari, M. 2008, *MNRAS*, 390, 71
Djorgovski, S. & Davis, M. 1987, *ApJ*, 313, 59
Dressler, A., *et al.* 1987, *ApJ*, 313, 42
Emsellem, E., Monnet, G., & Bacon, R. 1994, *A&A*, 285, 723