

THE JOURNAL OF AGRICULTURAL SCIENCE

EDITED BY

G. D. H. BELL, C.B.E., PH.D., F.R.S., Plant Breeding Institute, Cambridge

K. L. BLAXTER, PH.D., N.D.A. (HONS.), D.SC., F.R.S.E., F.R.S.,
Rowett Research Institute, Bucksburn, Aberdeen

G. W. COOKE, PH.D., F.R.I.C., F.R.S., Rothamsted Experimental Station, Harpenden

JOHN HAMMOND, Jr., M.A., School of Agriculture, Cambridge

PROF. SIR J. B. HUTCHINSON, C.M.G., SC.D., F.R.S., School of Agriculture,
Cambridge

H. L. PENMAN, O.B.E., PH.D., F.R.S., Rothamsted Experimental Station, Harpenden

W. J. RIDGMAN, M.A., School of Agriculture, Cambridge

H. H. ROGERS, B.SC., DIP.AG.SCI., Plant Breeding Institute, Cambridge

PROF. E. W. RUSSELL, C.M.G., PH.D., F.INST.P., Department of Soil Science,
University of Reading

F. YATES, C.B.E., SC.D., F.R.S., Rothamsted Experimental Station, Harpenden



CAMBRIDGE UNIVERSITY PRESS

BENTLEY HOUSE, 200 EUSTON ROAD, LONDON, N.W. 1

AMERICAN BRANCH: 32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Published once every two months

Annual subscription £9 net (U.S.A. \$29.00)

Single parts £2 net (U.S.A. \$6.00)

Second class postage paid at New York, N.Y.

INSTRUCTIONS TO AUTHORS

Failure to comply with the Instructions to Authors may delay publication

Papers intended for publication should be submitted to Dr G. D. H. Bell, C.B.E., F.R.S., Plant Breeding Institute, Cambridge, CB2 2LQ or to one of the Associate Editors. It must be understood that, if accepted by the Editorial Board, the paper will not be published elsewhere in the same form.

MANUSCRIPTS. Papers, written in English, are accepted from any country and should be typed in double-spacing on one side of the paper with a margin at least 4 cm wide on the left-hand side. Authors should instruct typists on the style required. A top copy and one carbon copy should be submitted.

SHORT NOTES may be accepted provided they are based on adequate experimental evidence; special provision is made for their publication with the least possible delay. MSS should not exceed 1500 words in length or their equivalent. For tabulated matter allow 25 words per line of the table (including headings). For line illustrations allow 225 words per quarter of a page.

TABLES must be self-explanatory. They should be typed on separate sheets, numbered consecutively and carry an appropriate title. When possible, tables should be arranged so that they can be printed in the normal orientation of the text and without rules.

LINE DRAWINGS (with photocopies) should be 25 cm wide and drawn in black waterproof ink on Bristol board, graph paper with blue lines or tracing paper. Legends should be typed on a separate copy and numbering inserted lightly and clearly in soft pencil on the drawing. Tables and figures should not reproduce the same data.

The approximate position of tables and figures should be noted in the text.

PLATES should make a definite contribution to the value of the paper and the number submitted should be kept to a minimum. They should be good quality, unmounted, glossy prints and be lightly numbered in pencil on the reverse side. If several, or coloured, plates are submitted the author may be asked to contribute to the cost of reproducing them.

TITLE. The title must be specific and suitable for indexing by the mechanical methods now being employed. The full name and address of the institution in which the research has been carried out should be stated. Change of address may be given as a footnote. A short title, not exceeding 50 characters, should be provided for the running headlines.

STYLE. Experimental details and results should be recorded in the past tense and there should be no unnecessary repetition or loose phrases. Manuscripts are likely to be returned for modification if the presentation is not clear and precise.

LAYOUT. Authors are recommended to study '*General Notes on the Preparation of Scientific Papers*' (Royal Society, London, 2nd edn, 1965). The Editorial Board do not insist on a rigid format but it is usually convenient to divide the paper into sections, e.g. Introduction, Materials and Methods, Results, and Discussion. An excess of headings and sub-headings should be avoided.

Authors are advised to note the following points: a detailed review of literature is not necessary; relevant details should be given of the plant or animal material, the experimental design and chemical or other techniques employed; mean results with their relevant standard errors should be presented rather than detailed data; the statistical methods used should be clearly stated; the discussion should relate the author's experiments to other work on the subject and give the author's conclusions. Footnotes should be avoided.

SUMMARIES of papers are placed at the beginning of the text and authors should submit MSS with the summaries so placed. The summary should be factual and suitable for use in abstracting journals; paragraphs should not be numbered.

REFERENCES. The bibliography must be given in the form—Surname of authors, initials, year of publication (in parentheses), title of paper, name of journal (abbreviated according to the *World List of Scientific Periodicals*, 4th edn, Butterworths, London), volume and pages of reference. References should be in alphabetical order. In the text a reference should be quoted by the author's name and date (in parentheses). Where there are more than two authors, the initial reference in the text should include the names of all authors, but subsequent citations should be in the form—first author followed by *et al.* Authors should check that all references in the text appear at the end of the paper and vice versa, and that names and dates correspond in the two places.

PROOF CORRECTION. Standard proof correction marks (British Standard 1219) should be made as legibly as possible in ink, not pencil. Directions to the printer which are not to be set up in type should be encircled. Captions to illustrations and all references should be checked. Queries marked by the printer should be answered. Proofs are provided in order that authors can check the correctness of the type-setting—excessive alterations may be charged to the author.

OFFPRINTS. Contributors will receive 25 copies of their papers free and can order others when they receive the proofs.

Gardona

the spray-safe insecticide

Gardona is a new insecticide from Shell. And it's a spray-safe insecticide because it can be sprayed without the use of special protective clothing. What's new about Gardona is its low toxicity to man and animals.

Gardona is especially effective against caterpillars, flies and beetles. It protects apples and pears from Codling moth, citrus fruits from fruit flies, and vegetables from caterpillars and flea-beetles.

And Gardona's selective activity against these pests means most beneficial insects and organisms survive the spraying.

Although Gardona will control pests for up to fourteen days, depending on the dosage, its low mammalian toxicity means crops may be safely harvested a very short time after treatment. In most cases the following day.

Spray-safe Gardona is also showing promising results in the control

of major pests in a variety of other crops such as cotton, rice and maize.

Gardona is another product of Shell's continuing research into the development of effective and safe insecticides. For further information or supplies of Gardona contact your Shell company or Shell Chemicals distributor.

Shell Chemicals



Azodrin

does its job inside out-outside in

... and that means double trouble for cotton pests, because Shell's powerful new organo-phosphorus insecticide Azodrin works *two* ways.

By contact, immediately destroying insects on the plant surface. And systemically, by penetrating foliage rapidly where it kills sucking and chewing pests as they try to eat.

This sophisticated dual action makes Azodrin by far the most wide-ranging insecticide yet devised giving *continuing* control over *more* pests *more* effectively.


Azodrin has already been employed with outstanding success in the United States and

Central America, and is now proving world-wide to be one of the cotton grower's most powerful allies.

Suddenly pests that took a bite out of profits are losing their appetite.

For further information or supplies of Azodrin contact your Shell company or Shell chemicals distributor. Shell works today for your more profitable tomorrow.

Shell Chemicals



(ii)

THE JOURNAL OF AGRICULTURAL SCIENCE

Because of the increasing number of papers of high standard received by the Editors, the size of each volume of The Journal of Agricultural Science will be increased in 1970. Cambridge University Press regret that there will also have to be an increase in price. The subscription will be £6 net per volume (£12 net per year). In the U.S.A. the subscription will be \$19.50 per volume (\$39.00 per year).

Aldrin

makes the earth a healthy place for crops to grow in

Aldrin is a highly effective Shell insecticide.

It works *underground* to destroy the full range of soil insect pests, which attack the vital root system of your crop. For the major groups of soil pests, wireworms, cutworms, white grubs and mole-crickets, it is recognised as the most efficient insecticide there is. It can be applied to any type of soil in the world, and is harmless to beneficial micro-organisms in the soil.


Used for row treatment, as an overall application, as a root dip, or as a seed-treatment, aldrin is especially suited for use with maize and other cereals, sugar cane, sugar beet, potatoes and tobacco.

Aldrin has excellent residual action. It is stable in the presence of other agricultural chemicals, does not affect germination, and has no taint. Used as directed, aldrin is harmless to both farmers and consumers. It is available as the technical material and high

concentrate dusts for further formulation, and also in forms suitable for the grower: emulsifiable concentrates, wettable powders, field strength dusts and granules.

For details of approved uses and advice on application of aldrin consult your Shell company or Shell chemicals distributor.

Shell Chemicals



CORRIGENDUM

**The effect of maturity and leafiness on the intake and digestibility
of alfalfas and grasses fed to sheep**

BY J. E. TROELSEN AND J. B. CAMPBELL

J. agric. Sci. **73**, 145–154

The Press regrets the following error:

P. 150, line 7 of caption to Fig. 1: $Y = 85.3 + 3.58X$

Should read: $Y = -85.3 + 3.58X$