BRIEF SUMMARY of PRESCRIBING INFORMATION INDICATIONS AND USAGE: Bipolar Mania: SEROQUEL is indicated for the short-term treatment of acute manic INDICATIONS AND USAGE Bypolar Mania: SERDOUEL is indicated for the short-term treatment of acuse manic episodes associated with bipolar indicated, as either monotherapy or adjunct therapy to lithium or deletioner. The efficacy of SERDOUEL in acuse bypolar mania was established in two 3-week monotherapy trails and one 3-week adjunct the SERDOUEL in acuse bypolar mania was established in two 3-week monotherapy trails and one 3-week monotherapy trails and one 3-week manifestable in the second of the second period and the second of the second period and the second of the se

CONTRAINDICATIONS: SEROQUEL is contraindicated in individuals with a known hypersensitivity to this medica-

Persistation of Confusion interfaces are present inselfusions of the throughout the individual patient.

CONTRAINBUCKTORIS: ENDOUGLE is contraindicated in individuals with a known hypersembility to this medication of any for its ingedieters.

WARRINGS: Neurolegic Malignant Syndrome (NMS): A potentially fatal symptom complex sometimes referred to a Neurolegic Malignant Syndrome (NMS): A potentially fatal symptom complex sometimes referred to a Neurolegic Malignant Syndrome (NMS): A potentially fatal symptom complex sometimes referred to a Neurolegic Malignant Syndrome (NMS): As been reported in association with administration of artiflexychotic drugs, including SEPDOUEL. Brain can see a complex sometimes referred to a new property of the complex sometimes and the complex sometimes and the complex sometimes and the complex sometimes are neglected with SEPDOUEL. Clinical manifestations of the complex must be complex sometimes and the soft sometimes and the complex some

PRECAUTIONS: General: Orthostatic Hypotension: SEROQUEL may induce orthostatic hypotension associated

pypergycenta king esperament with adaptival artitiscypcionics should undergo tasking lowed places lesting. It same cases, hyperglycernal has resolved when the adynctal antipsychotic was document, blood places lesting. It same cases, hyperglycernal has resolved when the adynctal antipsychotic was document of an advantage of communication of anti-diabetic retainment despte discontinuation of the suspect drug.

PRECAMTIONIS: General: Orthostatic Hypotensian: SEROOUEI, may induse orthostatic hypotensian associated with diaziness, subsycration and, in some petents, syrrouge, sepacially curing the initial doses thration period, probably reflecting its or, addressing antipage and properties. Syrrouge was reported in 1% (202567) of the patients streated with SEROOUEI, compared with 0% (000707) on patients and about of a control and system of the patients are advantaged and advantaged and about of the patients are advantaged and advantaged and advantaged and properties and advantaged and treatment with a minimal properties and advantaged and properties and advantaged and advanta

The racing patients of combustation in production in Information for Pollantic Physicians are advised to discuss the following issues with patients for whom they prescribe EFGOULE. Untributative Hypotesian Patients Should be advised of the rook of orbitation by protestic passessably during the 3-50 years of a rindown. Excitation of the patients of the rook of the rook of orbitation by technical bearest event association and School and the rook of the rook of the rindown of the rook of the rook

stration, and careful monitoring during the Intial dosing period in the elderly. The mean plasma clearance of SERCOULE was reduced by 30% to 50% in elderly patients when compared in younger patients. SERCOULE was reduced by 30% to 50% in elderly patients exposed to SERCOULE for the treatment of acute highest mainst (monitoring and adjunct the patients exposed to 15 RFOQUEL for the treatment of acute highest mainst (monitoring and adjunct the patients) and approximately 2000 patients and roman subjects exposed to 1 or more doses of SERCOULE for the treatment of schoolpreans. Of these approximately 3000 subjects, approximately 2001 patients and two participated in multiple dose effectiveness trials, and their experience corresponded to approximately 914.3 patient-years. The conditions and duration of treatment with SERCOULE (avend years) and intuitive for effectiveness trials, and their experience corresponded to approximately 914.3 patient-years. The conditions and cultarion for terminer with SERCOULE (avend years) and included in overlapping categories (open-table and double-blind phases of studies, inpatients and outpatients, fixed-dose and dose-firation studies, and short-term or longer-term exposer. Avenerse acrotions were assessed by collecting adverse exists, results of hypotracia carminiforms, vital signs, weights, isotracity analyses. ECIS, and results of ophthalmologic examinations, Afverse events for a significant of the proposition of the control of the proposition of the proposition of the control of the proposition of the proposition of the proposition of the proposition of individuals who experienced, at least once, a treatment-emergent adverse event of the type istent. An event was considered transverse events. The state for foreuries of adverse event of the type istent. An event was considered transverse events, and the extension of the proposition of individuals who experienced, at least once, a treatment-emergent adverse event of the type istent. An event was considered transverse events. The course or usual melancia practicities where papert characteristics are over a control course for usual melancia practicities where papert characteristics are over the control course of the circuit and the control course of the control course of the cours

2 to 12-West Explored Dorothold Chinal Tails for the Treatment of Schoodhersia and Acute Bipolar Mania (montherpy). Belgs as a White Headach, Pain, Asteria, Addination Pain, Lake Pain, Feeter Cardinassular Eurhyandia, Postular Hippolarison (Diegland: Purpholarison). Telepian (Painson). A miles and interfluent West Lake Pain, Feeter Cardinassular Eurhyandia, Postular Hippolarison (Diegland: Painson). A miles and interfluent West Lake Painson. Spring Painson, Painson,

Steven Johnson syndrome (SJS).

DRUG ABUSE AND DEPENDENCE: Centrolled Substance Class: SEROQUEL is not a controlled substance. Physical and Psychologic dependence: SEROQUEL has not been systematically studied, in animals on humans, for its other laft of abuse, to learned or physical dependence. While the clinical has did not reveal any tendency for any drug-selving behavior, these observations were not systematic and it is not possible to predict on the basis of the limit of earbier, to the central to whire IA SIS-desire drug will be revised, deverted and/or shared conserved closely for signs of misses or abuse of SEROQUEL, e.g., development of tolerance, increases in dose, drug-seesing behavior.

OVERDOSAGE: Human experience: Experience with SEROOUEL (quetiapine fumarate) in acute overdosage was limited in the clinical trial database (6 reports) with estimated doses ranging from 1200 mg to 9600 mg and natalities. In general, express dispins and symptoms were those resulting from a reaggeration of the flugs known pharmacological effects, i.e., drowsiness and sedation, tachycardia and hypotension. One case, involving an estimated overside of 9900 mg, was associated with hypotension and restrict dependence of 9900 mg, was associated with hypotension and restrict dependence of 9900 mg, was associated with hypotension and restrict and oversides of 9500 mg, were stableable and maritan an airway and ensure adequate oxygenation and verillation. Gastric lavage cfate inclusion, it option its unconsciously and administration of activated chances othered were the stable and maritanian an airway and ensure adequate oxygenation and verillation. Gastric lavage cfate inclusion, it options its unconsciously and administration of activated chances othered with a laxable whole the considered. The possibility of oblundation, setzure or dystonic rescition of the head and nick following overdose mg create a risk of assignation with induced emesis. Cardiovascular monofroing should comments insulidated comments of advantage of the stable oversible and administration of the designation and very an experience according and the stable and consideration of the stable and comments and consideration of the stable and consideration OVERDOSAGE: Human experience: Experience with SEROQUEL (quetiapine fumarate) in acute overdosage

SEROQUEL is a trademark of the AstraZeneca group of companies. © AstraZeneca 2004

Manufactured for: AstraZeneca Pharmaceuticals LP Wilmington, Delaware 19850-5437



The safety and efficacy of SEROQUEL in pediatric patients have not been established.

Patients should be periodically reassessed to determine the need for continued treatment.

Prescribing should be consistent with the need to minimize the risk of tardive dyskinesia, seizures, and orthostatic hypotension. A rare condition referred to as neuroleptic malignant syndrome (NMS) has been reported with this class of medications, including SEROQUEL.


There have been reports of diabetes mellitus and hyperglycemia-related adverse events associated with the use of atypical antipsychotics, including SEROQUEL.

The most common adverse events associated with the use of SEROQUEL were somnolence, dry mouth, dizziness, constigation, asthenia, abdominal pain, postural hypotension, pharyngitis, SGPT increase, dyspepsia, and weight gain.

In bipolar mania trials, withdrawal rates due to adverse events were similar to placebo for SEROQUEL as monotherapy (SEROQUEL 5.7%, placebo 5.1%) and adjunct therapy (SEROQUEL plus lithium or divalproex 3.6%, lithium or divalproex alone 5.9%).

References: 1. SEROQUEL® (quetiapine fumarate) Prescribing Information, Rev 01/04, AstraZeneca Pharmaceuticals LP, Wilmington, Delaware. 2. Data on file, DA-SER-13, AstraZeneca Pharmaceuticals LP, Wilmington. Delaware. 3. Data on file, DA-SER-15, AstraZeneca Pharmaceuticals LP, Wilmington, Delaware. 4. Data on file, DA-SER-14, AstraZeneca Pharmaceuticals LP, Wilmington, Delaware. 5. Data on file, DA-SER-16, AstraZeneca Pharmaceuticals LP, Wilmington, Delaware.





To prevent medication errors, write "SEROQUEL" clearly on your Rx pad. Spell "SEROQUEL" clearly over the phone.

First-line treatment

Please see Brief Summary of Prescribing Information on following page.

www.SEROQUEL.com