

Major stopover regions and migratory bottlenecks for Nearctic-Neotropical landbirds within the Neotropics: a review

NICHOLAS J. BAYLY, KENNETH V. ROSENBERG, WENDY E. EASTON,
CAMILA GÓMEZ, JAY CARLISLE, DAVID N. EWERT, ANNA DRAKE and
LAURIE GOODRICH

Summary

Nearly 300 species of landbirds, whose populations total billions, migrate between the Neotropics and North America. Many migratory populations are in steep decline, and migration is often identified as the greatest source of annual mortality. Identifying birds' needs on migration is therefore central to designing conservation actions for Nearctic-Neotropical migratory birds; yet migration through the Neotropics is a significant knowledge gap in our understanding of the full annual cycle. Here, we synthesise current knowledge of Neotropical stopover regions and migratory bottlenecks, focusing on long-distance, migratory landbirds that spend the boreal winter in South America. We make the important distinction between "true" stopover—involving multi-day refuelling stops—and rest-roost stops lasting < 24 hours, citing a growing number of studies that show individual landbirds making long stopovers in just a few strategic areas, to accumulate large energy reserves for long-distance flights. Based on an exhaustive literature search, we found few published stopover studies from the Neotropics, but combined with recent tracking studies, they describe prolonged stopovers for multiple species in the Orinoco grasslands (Llanos), the Sierra Nevada de Santa Marta (Colombia), and the Yucatan Peninsula. Bottlenecks for diurnal migrants are well described, with the narrowing Central American geography concentrating millions of migrating raptors at several points in SE Mexico, Costa Rica, Panama and the Darién. However, diurnally migrating aerial insectivores remain understudied, and determining stopover/roost sites for this steeply declining group is a priority. Despite advances in our knowledge of migration in the Neotropics, we conclude that major knowledge gaps persist. To identify stopover sites and habitats and the threats they face, we propose a targeted and collaborative research agenda at an expanded network of Neotropical sites, within the context of regional conservation planning strategies.

Introduction

Nearly 300 species of landbirds, whose combined populations represent billions of birds, migrate between the Neotropics and North America (Martin and Finch 1995, DeGraaf and Rappole 1996, Berlanga *et al.* 2010). As many as 50 of these species make an annual round trip of > 10,000 km but the number of individuals is declining year after year, especially among those species that migrate farthest (Sauer *et al.* 2014). The same is apparent in Afro-Palaearctic migrants (Sanderson *et al.* 2006). Global declines in migratory species have led to concerns that migration and the capacity of our ecosystems to maintain this process are in danger of disappearing (Wilcove and Wikelski 2008, Bauer and Hoyer 2014). For many species, the majority of their long-distance

migrations take place south of the Tropic of Cancer, yet our understanding of migration is strongly biased by studies in the temperate zone. Consequently the Neotropics continue to be described as the “black box” in our knowledge of migratory landbirds (Faaborg *et al.* 2010a).

As early as the 1980s, results from the North American Breeding Bird Survey highlighted widespread declines in Nearctic-Neotropical migratory landbirds (Terborgh 1980, Robbins *et al.* 1989), and considerable efforts have been made since to both understand and address the year-round conservation needs of this group (Keast and Morton 1980, Hagan and Johnston 1992, Holmes 2007, Faaborg *et al.* 2010a). Partners in Flight (PIF), for example, was created in 1990 to develop strategies for the conservation of migratory landbirds throughout their annual cycle, and today is the umbrella group for a diverse set of actors from across the Americas (Finch and Martin 1995, Rich *et al.* 2004, Berlanga *et al.* 2010, Rosenberg *et al.* 2016). More than 30 years of research into Neotropical migratory birds has substantially advanced our understanding of how populations may be limited by events on the breeding grounds (Holmes *et al.* 1996) or the wintering grounds (Latta and Faaborg 2002, Studds and Marra 2005, Johnson *et al.* 2006), and how effects may carry over from one season to the other (Nott *et al.* 2002, Norris *et al.* 2004, Norris and Marra 2007, Harrison *et al.* 2011, González-Prieto and Hobson 2013). Indeed, the formulation of the winter limitation hypothesis (Sherry and Holmes 1995) and the subsequent collection of empirical evidence for the limiting role of winter habitats have firmly placed Neotropical wintering grounds on the research and conservation agenda (Marra 2000, Latta and Faaborg 2002, Studds and Marra 2005). The same cannot be said, however, of the ecological needs and constraints of these same birds while migrating through the Neotropics (Faaborg *et al.* 2010a), despite the considerable distances that many long-distance migrants must traverse there.

For some bird species, migration is by far the greatest source of mortality during their annual cycle (Sillert and Holmes 2002, Newton 2006, Rockwell *et al.* 2016). For example, successive delayed arrivals or habitat degradation at even a single major stopover site can lead to significant declines, threatening the viability of populations across the Western Hemisphere (Baker *et al.* 2004). Although some en-route flexibility exists (Tottrup *et al.* 2008, Stanley *et al.* 2012), migration strategies are under strong genetic control (Gwinner 1996, Delmore and Irwin 2014), and migrants are expected to be limited in their ability to change routes if conditions change or stopover habitats are no longer available in the short term but may be able to adapt in the longer term (Sutherland 1998). Recognition of the critical importance of migration stopovers and bottlenecks has led to repeated calls for more studies and collaborations into the ecology and needs of birds during the migration periods (Hutto 1998, Heglund and Skagen 2005, Mehlman *et al.* 2005, Faaborg *et al.* 2010a, 2010b).

These appeals have led to more work within North America, especially along the USA Gulf Coast (Buler *et al.* 2007, Moore and Buler 2011, Laughlin *et al.* 2013, Cohen *et al.* 2017), the Pacific flyway (Carlisle *et al.* 2009), riparian forests in the south-west (Yong and Finch 1997, Yong *et al.* 1998) and around the Great Lakes (Dunn 2001, 2002, Ewert *et al.* 2011). For many migratory species, however, the majority of their migratory route lies south of these well-studied areas. For example, millions of migrating landbirds converge each year on the northern Neotropics (La Sorte *et al.* 2016), and species whose breeding range may stretch over 4,000 km from east to west concentrate into areas spanning less than 100 km as a result of the funnel-shaped geography of Central America (Bildstein 2004). Yet, despite this obvious importance, migration studies of Nearctic-Neotropical landbirds south of the United States of America (USA) and Canada remain rare, limiting our ability to design effective conservation strategies that address their needs throughout the annual cycle.

Nonetheless, the last few decades have seen studies from Mexico's Gulf Coast and the Yucatan Peninsula (Winker 1995, Deppe and Rotenberry 2005, Johnson and Winker 2008, Bayly and Gómez 2011, Shaw and Winker 2011), and from northern Colombia (Colorado 2010, Bayly *et al.* 2012a, 2013, Gómez *et al.* 2013, 2015), in some cases demonstrating the critical importance of stopover sites within the Neotropics (Bayly *et al.* 2012a, 2013). Enormous gaps remain, however, and in particular our knowledge of how nocturnally migrating birds use the region south of the

Yucatan Peninsula through Central America to northern South America; large Caribbean Islands; and especially regions within South America, is poor. In addition, for over 30 species that winter primarily in South America, much confusion persists around which regions are used during migration because of imprecisely mapped winter distributions and the difficulty of distinguishing migrating from wintering individuals (Renssen 2001). Some of these gaps are being partially filled as a result of innovative lightweight tracking technologies and their application to a growing number of Neotropical migratory landbirds (McKinnon *et al.* 2013a, Taylor *et al.* 2017). The migration of raptors is better understood, with multiple studies detailing the main migration routes (Martell *et al.* 2001, Bildstein 2004, Porras-Peñaranda *et al.* 2004, Colorado *et al.* 2006, Ruelas Inzunza *et al.* 2010a, Kochert *et al.* 2011, Bayly *et al.* 2014).

While general patterns of migration speed and routes have been shown to differ between spring and autumn migrations for many species (Gómez *et al.* 2013, La Sorte *et al.* 2014a, 2014b, Hobson and Kardynal 2015), detailed seasonal comparisons for most species and regions are lacking. In addition, the strategies and needs of diurnal and nocturnal migrating species are expected to differ markedly, making generalisations across species groups difficult. Given the critical importance of migration to annual survival and fitness (Sillert and Holmes 2002, Finch *et al.* 2014) and its ecological significance for long-term population viability, our lack of knowledge of stopover regions and migratory bottlenecks within the Neotropics poses a serious constraint on our ability to address declines in Nearctic-Neotropical migrants (Faaborg *et al.* 2010a, Sheehy *et al.* 2010).

In this review we synthesise current knowledge of stopover regions and migratory bottlenecks within the Neotropics, focusing on diurnal and nocturnally migrating landbirds. Our primary objective is to stimulate research in regions and for species that lack even baseline data. We outline potential threats at known and suspected stopover regions, and present an emerging picture of the migration strategies of long-distance migratory birds in the Neotropics. We also propose consistent terminology that, for example, can make the distinction between “true” stopover—where birds engage in multi-day stops with significant refuelling—and rest-roost stops lasting one day or less with no or limited refuelling. Then, based on a conceptual framework for understanding the needs of migratory birds, we propose a targeted and collaborative research agenda at an expanded network of sites across the Neotropics within the context of new regional conservation planning strategies (e.g. Rosenberg *et al.* 2016).

The needs of migrating landbirds in the Neotropics

To successfully migrate between their breeding and over-wintering grounds, where *over-wintering* is defined as the stationary period/s of the non-breeding season during the boreal winter (Table 1), Nearctic-Neotropical migrants depend on a series of sites along the length of their migratory route. These sites provide critical resources such as the fuel for migratory flights, safe roosting sites, and refuges during unfavourable climatic conditions (Mehlman *et al.* 2005, Newton 2008). A successful migration also depends on birds avoiding in-flight hazards such as collisions with human infrastructure (Longcore *et al.* 2013) and persecution by humans (Newton 2006). Migrating landbirds stop at multiple sites across regions, yet not all sites are of equal importance for determining the speed of migration or survival (Weber *et al.* 1998). Indeed, diurnal migrants typically stop on a nightly basis between daytime flights, while nocturnal migrants generally stop on a daily basis when migrating overland (Newton 2008). As a consequence, migrants can be encountered at numerous sites along the length of their migratory routes, making it difficult to differentiate between sites used to rest between flights and those used to accumulate energy reserves (Warnock 2010).

The lack of distinction between these two very different types of “stopover sites” has given rise to a general perception that most migratory landbirds use numerous stopover sites during any one migration (McKinnon *et al.* 2013a) and that population level movement across space is continuous. Because of this perception, research, conservation, and management activities

Table 1. Stopover and migration terminology adopted throughout this review and its equivalence to the categories proposed by Mehlman *et al.* (2005), which was used as the foundation.

| Type of site or region | Description | Equivalent Mehlman et al. |
|------------------------|--|-------------------------------------|
| Passage | Geographical points or corridors where birds pass in continuous flight during migration | N/A |
| Bottleneck | Passage points where geography constrains migration, leading to significant concentrations of populations that could elevate threats or risks. | N/A |
| Rest-Roost | Places associated with resting or roosting behaviours between successive migratory flights that do not involve significant refuelling (likely applies to the majority of sites used on migration where refuelling is equivalent to <5% of lean body mass). | Similar to <i>convenience store</i> |
| Fire escape | Places where migrants only make landfall when they encounter adverse conditions that prohibit onward flights; sites provide vital shelter for survival but are rarely used for refuelling. | <i>Fire escapes</i> |
| Stopover | Behaviour and places associated with multi-day stops giving rise to significant refuelling (>5% of lean body mass) during migration. | <i>Full service hotel</i> |
| Staging | Behaviours and places associated with prolonged stopover not entirely dedicated to refuelling, such as where large numbers of birds gather, often building in numbers until favourable cues promote subsequent migration. | <i>Full service hotel</i> |
| Moult-migration | Behaviour or places where a significant portion of a species' population pauses on migration to undergo flight feather moult before continuing on migration. | N/A |
| Pre-migratory | Behaviour or places where significant fuelling occurs immediately following breeding or over-wintering periods that may or may not involve short local movements (<100 km). | <i>Full service hotel</i> |
| Over-wintering | Behaviour or places associated with prolonged (>1 month) stationary non-breeding periods during the boreal winter season, usually between autumn and spring migration periods. Fuelling does not take place during this period. Given the prevalence of multiple over-wintering sites for some species, may blur with pre-migratory "staging". | N/A |

for birds on migration have often focused on sites where large numbers of birds are present (Moore *et al.* 1990, Simons *et al.* 2000, Dunn 2001), without considering the ecological function of those sites. Recent studies on a range of species have challenged this paradigm, showing that in reality, individual landbirds often make long stopovers at just three or four key areas along the migratory route (Figure 1) (Heckscher *et al.* 2011, Kochert *et al.* 2011, Delmore *et al.* 2012, Fraser *et al.* 2012, Stanley *et al.* 2012, Callo *et al.* 2013, McKinnon *et al.* 2013a, Renfrew *et al.* 2013), at which they accumulate large energy reserves (Figure 2) (Bayly *et al.* 2012a, 2013). These findings highlight an urgent need to identify major stopover regions (defined here as areas varying in size from 100 km² to 50,000 km² with relatively uniform environmental conditions and offering a certain set of resources) and assess the needs of birds within them. This need is magnified in the Neotropics where the funnel-shaped geography of Mexico and Central America acts as a bottleneck, concentrating millions of migratory landbirds (Porrás-Peñaranda *et al.* 2004, Batista *et al.* 2005, Ruelas Inzunza *et al.* 2010a, La Sorte *et al.* 2016). This also highlights a need to revise the current stopover terminology to better define the contribution of individual sites to migration strategies, and to this end we present an updated terminology in Table 1 that we utilize throughout the review.

One of the major factors driving the organisation, speed and, ultimately, the success of migration is the quality of resources at stopover sites and their impact on the rate of fuel deposition (Hedenström 2008). Indeed, with time-minimisation emerging as the main currency shaping the evolution of migration strategies (Alerstam 2011), migratory birds are expected to seek out sites that maximize their fuel deposition rate and, accordingly, their speed of migration (Buler *et al.* 2007, La Sorte *et al.* 2014a). It follows that one of the fundamental needs of migratory birds is therefore the availability of high quality habitats (those supporting high fuel deposition rates), in specific regions along their migratory route (Weber *et al.* 1999, Bayly *et al.* 2016). These habitats




Figure 1. Spring migration strategy (northward) and stopover site use of Red-eyed Vireos from a breeding site in the north-eastern United States as revealed by geolocators recovered from nine individuals (adapted from Callo *et al.* 2013). Eight of the nine individuals appeared to make a rest-roost stop close to the tip of the Yucatan peninsula before crossing the Gulf of Mexico.




Figure 2. Individual **A.** Veery and **B.** Grey-cheeked Thrush recaptured on more than one occasion during an autumn and spring stopover respectively, in the Sierra Nevada de Santa Marta in northern Colombia. Birds increased significantly in body mass, providing evidence for extensive fuelling in this region (*taken from Bayly *et al.* 2012a, 2013*).

must not only sustain high rates of fuel deposition but also offer low to moderate predation risk, as migratory birds may avoid sites with elevated densities of predators or trade-off their foraging rate against predator surveillance (Lindström 1990, Schmaljohann and Dierschke 2004, Pomeroy 2006, Warnock 2010). Recent work on *Catharus* thrushes in northern Colombia has shown that the energy reserves acquired at key stopovers may enable birds to cover 30% or more of their total migratory distance (Bayly *et al.* 2012a, Bayly *et al.* 2013, Gómez *et al.* 2017), again highlighting the enormous influence that individual stopover sites can have on a migratory journey (Figure 2).

Additional sites used by migrating birds between major stopover regions (which we will refer to as *rest-roost stops* hereafter) are believed to have far less impact on the speed and organisation of migration (Alerstam 2011); nevertheless they have the potential to affect survival (Newton 2006). Energetic requirements at rest-roost stops are generally low, as individuals may only be resting between successive nocturnal or diurnal flights, although the ability to top up reserves may constitute a key element of the migratory strategy in certain species (Robson *et al.* 2001, Delingat *et al.* 2006). The primary function of rest-roost stops is therefore to provide safe roosting conditions and access to resources such as water. Sites used as rest-roost stops may assume a greater importance when situated close to ecological barriers, such as the Gulf of Mexico or the Caribbean Sea (Moore *et al.* 1990). Under optimal conditions, birds typically overfly coastal habitats either side of these barriers (Lowery 1945, Moore *et al.* 1990) but when they encounter unfavourable conditions such as rain or strong headwinds, these habitats may temporarily support high concentrations of birds (Simons *et al.* 2000). Habitats used in such emergencies, often referred to as “fire escapes”, can also be important away from ecological barriers (Mehlman *et al.* 2005). For example, passing tropical storms in Central America occasionally ground migrating raptors for days at sites where normally they would not stop (Bildstein 2004) and possibly delay subsequent returns to the breeding grounds.

Just as stopover sites used for migratory fuelling affect the outcome of migration, the quality of sites occupied prior to the onset of migration can also have a fundamental impact on migration strategies (Bearhop *et al.* 2004, Norris and Marra 2007). For example, many species undergo pre-migratory fuelling on or near their Neotropical overwintering grounds, and the energy reserves

gained there may enable birds to cover a significant proportion of their northbound migration (Stutchbury *et al.* 2009, Heckscher *et al.* 2011, DeLuca *et al.* 2015). Further, the quality of overwintering habitats can influence the timing of migration, both in terms of initiation and arrival on the breeding grounds, with carry-over effects on territory acquisition and breeding performance (Norris *et al.* 2004, Studds and Marra 2005, Norris and Marra 2007, Guillemain *et al.* 2008). Overwintering habitats therefore likely constitute an essential element in many birds' migration strategies but, unlike stopover habitats, they tend not to support high concentrations of birds. As a consequence, the loss of stopover habitat along migration routes may have more severe population consequences than the loss of overwinter habitat used for pre-migratory fuelling (Baker *et al.* 2004, Newton 2004).

While moult rarely overlaps with migration, this is not true in all species and stopover or pre-migratory fuelling sites may also be used to meet the energetic requirements of feather replacement. Moulting-migration, whereby birds undergo a complete moult along their migration route, apparently "pausing" their migration to do so, occurs in the monsoon region of south-western USA and north-western Mexico in several species that breed in western North America (Leu and Thompson 2002, Rohwer *et al.* 2005). Energetic demands at moulting-migration sites are presumably higher than at typical stopover sites, yet the extent to which this phenomenon occurs elsewhere in the Neotropics is unknown. Similarly, pre-alternate moults that give rise to breeding plumages also likely occur at sites used for pre-migratory fuelling, but specific strategies or needs of birds during this critical phase of the life cycle have rarely been studied. In at least one long-distance migrant, the Bobolink *Dolichonyx oryzivorus*, areas used in South America for pre-alternate moult are part of a sequence of stationary non-breeding sites used by this species, blurring the concepts of overwintering, stopover, or staging sites (Renfrew *et al.* 2013).

Current knowledge of stopover regions and migratory bottlenecks for landbirds in the Neotropics

In this section we discuss our current knowledge of diurnal migrants and nocturnal migrants separately, given the differing needs of these two groups; e.g. diurnal migrants both migrate and feed by day and sleep at night, whereas nocturnal migrants migrate at night and feed and/or rest by day. This may not hold true, however, when birds make long over-water flights that take more than 12 hours to complete. Secondly, we discuss findings from geolocators and other tracking technologies independently of on-the-ground field studies, in light of the qualitative differences in the data they provide. Finally, although we discuss migrants from both eastern and western North America, we concentrate on the preponderance of eastern breeding species whose migration routes take them through Central and northern South America and across barriers such as the Gulf of Mexico and Caribbean Sea. In contrast, many species from western North America overwinter in the northern extent of the Neotropics (Pacific slope of Mexico and northern Central America; Kelly and Hutto 2005) and key stopover regions linked with survival and productivity occur outside of our region of interest (LaManna *et al.* 2012, Drake *et al.* 2014). Nonetheless, a handful of western species migrate to South America and likely also rely on Neotropical stopovers.

The studies presented in this section were compiled following an ongoing and exhaustive literature search, employing the Google Scholar and Web of Science search engines. We entered the following key words singly or in combination: fuelling, fuel load, fuel deposition rate, geocator, migration, migratory birds, migration strategy, stopover; in combination with geographical locations such as Neotropics, Central America, Costa Rica, Colombia, etc. In addition, the literature cited in relevant publications was surveyed for appropriate references. In compiling and summarising information, we specifically looked for papers in which evidence for stopovers such as stopover duration or changes in fuel loads were presented. Finally, we used the general Google search engine to discover technical reports published on the web.

Raptors and other diurnal migrants

Among the diurnally migrating species that use terrestrial habitats, perhaps the best-known group is the raptors (in which we include falcons and New World Vultures). In addition to long-term monitoring programmes at a variety of sites between Veracruz, Mexico (Ruelas Inzunza *et al.* 2010a), and the Canal Zone in Panama (Porrás-Peñaranda *et al.* 2004, Batista *et al.* 2005), the use of satellite transmitters has revealed considerable detail with respect to routes and stopovers (Fuller *et al.* 1998, Martell *et al.* 2001, Kochert *et al.* 2011).

More than 30 raptor species have been recorded migrating through Mesoamerica but the majority of the 4 to 6 million raptors that migrate to or through the Neotropics on an annual basis belong to just four species: Turkey Vulture *Cathartes aura* (2 million), Broad-winged Hawk *Buteo platypterus* (2.1 million), Swainson's Hawk *Buteo swainsonii* (1.2 million), and Mississippi Kite *Ictinia mississippiensis* (0.2 million) (Bildstein 2004, Ruelas Inzunza *et al.* 2010b). Other less abundant but visible species along the flyway include Osprey *Pandion haliaetus*, Peregrine Falcon *Falco peregrinus*, and Swallow-tailed Kite *Elanoides forficatus*. Acting like a natural bottleneck, the narrowing geography of Central America concentrates migrating raptors at several points. The most well studied bottlenecks include the narrow gap between the Sierra Madre Oriental and the Gulf of Mexico in Veracruz, Mexico (Ruelas Inzunza *et al.* 2009), the corridor between the Cordillera de Talamanca and the Caribbean Sea in Costa Rica and western Panama (Porrás-Peñaranda *et al.* 2004) and the Isthmus of Panama (Batista *et al.* 2005). Nevertheless, large numbers of migrating raptors can be seen at several other points along the route including the Darién of Panama and Colombia (Bayly *et al.* 2014), and further study will likely reveal a concentrated passage along the Pacific coasts of Guatemala, El Salvador and Nicaragua (Bildstein 2004).

The routes taken by raptors between Mexico and South America are well-defined, with the main flyway dividing in two in southern Mexico, after birds have passed through coastal Veracruz (Bildstein 2004). The majority of birds cross to the Pacific coast at this point, following it down through Guatemala, El Salvador and Nicaragua until crossing to the Caribbean lowlands in southern Nicaragua and northern Costa Rica (Porrás-Peñaranda *et al.* 2004, Kochert *et al.* 2011); however, Broad-winged Hawk, for example, may follow the Caribbean coast throughout. On entering Panama, many individuals cross again to the Pacific slope, although many others continue down the Caribbean coast, entering South America via the Darién. In South America routes are not well studied (Colorado *et al.* 2006), but satellite tracks suggest that while Swainson's Hawks continue on an almost direct southerly route (Kochert *et al.* 2011), Turkey Vultures head east-south-east towards the Llanos of Venezuela and Colombia. Many Osprey, Peregrine Falcon and Merlin *Falco columbarius* do not join the main flight through Mesoamerica until its lower portion, having left North America via the Florida peninsula and migrating over-sea via Cuba (Fuller *et al.* 1998, Martell *et al.* 2001, Rodríguez-Santana *et al.* 2014). Western Cuba has also been shown to be a major migratory route for USA-breeding Swallow-tailed Kites (Zimmerman and Meyer 2004).

Migrating raptors generally pass along the routes described above rapidly, rarely pausing to top up fuel reserves (Smith *et al.* 1986, Fuller *et al.* 1998, Martell *et al.* 2001, Kochert *et al.* 2011); however, a number of stopover regions have been identified. For example, while autumn migrating Swainson's Hawks appear to accumulate much of their energy requirements at stopovers in the southern USA, birds do make occasional stops in south-eastern Mexico, western Guatemala and Nicaragua, north-western Colombia and in the Colombian Llanos (Kochert *et al.* 2011). In contrast, the spring migration of Swainson's Hawks does not appear to involve Neotropical stopovers (Smith *et al.* 1986, Kochert *et al.* 2011). Swallow-tailed Kites make a prolonged stopover on the Yucatan peninsula during autumn migration (Zimmerman and Meyer 2004), seemingly to recover after an over-sea crossing from Florida or Cuba. The lack of information on other species and the general notion that most raptors can complete their migration through the Mesoamerican corridor without stopping to fuel (Smith *et al.* 1986), imply that their principal need may be safe roosting sites. However, anecdotal evidence aside (Hicks *et al.* 1966, Bildstein and Saborio 2000, Ruelas Inzunza *et al.* 2009), there is very little information regarding the use of roosting sites by

migrating raptors. Nonetheless, the highly concentrated flyway used by most raptors suggests that they likely have as yet unidentified needs, which in part may be dictated by their diet which impedes the *en masse* use of stopover sites.

Besides raptors, very little has been published on other diurnally migrating landbirds, although these species are routinely encountered during raptor migration monitoring. Hirundines are a highly abundant group of diurnal migrants and seasonal counts of Cliff *Petrochelidon pyrrhonota*, Barn *Hirundo rustica* and Bank *Riparia riparia* Swallows at bottlenecks in Veracruz and the Colombian Darién involve hundreds of thousands of birds (Winkler 2006, Bayly *et al.* 2014). Little is known, however, about stopover behaviour in these species. Given the declining status of many aerial insectivores (Table 2) (Nebel *et al.* 2010, Rosenberg *et al.* 2016), determining stopover and major roost sites for this group is a priority, especially if they adopt staging behaviours similar to those seen in Purple Martin *Progne subis* and Tree Swallow *Tachycineta bicolor* (Fraser *et al.* 2013, Laughlin *et al.* 2013), where birds remain at sites longer than would appear necessary for fuelling purposes. Many aerial insectivores migrate as far as the southern cone of South America and likely have complex migration strategies and differential strategies between populations of a single species (Table 2). Other abundant yet rarely reported diurnal migrants are even less well known, and aside from counts at concentration points in the Darién (Bayly *et al.* 2014), little is known of migration or even overwintering grounds of species such as Chimney Swift *Chaetura pelagica*, Black Swift *Cypseloides niger*, Common Nighthawk *Chordeiles minor*, Eastern Kingbird *Tyrannus tyrannus* and Dickcissel *Spiza americana*.

Nocturnal migrants

The vast majority of Nearctic-Neotropical migratory landbirds (e.g. warblers, thrushes, tanagers, cuckoos) migrate solely or primarily at night (DeGraaf and Rappole 1996, Alerstam 2009), and many traverse vast stretches of the Caribbean basin or Central and northern South America to reach wintering grounds in the Andes, the Amazon Basin, or farther south still. Despite the importance of this extensive geography to annual survival, many Neotropical regions are still completely unexplored in terms of where, when and why migratory landbirds stop (Faaborg *et al.* 2010b).

Table 2. Over-wintering destination and population status of diurnally migrating landbirds that winter in South America. Season counts are based on the highest single autumn counts from a single site in the Darién of Colombia (see Figure 4, region 3) reported in Bayly *et al.* (2014) or on counts from the same study site in 2014 and 2015. Three species (marked N/A) are rarely recorded at this watchsite presumably because their migration route takes them over the Caribbean Sea or the Pacific Ocean.

| Species | Over-wintering Destination | Population Status ¹ | Season high counts (Year) |
|---|-----------------------------------|--------------------------------|---------------------------|
| ² Common Nighthawk <i>Chordeiles minor</i> | S. South America ³ | -58% | 21,922 (2015) |
| Chimney Swift <i>Chaetura pelagica</i> | W. Amazon Basin | -67% | 6,922 (2014) |
| Black Swift <i>Cypseloides niger</i> | W. Amazon Basin | -94% | N/A |
| Eastern Kingbird <i>Tyrannus tyrannus</i> | W. Amazon Basin | -38% | 243,669 (2014) |
| Purple Martin <i>Progne subis</i> | S. American Lowlands ⁴ | -23% | N/A |
| Bank Swallow <i>Riparia riparia</i> | S. American Lowlands ⁴ | -89% | 22,898 (2011) |
| Barn Swallow <i>Hirundo rustica</i> | S. American Lowlands ⁴ | -38% | 87,655 (2014) |
| Cliff Swallow <i>Petrochelidon pyrrhonota</i> | S. South America ³ | +37% | 158,413 (2014) |
| Bobolink <i>Dolichonyx oryzivorus</i> | S. South America ³ | -60% | N/A |
| Dickcissel <i>Spiza americana</i> | N. South America | -14% | 30,268 (2015) |

¹Population loss from 1970-2014 based on Rosenberg *et al.* 2016.

²Nighthawks regularly initiate and terminate flights during daylight hours but may migrate primarily at night.

³Includes regions primarily south and east of the Amazon Basin.

⁴Includes various lowland areas including Amazon Basin and regions to south and east.

In this section, we describe current knowledge of Neotropical stopover sites for nocturnal migrants within six broad geographic regions (Fig. 3): (1) south-eastern Mexico and the Yucatan peninsula; (2) northern Central America including Honduras, Nicaragua, and Guatemala; (3) Costa Rica and Panama; (4) northern Colombia; (5) the Caribbean islands; and (6) western Mexico. Other regions in the Neotropics are no doubt used for stopover but they remain largely unstudied.

South-eastern Mexico and the Yucatan Peninsula — Perhaps the best studied region, with several studies from Veracruz (Winker 1995, Martínez Leyva *et al.* 2005, Shaw and Winker 2011), the northern tip of the Yucatan (Deppe and Rotenberry 2005), north-eastern Belize (Bayly and Gómez 2011, Gómez and Bayly 2011), and southern Belize (Johnson and Winker 2008). During autumn migration, species richness (≈ 70 species) and abundance of migratory landbirds is high and it is evident that the coastal geography of the region concentrates individuals of many species that cross the Gulf of Mexico or migrate around its western edge (Winker 1995, Bayly and Gómez 2011). However, mist-netting has revealed that individuals of most species carry moderate fuel reserves and therefore have little need to refuel at stopover sites (Winker 1995, Johnson and Winker 2008, Bayly and Gómez 2011). Low recapture probabilities and short between-capture durations also imply that most birds were making rest-roost stops between nocturnal flights (Bayly and Gómez 2011).

Winker (1995) and Johnson and Winker (2008) provide evidence that birds may be able to top up their reserves during these daytime stops but provide no evidence that birds stopover for longer. Together these findings suggest that most migrants (19 out of 20 transient species in north-east Belize, for example) passing through the region may accumulate sufficient energy reserves north of the Gulf of Mexico to cross both the Gulf and the Yucatan Peninsula (Bayly and Gómez 2011). Not all species or individuals follow this pattern and Willow Flycatchers *Empidonax traillii*, for example, probably make a short refuelling stop before continuing their migration (Gómez and Bayly 2011).

Spring migration sees a reduction in species richness in the region, and many transient species on route from South America, appear to be carrying sufficient reserves on arrival to reach North America without refuelling (Bayly and Gómez 2011). For a smaller number of Central American wintering species, fuel loads, recapture rates and fuel deposition rates point to a stopover both on




Figure 3. The Neotropical region lying between the Tropic of Cancer and the Tropic of Capricorn. Major regions discussed in the section on nocturnal migrants and referred to throughout the text are outlined in shades of grey and named following the text.

the Yucatan Peninsula and in south-eastern Mexico, including species of concern such as Wood Thrush *Hylocichla mustelina* and Kentucky Warbler *Geothlypis formosa* (Rogers and Odum 1966, Bayly and Gómez 2011, Shaw and Winker 2011).

Northern Central America — There are no published studies of autumn stopover site use in northern Central America, despite the potential importance of this region for a large number of species that cross the Gulf of Mexico (Cohen *et al.* 2017). A spring study focused on Cerulean Warblers *Setophaga cerulea* in the mountains of northern Honduras, eastern Guatemala and southern Belize, highlighted the importance of this area for this species of conservation concern (Welton *et al.* 2012). Cerulean Warblers are rarely recorded anywhere along their migration route between North American breeding grounds and Andean wintering grounds, but Welton *et al.* (2012) documented the presence of birds during a two-week period in early April, largely on Caribbean-facing mountains in Honduras and Guatemala. Several other warbler species, including the 'Near Threatened' Golden-winged Warbler *Vermivora chrysoptera*, were also recorded in this study. Given its geographic position and intriguing unpublished records submitted to the online public database eBird (e.g. 400 Bay-breasted Warblers, *Setophaga castanea*, counted at a single location in Nicaragua in April) (Sullivan *et al.* 2014), northern Central America will likely prove to be a critical stopover region for many species.

Costa Rica and Panama — Stopover studies from Costa Rica and Panama are also surprisingly rare, and aside from those dating back 50 years (Rogers and Odum 1966) and a recent study on Swainson's Thrush *Catharus ustulatus* (Wilson *et al.* 2008), we could find no reference to the stopover behaviour of Nearctic-Neotropical migrants in this region. There is, however, a long-term migration monitoring station at Tortuguero, Costa Rica (Elizondo-Camacho and Ralph 2012), and the data collected there would no doubt shed light on the use of the Caribbean lowlands for stopovers, beyond the habitat selection inferences thus far reported (Wolfe *et al.* 2014). Further, unpublished records in eBird and elsewhere suggest that a variety of species (e.g. Cerulean Warbler) may depend on montane forest habitats in Costa Rica and Panama, especially in early autumn. Studies of autumn migrants on the Caribbean coast of Panama found a mixture of exhausted birds and individuals with sufficient reserves to continue their migration well into South America (Rogers 1965, Rogers and Odum 1966). There was some evidence that fuel-depleted birds stopped over in Panama to rebuild their energy reserves (Galindo *et al.* 1963). In a spring study of Swainson's Thrushes on the Pacific coast of Costa Rica, there was no evidence of birds making prolonged stopovers (Wilson *et al.* 2008).

Northern Colombia — In contrast to the above regions, a growing body of evidence on stopover site use in northern Colombia is emerging, with studies on *Empidonax* flycatchers in the Darién (Colorado 2010) and on *Catharus* thrushes, vireos, and warblers in the Sierra Nevada de Santa Marta (SNSM) and the Darién revealing new insights on the importance of this region to migrants (Gómez *et al.* 2013, 2014). During autumn, *Empidonax* flycatchers stopping in the Darién did not appear to build energy reserves (Colorado 2010), whereas three species of *Catharus* thrushes exhibited a mixed strategy, with some individuals passing through rapidly and others making multi-day stopovers to refuel (Gomez *et al.* 2014). In contrast, migrants passing through the SNSM frequently made prolonged stopovers in both autumn and spring. For example, Veery *Catharus fuscescens* on stopover in the SNSM stayed for an average of nine days in autumn, during which time they increased their energy reserves by around 30% (Figure 2) and were subsequently capable of flights of up to 2,000 km towards their South American overwintering grounds (Bayly *et al.* 2012a). A similar situation was found for Grey-cheeked Thrush *Catharus minimus* in spring, with birds stopping in the SNSM for up to two weeks and storing sufficient energy to reach North America (>2,500 km) without refuelling (Bayly *et al.* 2013). In a third species, the Tennessee Warbler (*Oreothlypis peregrina*), long spring stopovers in the SNSM resulted in

smaller fuel loads compared to Grey-checked Thrushes, but they were still sufficient for flights to northern Central America and, in some individuals, to the southern USA (Bayly *et al.* 2016).

These recent results from studies in northern Colombia suggest that migratory landbirds are capable of adopting a migratory strategy similar to that of several shorebirds (Gómez *et al.* 2017); i.e. using a small number of sites to store large fuel reserves for long-distance flights (Piersma 1987, Gill *et al.* 2009, Klaassen *et al.* 2011). The strategy of species like the Grey-cheeked Thrush also has similarities with those adopted by migrating raptors, which pass over Mesoamerica without the need for refuelling stops (Smith *et al.* 1986, Kochert *et al.* 2011). These strategies do not appear to be restricted to the species studied in Colombia to date, as the results from geolocators described below demonstrate. Another key lesson from studies in Colombia is that migratory landbirds are far more abundant in certain stopover habitats (e.g. pre-montane forest) compared to others and that different habitats support different fuelling opportunities and rates, which can make the difference between individuals successfully crossing a water barrier or needing to take a longer route or make additional stopovers (Gómez *et al.* 2013, 2015, Bayly *et al.* 2016).

Caribbean islands — Despite many long-term studies of overwintering migrants in the West Indies, studies of stopover behaviour in the Caribbean are rare. General texts on Caribbean birds indicate that most species migrating to South America are rarely encountered on Caribbean islands and that most records are thought to be weather-related (Garrido and Kirkconnell 2000, Raffaele *et al.* 2003, Latta *et al.* 2006); thus the optimal strategy seems to be to overfly the Caribbean, rather than stopping to refuel on islands. This hypothesis is supported by a study of Blackpoll Warbler *Setophaga striata* in the Dominican Republic, where birds primarily undertook daytime rest-roost stops, often in association with unfavourable weather (Latta and Brown 1999). Geolocators suggest a similar role for Caribbean islands (see below), but further study may reveal stopover areas, particularly on large islands such as Cuba where information is scarce.

Western Mexico — Few studies of landbird migration stopover have occurred in western Mexico, yet the diversity of habitats, ranging from coastal lowlands to high mountains, combined with the large number of species migrating through or overwintering in Mexico, strongly suggests its importance as a stopover region, especially for species breeding in the western USA and Canada (Howell and Webb 1995). Recent studies of moult migration in north-western Mexico, in which individuals of many species undergo an extended stopover to complete feather moult during and after the summer monsoon season and then continue to overwintering areas (Leu and Thompson 2002, Rohwer *et al.* 2005, Pyle *et al.* 2009), is one key example showing the importance of western Mexico to a critical life history stage.

Findings from geolocators and other tracking technologies

Geolocators have revolutionised our ability to track small landbirds during migration, revealing a wealth of information that can be used to identify migratory routes and stopover regions. Nonetheless, due to the relatively low precision of light-level geolocators (typically ± 100 km but up to 400 km in the tropics) (McKinnon *et al.* 2013a, 2013b), it is only by combining findings from geolocators with detailed on-the-ground field studies that habitat use and quality within broad stopover regions can be assessed. Indeed, the array of habitats present within polygons or confidence ellipses identified by geolocators can be extremely large. For example, in northern Colombia, where geolocators have identified a general region used by Veery for autumn stopover (Heckscher *et al.* 2011, Hobson and Kardynal 2015), it is possible to travel from snow-capped peaks, through humid montane forest, to xeric scrub in less than 100 km. Only through detailed mark-recapture studies within that region has the high value of foothill and pre-montane forest been demonstrated for Veery and other Nearctic-Neotropical migrants (Bayly *et al.* 2012a, 2016).

Geolocators have revealed a number of unexpected patterns in the migratory strategies of small landbirds, including faster than expected travel speeds and a general pattern of birds making

prolonged stopovers in a small number of regions (McKinnon *et al.* 2013a). For example, eastern-breeding Veery were found to make multi-day stopovers in three main regions, the south-eastern USA, northern Colombia, and the northern Amazon, when on route from a Delaware breeding site to overwintering sites in the southern Amazon basin (Heckscher *et al.* 2011). Remarkably, western-breeding Veery also converge on the same northern Colombia stopover region in autumn (Hobson and Kardynal 2015), as results from an analysis of stable isotopes in feathers had predicted (González-Prieto *et al.* 2011). Bobolink use fewer regions still, with many birds making one long stopover in the Llanos (grasslands) of Venezuela and Colombia, while on route between North America and their southern South American overwintering grounds (Renfrew *et al.* 2013).

The pattern of long stopovers at a relatively small number of sites is emerging from landbird studies in both the Americas and Europe (Stanley *et al.* 2012, Kristensen *et al.* 2013, Lemke *et al.* 2013, DeLuca *et al.* 2015). This strategy may be used primarily by nocturnally migrating species, as aerial insectivores (Beason *et al.* 2012) and other diurnal migrant birds appear to behave differently (Jahn *et al.* 2013): although Fraser *et al.* (2013) documented prolonged stopovers in Purple Martin. Presumably, the ability of diurnal migrants to feed while actively migrating explains part of this difference. Aside from revealing the widespread occurrence of long stopovers, geolocators also show why landbirds were previously considered to use numerous sites/regions during migration. Indeed, although birds move rapidly between these major stopover regions, often at rates of 500 km/day (Callo *et al.* 2013, Fraser *et al.* 2013), flights are interrupted at a series of sites along the route, which most likely act as diurnal or nocturnal rest-roost stops in order to avoid migrating at unfavourable times of day (Stutchbury *et al.* 2009, Delmore *et al.* 2012).

There are still too few published geocator studies to draw conclusions about the Neotropical stopover regions used by the wider community of migratory landbirds, or by species with shared wintering grounds, or even those from different breeding regions (e.g. eastern vs. western North America). Nevertheless, several regions are now known to be important for two or more species (Table 3). For example, Purple Martin (Fraser *et al.* 2013), Swainson's Thrush (Delmore *et al.* 2012), and to a lesser extent Wood Thrush (Stutchbury *et al.* 2009) make an extended autumn stopover on the Yucatan Peninsula, joining the list of species found to make short stops there by banding studies (see above). Northern Colombia hosts extended stopovers by at least three species, Veery (autumn), Red-eyed Vireo *Vireo olivaceus* (spring), and Swainson's Thrush (autumn), adding further evidence for this region's importance to South American wintering species (Heckscher *et al.* 2011, Delmore *et al.* 2012, Callo *et al.* 2013). Geolocators further indicate that both Red-eyed Vireo (spring) and Swainson's Thrush (autumn and spring) also make long stopovers in northern Central America, confirming our prediction that this region should figure prominently in migration strategies. Finally, it is worth emphasizing the critical importance of the Llanos of Venezuela and Colombia as an autumn and spring stopover site for Bobolink (Renfrew *et al.* 2013).

GPS tags and coded nano-tags (radio transmitters) also hold promise for determining the precise location of new stopover sites. Nano-tags, in particular, are an excellent tool for studying stopover behaviour and the outcome of subsequent migratory flights (Taylor *et al.* 2011, 2017), such as the confirmation of non-stop flights >3,000 km by Grey-cheeked Thrush on leaving a stopover site in Northern Colombia (Gómez *et al.* 2017).

Emerging picture of stopover regions in the Neotropics

The general pattern emerging from both on-the-ground field studies and the deployment of geolocators is that long-distance migration strategies in many small landbirds are actually similar to those of certain shorebirds (Atkinson *et al.* 2007, Gill *et al.* 2009, Lindström *et al.* 2016), involving long flights and prolonged multi-day stopovers at a relatively small number of sites. We are just beginning to understand the generality of this pattern, as well as which regions are of greatest importance to the largest number of migratory birds.

Table 3. Evidence for major stopover regions and bottlenecks for migratory landbirds in the Neotropics. Numbers following region names relate to Fig. 3 & 4.

| Region (see Fig. 3) | Species | Season | Method | Evidence for | Reference |
|---|---|--------|----------------------------|---|---|
| Yucatan/SE Mexico (8 & 9) | | | | | |
| Veracruz | Raptors, diurnal landbirds | Autumn | Observations | Major bottleneck for several raptor species | Ruelas I. <i>et al.</i> 2010a |
| Yucatan Peninsula | Purple Martin | Autumn | Geolocators | Stopover: mean stopover duration 16 days | Fraser <i>et al.</i> 2013 |
| SE Mexico | Kentucky Warbler & others | Spring | Mist-netting | Mass gains and recapture rate suggest stopover in Kentucky & Hooded Warbler | Shaw & Winkler 2011 |
| NE Belize | Magnolia & Yellow Warbler | Spring | Mist-netting | Increased abundance, mass gains in recaptures | Bayly & Gomez 2011 |
| Northern Central America (6 & 7) | | | | | |
| Nicaragua | Red-eyed Vireo | Spring | Geolocators | Stopover averaging 8 days | Callo <i>et al.</i> 2013 |
| Honduras/Guatemala | Cerulean Warbler | Spring | Observations | Stopover suspected in early-mid April | Welton <i>et al.</i> 2012 |
| Guatemala | Swainson's Thrush | Spring | Geolocators | Stopover averaging 11 days | Delmore <i>et al.</i> 2012 |
| Panama & Costa Rica (5) | | | | | |
| Costa Rica | Raptors | Both | Observations | Bottleneck on Caribbean coast | Porras-Peñaranda <i>et al.</i> 2004 |
| Panama | Raptors | Both | Observations | Bottleneck mainly on Pacific coast | Bastista <i>et al.</i> 2005 |
| Panama | Swainson's Thrush | Spring | Geolocators | Stopover averaging 11 days | Delmore <i>et al.</i> 2012 |
| Northern Colombia (3 & 4) | | | | | |
| Sierra Nevada de Santa Marta (SNSM) | Veery | Autumn | Mist-netting & Geolocators | Stopover: body mass gains up to 40% of lean body mass | Bayly <i>et al.</i> 2012a; Heckscher <i>et al.</i> 2012 |
| Darién | Catharus thrushes | Autumn | Mist-netting | Short stopover in some individuals | Gomez <i>et al.</i> 2014 |
| Darién | Diurnal migrants | Autumn | Observations | Bottleneck for raptors, swallows, Chimney Swift, Dickcissel, Common Nighthawk | Bayly <i>et al.</i> 2014 |
| SNSM | Grey-cheeked Thrush & Tennessee Warbler | Spring | Mist-netting | Stopover: body mass gains of 40% and 30% of lean body mass respectively. | Bayly <i>et al.</i> 2013; Bayly <i>et al.</i> 2016 |
| NW Colombia | Red-eyed Vireo | Spring | Geolocators | Stopover averaging 6 days | Callo <i>et al.</i> 2013 |
| Amazon & Llanos (1 & 2) | | | | | |
| Northern edge of Amazon | Red-eyed Vireo & Veery | Spring | Geolocators | Prolonged stopover at north edge of Amazon | Callo <i>et al.</i> 2013; Heckscher <i>et al.</i> 2012 |
| Llanos Colombia / Venezuela | Bobolink | Both | Geolocators | Staging during autumn and spring | Renfrew <i>et al.</i> 2013 |

Along the major flyways used by species breeding in eastern North America, a number of major stopover regions are emerging (Table 3; Figure. 4), including areas inland of the Gulf Coast and northern Florida in North America and northern Colombia in South America, both of which are used to store sufficient energy to make unbroken flights across the Gulf of Mexico and/or the Caribbean Sea. In particular, the Sierra Nevada de Santa Marta, an isolated massif on Colombia's Caribbean coast, is of considerable strategic importance for a number of species (Bayly *et al.* 2012a, 2016, Gómez *et al.* 2015). Not all birds are capable of such long-distance flights and various points in Central America, such as the Yucatan/south-east Mexico and northern Central America host stopover regions for certain species but we lack sufficient information to highlight regions used by multiple species. No doubt the deployment of geolocators and GPS tags on additional species and the establishment of migration banding stations in little-studied regions such as eastern Nicaragua and Honduras will reveal other critical fuelling regions.

Neotropical stopover regions for western-breeding migrants are almost completely unknown, but sites within south-western Mexico and the Pacific slope of Central America are likely to prove important for species on passage to South America, as they have for western populations of Swainson's Thrush (Ruegg and Smith 2002, Delmore *et al.* 2012), and may mirror findings in north-west Mexico where moult-migration is integral to migration strategies (Rohwer *et al.* 2005).




Figure 4. Major Neotropical stopover regions (dashed circles) and bottlenecks (closed circles) for migratory landbirds identified to date. **1.** Northern Amazon (Veery, Red-eyed Vireo); **2.** Llanos of Venezuela and Colombia (Bobolink); **3.** NW Colombia (bottleneck; also stopovers by Red-eyed Vireo & *Catharus* thrushes); **4.** Sierra Nevada de Santa Marta (Tennessee Warbler, Grey-checked Thrush, Veery); **5.** Costa Rica Caribbean slope & Panama Canal Zone (bottleneck for raptors); **6.** Western Nicaragua (Red-eyed Vireo); **7.** Highlands Honduras, Guatemala and Belize (Cerulean Warbler); **8.** Yucatan Peninsula (Purple Martin, Red-eyed Vireo, Magnolia Warbler); **9.** Veracruz (bottleneck for raptors, Kentucky and Hooded Warblers, Scissor-tailed Flycatcher); **10.** SW Mexico (Swainson's Thrush). See Table 3 for references.

Indeed, while this region is known to hold a diverse assemblage of over-wintering migrants (Howell and Webb 1995, Hutto 1997, Fagan and Komar 2016), it may also host stopover populations of long-distance migrant species such as Yellow-billed Cuckoo *Coccyzus americanus*, Western Wood-Pewee *Contopus sordidulus* and Olive-sided Flycatcher *Contopus cooperi*.

Threats facing landbirds on migration in the Neotropics

Just as we learn about the importance of key regions to long-distance migratory birds within the Neotropics, new concerns are being raised about threats to specific sites and vital habitats within those regions, as well as the predicted effects of climate change. Habitat loss continues to be the leading threat to migratory bird populations throughout their annual cycle (Faaborg *et al.* 2010a, Rosenberg *et al.* 2016), and in areas where birds are highly concentrated during migration stopover, loss or degradation of critical habitats may contribute disproportionately to overall population declines (Baker *et al.* 2004, Newton 2006). In addition, direct mortality of migrating birds due to collisions with buildings, communication towers and other structures could be magnified in areas where large proportions of species' populations concentrate. Although astounding numbers of birds are thought to perish through collisions in North America alone (Longcore *et al.* 2013), including significant percentages of global populations of species of concern such as Golden-winged Warbler (Arnold and Zink 2011), equivalent data from south of the USA are few and far between (Agudelo-Álvarez *et al.* 2010). Nonetheless, urban and resort development along coastlines, and proliferation of wind-energy and communications infrastructure along coasts, narrow landmasses, and mountain passes could pose major challenges for migratory bird conservation (Cohen *et al.* 2017).

Research to date indicates that the majority of landbirds stopping over in Central and northern South America rely on native forests, especially pre-montane forests on Caribbean-facing slopes of coastal mountains, such as in the Sierra Nevada de Santa Marta in Colombia, as well as lowland tropical wet and dry forests along the coasts (Johnson and Winker 2008, Welton *et al.* 2012, Gómez *et al.* 2015). As elsewhere in the Neotropics, these tropical forests are under severe threat from expanding agriculture and development, and several of the stopover regions identified to date are experiencing high rates of deforestation and are considered hotspots for future deforestation (Wassenaar *et al.* 2007). In addition, bird-friendly practices which have the potential to mitigate forest loss, such as shade-grown coffee, may not be as beneficial to birds on stopover (Bayly *et al.* 2016) as they are considered to be for over-wintering migrants (Komar 2006).

Much of south-eastern Mexico, such as in the state of Veracruz, has been deforested and although this trend has been reversed in recent years (Ruelas Inzunza *et al.* 2009), the extremely low percentage of remaining forest cover likely remains a barrier to a successful migration for many species. The expansion of cattle pastures in north-western and eastern Honduras and eastern Nicaragua also represents a major threat to forests that are expected to constitute important stopover habitats. In particular, the montane forests of north-western Honduras that are used by the threatened Cerulean Warbler during spring migration (Welton *et al.* 2012) are under considerable pressure. Likewise, in Colombia, the Darién bottleneck is a hotspot for deforestation to make way for cattle pastures. The Sierra Nevada de Santa Marta is also under pressure, as both coffee cultivation and cattle pastures move up slope (Bayly *et al.* 2012b). In western Mexico, loss of dry tropical forest (Trejo and Dirzo 2000) has likely diminished habitat, where many species have been shown to require primary forest (Hutto 1989).

While populations of most raptor species that migrate through the Neotropics are stable or increasing (Ruelas Inzunza *et al.* 2010a), dramatic changes along the main flyway could reverse current fortunes, particularly for species concentrating in large proportion in narrow corridors. The development of wind farms, in particular, has been cited as a potential hazard for migrating raptors but mortality rates reported elsewhere do not support this claim (De Lucas *et al.* 2008, Marques *et al.* 2014). Given our excellent knowledge of the routes taken by raptors, however, any

potentially detrimental projects can be identified rapidly and measures to mitigate their impact can be proposed based on sound scientific justification.

The next steps: research and conservation needs

The last five years have seen an upsurge in new information on the migratory strategies of a variety of landbirds, largely due to the innovation of geolocator technology (McKinnon *et al.* 2013a), but also as a result of on-the-ground field studies and monitoring programmes in northern Central America (Martínez Leyva *et al.* 2005), Colombia (Bayly *et al.* 2013) and Costa Rica (Elizondo-Camacho and Ralph 2012). Despite our growing understanding of migratory strategies and stopover site use, we are still only scratching the surface of a topic that will require decades of dedicated research to determine the needs of most species. To increase our understanding of migration strategies, and to identify the most important stopover sites and habitats for declining Nearctic-Neotropical migratory birds, we recommend a coordinated research initiative combining targeted field studies with the latest technologies across key regions of the Neotropics: including regions 1, 2, 3, 6, 7, and 10 in Figure 3, premontane forests (600–1,500 m) in Panama and Costa Rica, Pacific dry forests in Costa Rica and Nicaragua, lowland dry forests in northern Colombia, and a general exploration of regions south of the equator. The goals of this initiative would be to (1) identify new and previously unknown stopover regions for migratory landbirds; (2) evaluate sites and habitats within those regions that are most critical for successful migration; and (3) develop strategies to conserve these sites and habitats.

The deployment of geolocators or GPS tags on a wider range of landbird species, especially those of conservation concern, is a clear research priority in order to identify new stopover regions, and studies are already underway on Cerulean, Golden-winged, Canada *Cardellina canadensis*, Kirtland's *Setophaga kirtlandii* and Prothonotary *Protonotaria citrea* Warblers. It is also necessary to build on successful geolocator studies by ensuring representative sampling across breeding/wintering ranges, thereby shedding light on how stopover use varies between populations (Jahn *et al.* 2013, Fraser *et al.* 2013, Hobson and Kardynal 2015) or the degree to which populations converge on major stopover regions, as occurs in Purple Martin (Fraser *et al.* 2013), Veery (González-Prieto *et al.* 2011, Heckscher *et al.* 2011, Hobson and Kardynal 2015) and Bobolink (Renfrew *et al.* 2013).

While geolocators enable us to rapidly identify broad stopover regions, their lack of precision and inability to tell us about the habitats and resources that birds are using, means that more precise technology (e.g., GPS tags) and on-the-ground field studies in the Neotropics are essential. Such studies will not only help to pinpoint the actual location of birds within the broad regions identified by geolocators (McKinnon *et al.* 2013b) but also determine what habitats are used and how their quality varies for birds on stopover (Bayly *et al.* 2016). We therefore recommend that geolocator studies be followed up or complemented by standardised surveys within these broad regions, in combination with multi-year mark-recapture studies, that can: (1) estimate occupancy and persistence of landbirds by region and habitat during migration; (2) determine the energetic contribution of a given stopover region to migration strategies; (3) provide site-specific habitat and resource associations for defining realistic conservation priorities; and (4) provide estimates of use at the population level for communities of migrants.

The recapture of individually marked birds, for example, is essential to estimate stopover durations (Schaub *et al.* 2001), rates of energy accumulation (Schaub and Jenni 2000), and ultimately the contribution that a given stopover site makes to the overall energetic needs of migration (Bayly *et al.* 2012a). By comparing rates of mass gain (a surrogate for energy deposition) between habitats, we can also examine habitat quality (Dunn 2001, Bayly *et al.* 2016). Mark-recapture studies can also make use of innovative techniques to gather data on difficult-to-catch and canopy-dwelling species, including the use of coded nano-tags and tower arrays to determine stopover durations and subsequent movements of tagged individuals (Gómez *et al.* 2017, Taylor *et al.* 2017).

Once we know where to focus conservation resources, the scale of action required to ensure adequate habitat availability in major stopover regions throughout the Neotropics is potentially overwhelming. To be successful, strategies to conserve stopover habitats need to be integrated within regional conservation efforts, including those focused on Neotropical resident and threatened species, protection of watersheds, reducing deforestation and socio-economic development (Barlow *et al.* 2016). Any approach must go above and beyond the strategic purchase of private reserves, given the considerable areas occupied by migratory birds, and include activities such as lobbying for changes in detrimental agricultural and land-use policies, community-based conservation efforts, ecotourism activities, promotion of migrant friendly agricultural practices such as shade-grown coffee and silvopastures, and the use of tools such as Payment for Environmental Services. Indeed, as we identify new stopover sites, the actions required to ensure that they continue to function as key links in the lives of migratory birds will have to be tailored to the unique conditions in each.

As bird conservation plans have focused largely on breeding habitat, and to a lesser extent on overwintering sites, new information on the specific needs of birds on migration must be incorporated into full life-cycle bird conservation plans (Berlanga *et al.* 2010, Rosenberg *et al.* 2016). Only through large-scale and highly collaborative efforts, such as the GoMAMN (<https://www.gomamn.org/>, Cohen *et al.* 2017) or the *Neotropical Flyways Project* (<http://selva.org.co/research-programs/migratory-species/nfp/>) will we be able to understand and conserve the spectacle of migration throughout the Western Hemisphere.

While our suggestions for continued and future research will yield important advances that can help guide conservation, some habitats and regions are so threatened that we can't afford to wait for the next generation of data. Thus, we want to emphasise the importance of continuing ongoing habitat conservation and restoration efforts throughout the region, especially in natural habitats already in decline, as these efforts will undoubtedly benefit Neotropical migrant and resident species alike.

Acknowledgements

This review grew out of the Migration Stopover Special Session at the *Partners in Flight V Conference and Conservation Workshop* in Snowbird, Utah in August 2013, and we thank the organisers of Partners in Flight V, especially American Bird Conservancy, for including the session within the programme. Having begun as an overview of our knowledge prior to the special session, the discussions during the session highlighted the need for this review. We thank all participants in the session for their contributions and insights and subsequent discussions, especially, Rosa M. Vidal, Elisa Peresbarbosa, C. J. Ralph, Melinda Welton and Pablo Elizondo. This is contribution No. 1 of the Neotropical Flyways Project and Hawk Mountain conservation science contribution number 285. We thank an anonymous reviewer for suggestions that improved this manuscript. Attendance at PIFV by NB and KVR was supported by the Cornell Lab of Ornithology, and travel support for other conference participants was provided by American Bird Conservancy.

References

- Agudelo-Álvarez, L., Moreno-Velasquez, J. and Ocampo-Peñuela, N. (2010) Collisions of birds with windows in a university campus in Bogotá, Colombia. *Ornitol. Colomb.* 10: 3–10.
- Alerstam, T. (2009) Fly by night or day? Optimal daily timing of bird migration. *J. Theor. Biol.* 258: 530–536.
- Alerstam, T. (2011) Optimal bird migration revisited. *J. Ornithol.* 152: 5–23.
- Arnold, T. and Zink, R. (2011) Collision mortality has no discernible effect on population trends of North American birds. *PLoS One* 6: e24708.
- Atkinson, P. W., Baker, A. J., Bennett, K. A., Clark, N. A., Clark, J. A., Cole, K. B., *et al.*

- (2007) Rates of mass gain and energy deposition in red knot on their final spring staging site is both time- and condition-dependent. *J. Appl. Ecol.* 44: 885–895.
- Baker, A. J., González, P. M., Piersma, T., Niles, L. J., do Nascimento, I., Atkinson, P. W., et al. (2004) Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proc. R. Soc. London B* 271: 875–882.
- Barlow, J., Lennox, G. D., Ferreira, J., Berenguer, E., Lees, A. C., Mac Nally, R., Thomson, J. R., de Barros Ferraz, S. F., Louzada, J., Oliveira, V. H. F. and Parry, L. (2016) Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature* 535: 144–147.
- Batista, C., Miro, G., Angehr, G. and Bildstein, K. L. (2005) More than three million migrating raptors counted ocean-to-ocean in Panama, autumn 2004. *Hawk Migr. Stud.* 31: 5–6.
- Bauer, S. and Hoye, B. J. (2014) Migratory animals couple biodiversity and ecosystem functioning Worldwide. *Science* 344: 1242–1252.
- Bayly, N. J., Cárdenas-Ortiz, L., Rubio, M. and Gómez, C. (2014) Migration of raptors, swallows and other diurnal migratory birds through the Darién of Colombia. *Ornitol. Neotrop.* 25: 63–71.
- Bayly, N. J. and Gómez, C. (2011) Comparison of autumn and spring migration strategies of Neotropical migratory landbirds in north-east Belize. *J. Field Ornithol.* 82: 117–131.
- Bayly, N. J., Gómez, C. and Hobson, K. A. (2013) Energy reserves stored by migrating Gray-cheeked Thrushes *Catharus minimus* at a spring stopover site in northern Colombia are sufficient for a long-distance flight to North America. *Ibis* 155: 271–283.
- Bayly, N. J., Gómez, C., Hobson, K. A., González, A. M. and Rosenberg, K. V. (2012a) Fall migration of the Veery (*Catharus fuscescens*) in northern Colombia: determining the energetic importance of a stopover site. *Auk* 129: 449–459.
- Bayly, N. J., Paez, A., Gómez, C. and Mora, C. (2012b) *Conservation hotspots for migratory landbirds in the Sierra Nevada de Santa Marta, Colombia*. Bogotá: SELVA: Investigación para la Conservación en el Neotropico.
- Bayly, N. J., Gómez, C., Hobson, K. A. and Rosenberg, K. V. (2016) Prioritizing tropical habitats for long-distance migratory songbirds - an assessment of habitat quality at a stopover site in Colombia. *Avian Conserv. Ecol.* 11: 5.
- Bearhop, S., Hilton, G. M., Votier, S. C. and Waldron, S. (2004) Stable isotope ratios indicate that body condition in migrating passerines is influenced by winter habitat. *Proc. R. Soc. London B* 271: S215–218.
- Beason, J. P., Gunn, C., Potter, K. M., Sparks, R. A. and Fox, J. W. (2012) The Northern Black Swift: Migration path and wintering area revealed. *Wilson J. Ornithol.* 124: 1–8.
- Berlanga, H., Kennedy, J. A., Rich, T. D., Arizmendi, M. C., Beardmore, C. J., Blancher, P. J., et al. (2010) *Saving Our shared birds: Partners in flight tri-national vision for landbird conservation*. Ithaca: Cornell Lab of Ornithology.
- Bildstein, K. L. (2004) Raptor migration in the Neotropics: patterns, processes, and consequences. *Ornitol. Neotrop.* 15: 83–99.
- Bildstein, K. L. and Saborio, M. (2000) Spring migration counts of raptors and New World Vultures in Costa Rica. *Ornitol. Neotrop.* 11: 197–205.
- Buler, J. J., Moore, F. R. and Woltmann, S. (2007) A multi-scale examination of stopover habitat use by birds. *Ecology* 88: 1789–1802.
- Callo, P. A., Morton, E. S. and Stutchbury, B. J. M. (2013) Prolonged spring migration in the Red-Eyed Vireo (*Vireo olivaceus*). *Auk* 130: 240–246.
- Carlisle, J. D., Skagen, S. K., Kus, B. E., van Riper, C., Paxton, K. L. and Kelly, J. F. (2009) Landbird migration in the American West: Recent progress and future research directions. *Condor* 111: 211–225.
- Cohen, E. B., Barrow, W. C., Buler, J. J., Deppe, J. L., Farnsworth, A., Marra, P. P., McWilliams, S. R., Mehlman, D. W., Wilson, R. R., Woodrey, M. S. and Moore, F. R. (2017) How do en route events around the Gulf of Mexico influence migratory landbird populations. *Condor* 119: 327–343.
- Colorado, G. J. (2010) Fall migration of Empidonax flycatchers in northwestern Colombia. *J. f. Ornithol.* 81: 259–266.
- Colorado, G. J., Bechard, M. J., Márquez, C. and Castaño, A. (2006) Raptor migration in the

- Cauca river valley of northern Colombia. *Ornitol. Neotrop.* 17: 161–172.
- De Lucas, M., Janss, G., Whitfield, D. P. and Ferrer, M. (2008) Collision fatality of raptors in wind farms does not depend on raptor abundance. *J. Appl. Ecol.* 45: 1695–1703.
- DeGraaf, R. M. and Rappole, J. H. (1996) *Neotropical migratory birds*. Ithaca: Cornell University Press.
- Delingat, J., Dierschke, V., Schmaljohann, H., Mendel, B., Bairlein, F., Delingat, J., *et al.* (2006) Daily stopovers as optimal migration strategy in a long-distance migrating passerine: the Northern Wheatear *Oenanthe oenanthe*. *Ardea* 94: 593–605.
- Delmore, K. E., Fox, J. W. and Irwin, D. E. (2012) Dramatic intraspecific differences in migratory routes, stopover sites and wintering areas, revealed using light-level geolocators. *Proc. R. Soc. London B* 279: 4582–4589.
- Delmore, K. E. and Irwin, D. E. (2014) Hybrid songbirds employ intermediate routes in a migratory divide. *Ecol. Lett.* 17: 1211–1218.
- DeLuca, W. V., Woodworth, B. K., Rimmer, C. C., Marra, P. P., Taylor, P. D., McFarland, K. P., *et al.* (2015) Transoceanic migration by a 12 g songbird. *Biol. Lett.* 11.
- Deppe, J. L. and Rotenberry, J. T. (2005) Temporal patterns in fall migrant communities in the Yucatan, Mexico. *Condor* 107: 228–243.
- Drake, A., Rock, C., Quinlan, S., Martin, M. and Green, D. (2014) Wind speed during migration influences the survival, timing of breeding, and productivity of a neotropical migrant, *Setophaga petechia*. *PLoS One* 9: 92–114.
- Dunn, E. H. (2001) Mass change during migration stopover: a comparison of species groups and sites. *J. Field Ornithol.* 72: 419–432.
- Dunn, E. H. (2002) A cross-Canada comparison of mass change in birds during migration stopover. *Wilson Bull.* 114: 368–379.
- Elizondo-Camacho, P. and Ralph, C. J. (2012) Contributions of two decades of bird banding and monitoring to the science and conservation of birds in Costa Rica. *North Am. Bird Bander* 37: 126.
- Ewert, D. N., Hamas, M. J., Smith, R. J., Dallman, M. E. and Jorgensen, S. W. (2011) Distribution of migratory landbirds along the northern Lake Huron shoreline. *Wilson J. Ornithol.* 123: 536–547.
- Faaborg, J., Holmes, R. T., Anders, A. D., Bildstein, K. L., Dugger, K. M., Gauthreaux, S. A., Heglund, P., Hobson, K. A., Jahn, A. E., Johnson, D. H., Latta, S. C., Levey, D. J., Marra, P. P., *et al.* (2010a) Conserving migratory land birds in the New World: Do we know enough? *Ecol. Appl.* 20: 398–418.
- Faaborg, J., Holmes, R. T., Anders, A. D., Bildstein, K. L., Dugger, K. M., Gauthreaux, S. A., Heglund, P., Hobson, K. A., Jahn, A. E., Johnson, D. H., Latta, S. C., Levey, D. J., Marra, P. P., *et al.* (2010b) Recent advances in understanding migration systems of New World land birds. *Ecol. Monogr.* 80: 3–48.
- Fagan, J. and Komar, O. (2016) *Peterson field guide to birds of Northern Central America*. New York: Houghton Mifflin Harcourt Publishing Co.
- Finch, D. M. and Martin, T. E., eds. (1995) *Ecology and management of Neotropical migratory birds: A synthesis and review of critical issues*. New York: Oxford University Press.
- Finch, T., Pearce-Higgins, J., Leech, D. I. and Evans, K. (2014) Carry-over effects from passage regions are more important than breeding climate in determining the breeding phenology and performance of three avian migrants of conservation concern. *Biodivers. Conserv.* 23: 2427–2444.
- Fraser, K. C., Stutchbury, B. J. M., Kramer, P., Silverio, C., Barrow, J., Newstead, D., *et al.* (2013) Consistent range-wide pattern in fall migration strategy of Purple Martin (*Progne subis*), despite different migration routes at the Gulf of Mexico. *Auk* 130: 291–296.
- Fraser, K. C., Stutchbury, B. J. M., Silverio, C., Kramer, P. M., Barrow, J., Newstead, D., *et al.* (2012) Continent-wide tracking to determine migratory connectivity and tropical habitat associations of a declining aerial insectivore. *Proc. R. Soc. London B* 279: 4901–4906.
- Fuller, M. R., Seegar, W. S. and Schueck, L. S. (1998) Routes and travel rates of Peregrine Falcons and Swainson's Hawks in the Western Hemisphere. *J. Avian Biol.* 29: 433–440.
- Galindo, P., Méndez, E. and Adames, A. (1963) Banding of migrant thrushes in Almirante, Panama. *Bird Band.* 34: 202–209.

- Garrido, O. H. and Kirkconnell, A. (2000) *Field guide to the birds of Cuba*. Ithaca: Cornell University Press.
- Gill, R. E., Tibbitts, T. L., Douglas, D. C., Handel, C. M., Mulcahy, D. M., Gottschalck, J. C., *et al.* (2009) Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier? *Proc. R. Soc. B Biol. Sci.* 276: 447–457.
- Gómez, C., Bayly, N. J., Norris, R. D., Mackenzie, S. A., Rosenberg, K. V., Taylor, P. D., Hobson, K. A., Cadena, C. D. (2017) Fuel loads acquired at a stopover site influence the pace of intercontinental migration in a boreal songbird. *Sci. Reports* 8: 3405.
- Gómez, C., Gómez-Bahamón, V., Cárdenas-Ortiz, L. and Bayly, N. J. (2015) Distribution of Nearctic–Neotropical migratory birds along a South American elevation gradient during spring migration. *Wilson J. Ornithol.* 127: 72–86.
- Gómez, C., Bayly, N. J. and Rosenberg, K. V. (2014) Fall stopover strategies of three species of thrush (*Catharus*) in northern South America. *Auk* 131: 702–717.
- Gómez, C., Bayly, N. J. and Rosenberg, K. V. (2013) Seasonal variation in stopover site use: *Catharus* thrushes and vireos in northern Colombia. *J. Ornithol.* 154: 107–117.
- Gómez, C. and Bayly, N. J. (2011) Migration of Empidonax flycatchers through northeast Belize. *Ornitol. Neotrop.* 22: 339–345.
- González-Prieto, A. M. and Hobson, K. A. (2013) Environmental conditions on wintering grounds and during migration influence spring nutritional condition and arrival phenology of Neotropical migrants at a northern stopover site. *J. Ornithol.* 154: 1067–1078.
- González-Prieto, A. M., Hobson, K. A., Bayly, N. J. and Gómez, C. (2011) Geographic origins and timing of fall migration of the Veery in northern Colombia *Condor* 113: 860–868.
- Guillemain, M., Elmer, J., Arzel, C., Johnson, A. R. and Simon, G. (2008) The income-capital breeding dichotomy revisited: late winter body condition is related to breeding success in an income breeder. *Ibis* 150: 172–176.
- Gwinner, E. (1996) Circadian and circannual programmes in avian migration. *J. Exp. Biol.* 199: 39–48.
- Hagan, J. M. and Johnston, D. W., eds. (1992) *Ecology and conservation of Neotropical migrant landbirds*. Washington DC: Smithsonian Institution Press.
- Harrison, X. A., Blount, J. D., Inger, R., Norris, D. R. and Bearhop, S. (2011) Carry-over effects as drivers of fitness differences in animals. *J. Anim. Ecol.* 80: 4–18.
- Heckscher, C. M., Taylor, S. M., Fox, J. W. and Afanasyev, V. (2011) Veery (*Catharus fuscescens*) wintering locations, migratory connectivity, and a revision of its winter range using geolocator technology. *Auk* 128: 531–542.
- Hedenström, A. (2008) Adaptations to migration in birds: behavioural strategies, morphology and scaling effects. *Philos. Trans. R. Soc. London* 363: 287–299.
- Heglund, P. J. and Skagen, S. K. (2005) Ecology and physiology of en route Nearctic–Neotropical migratory birds: a call for collaboration. *Condor* 107: 193–196.
- Hicks, D., Rogers, D. and Child, G. (1966) Autumnal hawk migration through Panama. *Bird Band.* 37: 121–123.
- Hobson, K. A. and Kardynal, K. J. (2015) Western Veeries use an eastern shortest-distance pathway: new insights to migration routes and phenology using light-level geolocators. *Auk* 132: 540–550.
- Holmes, R. T. (2007) Understanding population change in migratory songbirds: long-term and experimental studies of Neotropical migrants in breeding and wintering areas. *Ibis* 149: 2–13.
- Holmes, R. T., Marra, P. P. and Sherry, T. W. (1996) Habitat-specific demography of breeding Black-throated Blue Warblers (*Dendroica caerulescens*): implications for population dynamics. *J. Anim. Ecol.* 65: 183–196.
- Howell, S. and Webb, S. (1995) *A guide to the birds of Mexico and Northern Central America*. Oxford: Oxford University Press.
- Hutto, R. L. (1989) The effect of habitat alteration on migratory land birds in a west Mexican tropical deciduous forest: A conservation perspective. *Conserv. Biol.* 3: 138–148.
- Hutto, R. L. (1997) A description of mixed-species insectivorous bird flocks in western Mexico. *Condor* 89: 282–292.

- Hutto, R. L. (1998) On the importance of stopover sites to migrating birds. *Auk* 115: 823–825.
- Jahn, A. E., Cueto, V. R., Fox, J. W., Husak, M. S., Kim, D. H., Landoll, D. V., *et al.* (2013) Migration timing and wintering areas of three species of flycatchers (*Tyrannus*) breeding in the Great Plains of North America. *Auk* 130: 247–257.
- Johnson, A. B. and Winker, K. (2008) Autumn stopover near the Gulf of Honduras by Nearctic–Neotropical migrants. *Wilson J. Ornithol.* 120: 277–285.
- Johnson, M. D., Sherry, T. W., Holmes, R. T. and Marra, P. P. (2006) Assessing habitat quality for a migratory songbird wintering in natural and agricultural habitats. *Conserv. Biol.* 20: 1433–1444.
- Keast, A. and Morton, E. S., eds. (1980) *Migrant birds in the Neotropics: ecology, behavior, distribution, and conservation*. Washington DC: Smithsonian Institution Press.
- Kelly, J. F. and Hutto, R. L. (2005) An east-west comparison of migration in North American Wood Warblers. *Condor* 107: 197–211.
- Klaassen, R. H. G., Alerstam, T., Carlsson, P., Fox, J. W. and Lindström, Å. (2011) Great flights by Great Snipes: long and fast non-stop migration over benign habitats. *Biol. Lett.* 7: 833–835.
- Kochert, M. N., Fuller, M. R., Schueck, L. S., Bond, L., Bechard, M. J., Woodbridge, B., *et al.* (2011) Migration patterns, use of stopover areas, and austral summer movements of Swainson's Hawks. *Condor* 113: 89–106.
- Komar, O. (2006) Ecology and conservation of birds in coffee plantations: a critical review. *Bird Conserv. Interantn.* 16: 1–23.
- Kristensen, M. W., Tøttrup, A. P. and Thorup, K. (2013) Migration of the Common Redstart (*Phoenicurus phoenicurus*): a Eurasian songbird wintering in highly seasonal conditions in the West African Sahel. *Auk* 130: 258–264.
- La Sorte, F. A., Fink, D., Hochachka, W. M., DeLong, J. P. and Kelling, S. (2014a) Spring phenology of ecological productivity contributes to the use of looped migration strategies by birds. *Proc. R. Soc. London B* 281: 20140984.
- La Sorte, F. A., Fink, D., Hochachka, W. M., Farnsworth, A., Rodewald, A. D., Rosenberg, K. V., *et al.* (2014b) The role of atmospheric conditions in the seasonal dynamics of North American migration flyways. *J. Biogeogr.* 41: 1685–1696.
- La Sorte, F. A., Fink, D., Hochachka, W. M. and Kelling, S. (2016) Convergence of broad-scale migration strategies in terrestrial birds. *Proc. R. Soc. London B* 283: 20152588.
- LaManna, J., George, T., Saracco, J. F., Nott, M. P. and Desante, D. F. (2012) El Niño–Southern Oscillation influences annual survival of a migratory songbird at a regional scale. *Auk* 129: 1–10.
- Latta, S. C. and Brown, C. (1999) Autumn stopover ecology of the Blackpoll Warbler (*Dendroica striata*) in thorn scrub forest of the Dominican Republic. *Can. J. Zool.* 77: 1147–1156.
- Latta, S. C. and Faaborg, J. (2002) Demographic and population responses of Cape May Warblers wintering in multiple habitats. *Ecology* 83: 2502–2515.
- Latta, S. C., Rimmer, C. C., Keith, A., Wiley, J., Raffaele, H., McFarland, K. P., *et al.* (2006) *Birds of the Dominican Republic and Haiti*. Princeton: Princeton University Press.
- Laughlin, A. J., Taylor, C. M., Bradley, D. W., LeClair, D., Clark, R. G., Dawson, R. D., *et al.* (2013) Integrating information from geolocators, weather radar, and citizen science to uncover a key stopover area of an aerial insectivore. *Auk* 130: 230–239.
- Lemke, H., Tarka, M., Klaassen, R., Akesson, M., Bensch, S. and Hasselquist, D. (2013) Annual cycle and migration strategies of a trans-Saharan migratory songbird: a geolocator study in the Great Reed Warbler. *PLoS One* 8: e79209.
- Leu, M. and Thompson, C. W. (2002) The potential importance of migratory stopover sites as flight feather molt staging areas: a review for neotropical migrants. *Biol. Conserv.* 106: 45–56.
- Lindström, Å. (1990) The role of predation risk in stopover habitat selection in migrating Bramblings, *Fringilla montifringilla*. *Behav. Ecol.* 1: 102–106.
- Lindström, Å., Alerstam, T., Bahlenberg, P., Ekblom, R., Fox, J. W., Råghall, J., *et al.* (2016) The migration of the Great Snipe

- Gallinago media*: intriguing variations on a grand theme. *J. Avian Biol.* 47: 321–334.
- Longcore, T., Rich, C., Mineau, P., MacDonald, B., Bert, D. G., Sullivan, L. M., *et al.* (2013) Avian mortality at communication towers in the United States and Canada: which species, how many, and where? *Biol. Conserv.* 158: 410–419.
- Lowery, G. H. (1945) Trans-Gulf spring migration of birds and the coastal hiatus. *Wilson Bull.* 57: 92–121.
- Marques, A. T., Batalha, H., Rodrigues, S., Costa, H., Pereira, M. J. R., Fonseca, C., *et al.* (2014) Understanding bird collisions at wind farms: An updated review on the causes and possible mitigation strategies. *Biol. Conserv.* 179: 40–52.
- Marra, P. P. (2000) The role of behavioral dominance in structuring patterns of habitat occupancy in a migrant bird during the non-breeding season *Behav. Ecol.* 11: 299–308.
- Martell, M. S., Henny, C. J., Nye, P. E. and Solensky, M. J. (2001) Fall migration routes, timing, and wintering sites of North American Ospreys as determined by satellite telemetry. *Condor* 103: 715–724.
- Martin, T. E. and Finch, D. M. (eds) (1995) *Ecology and management of Neotropical migratory birds*. New York: Oxford University Press.
- Martínez Leyva, E., Ruelas Inzunza, E., Cruz Carretero, O., Barr, J. L., Peresbarbosa Rojas, E., Chávez Domínguez, I., *et al.* (2005) Dynamics of passerine migration in Veracruz, Mexico. Pp. 62–70 in: *Proceedings of the Fourth International Partners in Flight Conference: Tundra to Tropics*. Washington: Partners in Flight.
- McKinnon, E. A., Fraser, K. C. and Stutchbury, B. J. M. (2013a) New discoveries in landbird migration using geolocators, and a flight plan for the future. *Auk* 130: 211–222.
- McKinnon, E. A., Stanley, C. Q., Fraser, K. C., MacPherson, M., Casbourn, G., Marra, P. P., *et al.* (2013b) Estimating geocator accuracy for a migratory songbird using live ground-truthing in tropical forest. *Anim. Migr.* 1: 31–38.
- Mehlman, D. W., Mabey, S. E., Ewert, D. N. and Duncan, C. (2005) Conserving stopover sites for forest-dwelling migratory landbirds. *Auk* 122: 1281–1290.
- Moore, F. R. and Buler, J. J. (2011) Migrant-habitat relationships during stopover along an ecological barrier: extrinsic constraints and conservation implications. *J. Ornithol.* 152 (Suppl.): 101–112.
- Moore, F. R., Kerlinger, P. and Simmons, T. R. (1990) Stopover on a Gulf coast barrier island by spring trans-Gulf migrants. *Wilson Bull.* 102: 487–500.
- Nebel, S. A., Mills, J. D., McCracken, J. D. and Taylor, P. D. (2010) Declines of aerial insectivores in North America follow a geographic gradient. *Avian Conserv. Ecol.* 5: 1.
- Newton, I. (2004) Population limitation in migrants. *Ibis* 146: 197–226.
- Newton, I. (2006) Can conditions experienced during migration limit the population levels of birds? *J. Ornithol.* 147: 146–166.
- Newton, I. (2008) *The migration ecology of birds*. London: Elsevier.
- Norris, D. R. and Marra, P. P. (2007) Seasonal interactions, habitat quality, and population dynamics in migratory birds. *Condor* 109: 535–547.
- Norris, D. R., Marra, P. P., Kyser, T. K., Sherry, T. W. and Ratcliffe, L. M. (2004) Tropical winter habitat limits reproductive success on the temperate breeding grounds in a migratory bird. *Proc. R. Soc. London B* 271: 59–64.
- Nott, M. P., Desante, D. F., Siegel, R. B. and Pyle, P. (2002) Influences of the El Niño/Southern Oscillation and the North Atlantic Oscillation on avian productivity in forests of the Pacific Northwest of North America. *Glob. Ecol. Biogeogr.* 11: 333–342.
- Piersma, T. (1987) Hop, skip or jump: constraints on migration of Arctic waders by feeding, fattening and flight speed. *Limosa* 60: 185–194.
- Pomeroy, A. C. (2006) Experimental evidence that migrants adjust usage at a stopover site to trade off food and danger. *Behav. Ecol.* 17: 1041–1045.
- Porras-Peñaranda, P., Robichaud, L. and Branch, F. (2004) New full-season count sites for raptor migration in Talamanca, Costa Rica. *Ornitol. Neotrop.* 15 (suppl.): 267–278.
- Pyle, P., Leitner, W., Lozano-Angulo, L., Avilez-Teran, F., Swanson, H., Gómez Limón, E., *et al.* (2009) Temporal, spatial, and annual variation in the occurrence of molt-migrant Passerines in the Mexican monsoon region. *Condor* 111: 583–590.

- Raffaele, H., Wiley, J., Garrido, O., Keith, A. and Raffaele, J. (2003) *Birds of the West Indies*. Princeton: Princeton University Press.
- Remsen, J. V. R. (2001) True winter range of the Veery (*Catharus fuscescens*): lessons for determining winter ranges of species that winter in the tropics. *Auk* 118: 838–848.
- Renfrew, R. B., Kim, D., Perlut, N., Smith, J., Fox, J. and Marra, P. P. (2013) Phenological matching across hemispheres in a long-distance migratory bird. *Divers. Distrib.* 19: 1008–1019.
- Rich, T. D., Beardmore, C. J., Berlanga, H., Blancher, P. J., Bradstreet, M. S., Butcher, G. S., *et al.* (2004) *Partners in Flight: North American landbird conservation plan*. Ithaca: The Cornell Lab of Ornithology.
- Robbins, C. S., Sauer, J. R., Greenberg, R. S. and Droege, S. (1989) Population declines in North American birds that migrate to the neotropics. *Proc. Natl. Acad. Sci. USA* 86: 7658–7662.
- Robson, D., Barriocanal, C., Garcia, O. and Villena, O. (2001) The spring stopover of the Reed Warbler *Acrocephalus scirpaceus* in northeast Spain. *Ring. Migr.* 20: 233–238.
- Rockwell, S. M., Wunderle, J. M., Sillett, T. S., Bocetti, C. I., Ewert, D. N., Currie, D., *et al.* (2016) Seasonal survival estimation for a long-distance migratory bird and the influence of winter precipitation. *Oecologia Online*: 1–12.
- Rodríguez-Santana, F., Vega, Y. S., Padilla, M. S., Plasencia León, C., Adán, Y. E. T., Losada, M. S., *et al.* (2014) Magnitude and timing of autumn Osprey migration in southeastern Cuba. *J. Raptor Res.* 48: 334–344.
- Rogers, D. T. (1965) Fat levels and estimated flight-ranges of some autumn migratory birds killed in Panama during a nocturnal rainstorm. *Bird Band.* 36: 115–116.
- Rogers, D. T. and Odum, E. P. (1966) A study of autumnal postmigrant weights and vernal fattening of North American migrants in the tropics. *Wilson Bull.* 78: 415–433.
- Rohwer, S., Butler, L. and Froehlich, D. (2005) Ecology and demography of east-west differences in molt scheduling of neotropical migrant passerines. Pp. 87–105 in R. Greenberg and P. Marra, eds. *Birds of two worlds: the ecology and evolution of migration*. Baltimore: Johns Hopkins University Press.
- Rosenberg, K. V., Kennedy, J. A., Dettmers, R., Ford, R. P., Reynolds, D., Alexander, J. D., *et al.* (2016) *Partners in Flight landbird conservation plan: 2016 revision for Canada and Continental United States*. Ithaca: Cornell Lab of Ornithology.
- Ruegg, K. C. and Smith, T. B. (2002) Not as the crow flies: a historical explanation for circuitous migration in Swainson's thrush (*Catharus ustulatus*). *Proc. R. Soc. Lond. B* 269: 1371–1381.
- Ruelas Inzunza, E., Goodrich, L., Hoffman, S., Martínez, E., Smith, J. P., Peresbarbosa, E., *et al.* (2009) Long-term conservation of migratory birds in México: the Veracruz River of Raptors project. Pp. 577–589 in: *Tundra to tropics: connecting birds, habitats and people. Proceedings of the 4th International Partners in Flight Conference, 13-16 February 2008, McAllen, Texas*. Texas: University of Texas-Pan American Press.
- Ruelas Inzunza, E., Goodrich, L. J. and Hoffman, S. W. (2010a) Changes in migratory raptor populations in Veracruz, Mexico. *Acta Zoológica Mex.* 26: 495–525.
- Ruelas Inzunza, E., Goodrich, L. J. and Hoffman, S. W. (2010b) North American population estimates of waterbirds, vultures and hawks from migration counts in Veracruz, Mexico. *Bird Conserv. Int.* 20: 124–133.
- Sanderson, F. J., Donald, P. F., Pain, D. J., Burfield, I. J. and van Bommel, F. P. J. (2006) Long-term population declines in Afro-Palaearctic migrant birds. *Biol. Conserv.* 131: 93–105.
- Sauer, J. R., Hines, J. E., Pardieck, K. L., Ziolkowski, D. J. and Link, W. A. (2014) *The North American Breeding Bird Survey, Results and Analysis 1966-2013. Version 01.30.2015*. Laurel, MD: USGS Patuxent Wildlife Research Center.
- Schaub, M. and Jenni, L. (2000) Fuel deposition of three passerine bird species along the migration route. *Oecologia* 122: 306–317.
- Schaub, M., Pradel, R., Jenni, L. and Lebreton, J. D. (2001) Migrating birds stop over longer than usually thought: an improved capture-recapture analysis. *Ecology* 82: 852–859.
- Schmaljohann, H. and Dierschke, V. (2004) Optimal bird migration and predation risk: a field experiment with Northern Wheatears *Oenanthe oenanthe*. *J. Anim. Ecol.* 74: 131–138.

- Shaw, D. W. and Winker, K. (2011) Spring stopover and refueling among migrant passerines in the Sierra de Los Tuxtlas, Veracruz, Mexico. *Wilson J. Ornithol.* 123: 575–587.
- Sheehy, J., Taylor, C. M., McCann, K. S. and Norris, D. R. (2010) Optimal conservation planning for migratory animals: integrating demographic information across seasons. *Conserv. Lett.* 3: 192–202.
- Sherry, T. W. and Holmes, R. T. (1995) Summer versus winter limitation of populations: What are the issues and what is the evidence? Pp. 85–120 in T. E. Martin and D. M. Finch, eds. *Ecology and management of Neotropical migratory birds*. New York: Oxford University Press.
- Sillett, T. S. and Holmes, R. T. (2002) Variation in survivorship of a migratory songbird throughout its annual cycle. *J. Anim. Ecol.* 71: 296–308.
- Simons, T. R., Moore, F. R. and Gauthreaux, S. A. (2000) Mist netting trans-Gulf migrants at coastal stopover sites: the influence of spatial and temporal variability on capture data. *Stud. Avian Biol.* 135–143.
- Smith, N. G., Goldstein, D. L. and Bartholomew, G. A. (1986) Is long-distance migration possible for soaring hawks using only stored fat? *Auk* 103: 607–611.
- Stanley, C. Q., MacPherson, M., Fraser, K. C., McKinnon, E. A. and Stutchbury, B. J. M. (2012) Repeat tracking of individual songbirds reveals consistent migration timing but flexibility in route. *PLoS One* 7: e40688.
- Studds, C. E. and Marra, P. P. (2005) Nonbreeding habitat occupancy and population processes: an upgrade experiment with a migratory bird. *Ecology* 86: 2380–2385.
- Stutchbury, B. J. M., Tarof, S., Done, T., Gow, E., Kramer, P. M., Tautin, J., et al. (2009) Tracking long-distance songbird migration by using geolocators. *Science* 323: 896.
- Sullivan, B. L., Aycrigg, J. A., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., Damoulas, T., Dhondt, A. A., Dietrich, T., Farnsworth, A., Fink, D., Fitzpatrick, J. W., Fredricks, T., Gerbracht, J., Gomes, C., Hochachka, W. M., Iliff, M. J., Lagoze, C., La Sorte, F. A., Merrifield, M., Morris, W., Phillips, T. B., Reynolds, M., Rodewald, A. D., Rosenberg, K. V., Trautmann, N. M., Wiggins, A., Winkler, D. W., Wong, W.-K., Wood, C. L., Yu, J. and Kelling, S. (2014) The eBird enterprise: An integrated approach to development and application of citizen science. *Biol. Conserv.* 169: 31–40.
- Sutherland, W. J. (1998) Evidence for flexibility and constraint in migration systems. *J. Avian Biol.* 29: 441–446.
- Taylor, P. D., Crewe, T. L., Mackenzie, S. A., Lepage, D., Aubry, Y., Crysler, Z., Finney, G., Francis, C. M., Guglielmo, C. G., Hamilton, D. J., Holberton, R. L., Loring, P. H., Mitchell, G. W., Norris, D., Paquet, J., Ronconi, R. A., Smetzer, J., Smith, P. A., Welch, L. J., and Woodworth, B. K. (2017) The Motus Wildlife Tracking System: a collaborative research network to enhance the understanding of wildlife movement. *Avian Conserv. Ecol.* 12: 8.
- Taylor, P. D., Mackenzie, S. A., Thurber, B. G., Calvert, A. M., Mills, A. M., McGuire, L. P., et al. (2011) Landscape movements of migratory birds and bats reveal an expanded scale of stopover. *PLoS One* 6: e27054.
- Terborgh, J. W. (1980) The conservation status of Neotropical migrants: Present and future. In A. Keast and E. S. Morton, eds. *Migrant birds in the Neotropics: Ecology, behavior, distribution and conservation*. Washington DC: Smithsonian Inst. Press.
- Tottrup, A. P., Thorup, K., Rainio, K., Yosef, R., Lehtikoinen, E. and Rahbek, C. (2008) Avian migrants adjust migration in response to environmental conditions en route. *Biol. Lett.* 4: 685–688.
- Trejo, I. and Dirzo, R. (2000) Deforestation of seasonally dry tropical forest: a national and local analysis in Mexico. *Biol. Conserv.* 94: 133–142.
- Warnock, N. (2010) Stopping vs. staging: the difference between a hop and a jump. *J. Avian Biol.* 41: 621–626.
- Wassenaar, T., Gerber, P., Verbury, P., Rosales, M., Ibrahim, M. and Steinfeld, H. (2007) Projecting land use changes in the Neotropics: the geography of pasture expansion into forest. *Glob. Environ. Chang.* 17: 86–104.
- Weber, T., Ens, B. and Houston, A. (1998) Optimal avian migration: a dynamic model of fuel stores and site use. *Evol. Ecol.* 12: 377–401.
- Weber, T., Houston, A. and Ens, B. (1999) Consequences of habitat loss at migratory stopover sites: a theoretical investigation. *J. Avian Biol.* 30: 416–426.

- Welton, M. J., Anderson, D. L., Colorado, G. J., Hamel, P. B. and Calderón-F, D. (2012) Spring migration stopover by Cerulean Warblers in northern Middle America. *Ornitol. Neotrop.* 23: 289–305.
- Wilcove, D. S. and Wikelski, M. (2008) Going, going, gone: is animal migration disappearing. *PLoS One* 6: e1888.
- Wilson, S., Hobson, K. A., Collister, D. M. and Wilson, A. G. (2008) Breeding destinations and spring migration patterns of Swainson's Thrush (*Catharus ustulatus*) at a Costa Rican stopover site. *Auk* 125: 95–104.
- Winker, K. (1995) Autumn stopover on the isthmus of Tehuantepec by woodland Nearctic–Neotropical migrants. *Auk* 112: 690–700.
- Winkler, D. W. (2006) Roosts and migrations of swallows. *Hornero* 21: 85–97.
- Wolfe, J. D., Johnson, M. D. and Ralph, C. J. (2014) Do birds select habitat or food resources? Nearctic–Neotropical migrants in northeastern Costa Rica. *PLoS One* 9: e86221.
- Yong, W. and Finch, D. M. (1997) Migration of the Willow Flycatcher along the middle Rio Grande. *Wilson Bull.* 109: 253–268.
- Yong, W., Finch, D. M., Moore, F. R. and Kelly, J. F. (1998) Stopover ecology and habitat use of migratory Wilson's Warblers. *Auk* 115: 829–842.
- Zimmerman, G. M. and Meyer, K. D. (2004) *Migration ecology of Florida's Swallow-tailed Kites in Cuba, Mexico, and Belize*. Tallahassee, Florida: Florida Fish and Wildlife Conservation Commission.

NICHOLAS J. BAYLY*

SELVA: Investigación para la Conservación en el Neotropico, Bogotá D.C., Colombia.

CAMILA GÓMEZ

SELVA: Investigación para la Conservación en el Neotropico, Bogotá D.C., Colombia, and Laboratorio de Biología Evolutiva de Vertebrados, Universidad de los Andes, Bogotá, Colombia.

KENNETH V. ROSENBERG

Cornell Lab of Ornithology, Ithaca, New York, USA, and American Bird Conservancy, The Plains, Virginia, USA.

WENDY E. EASTON

Environment and Climate Change Canada, Canadian Wildlife Service, Delta, Canada.

JAY CARLISLE

Intermountain Bird Observatory, Boise State University, Boise, Idaho, USA.

DAVID N. EWERT

The Nature Conservancy, Lansing, Michigan, USA.

ANNA DRAKE

Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, Canada.

LAURIE GOODRICH

Hawk Mountain Sanctuary Association, Orwigsburg, Pennsylvania.

*Author for correspondence; e-mail: nick.bayly@selva.org.co

Received 25 January 2017; revision accepted 17 July 2017;

Published online 30 October 2017