

## Risk factors for *Escherichia coli* O157 on beef cattle ranches located near a major produce production region

L. A. BENJAMIN<sup>1</sup>, M. T. JAY-RUSSELL<sup>1\*</sup>, E. R. ATWILL<sup>1</sup>, M. B. COOLEY<sup>2</sup>,  
D. CARYCHAO<sup>2</sup>, R. E. LARSEN<sup>3</sup> AND R. E. MANDRELL<sup>2</sup>

<sup>1</sup> Western Center for Food Safety, University of California–Davis, Davis, CA, USA

<sup>2</sup> Produce Safety and Microbiology Research Unit, U.S. Department of Agriculture, Agricultural Research Service, Western Regional Research Center, Albany, CA, USA

<sup>3</sup> University of California Cooperative Extension, Templeton, CA, USA

Received 29 September 2013; Final revision 16 February 2014; Accepted 19 February 2014;  
first published online 20 March 2014

### SUMMARY

Our goal was to identify climate variables and management practices associated with the presence of *E. coli* O157 in rangeland cow-calf operations located in a major leafy green production region in the California Central Coast. *E. coli* O157 was present in 2·6% (68/2654) of faecal, 1·5% (3/204) of water and 1·1% (1/93) of sediment samples collected on eight ranches over 2·5 years. Five (62·5%) ranches were positive at least once during the study. The odds of detecting *E. coli* O157 in faecal samples was higher during periods of higher maximum soil temperature, higher maximum relative humidity, and larger herd sizes, but decreased as wind speed increased. Molecular subtyping of isolates from cattle faeces and streams/sediments suggested minimal movement of strains between ranches. The findings suggest that *E. coli* O157 prevalence is relatively low on cow-calf ranches in this region, spatially constrained, but may vary by weather conditions and herd size.

**Key words:** Cattle, *Escherichia coli* O157, multi-locus variable number tandem repeat analysis, risk factors.

### INTRODUCTION

An increasing number of outbreaks over the past decade have been attributed to the contamination of leafy green vegetables with *E. coli* O157 [1]. Consequently, there is global interest in developing strategies to prevent similar outbreaks from occurring in the future [2]. Most recent leafy green-related *E. coli* O157 outbreaks were traced to fresh spinach and lettuce grown in the California Central Coast [3]. Direct contamination

of produce fields by domestic or wild animal hosts, and indirect contamination of irrigation water by faecal material, have been implicated as potential contributing factors to pre-harvest contamination of leafy greens in California [4–8].

Because beef cattle, or their immediate environment, are believed to be reservoirs of these bacteria [9, 10], it is possible that *E. coli* O157 from beef cattle ranches can be conveyed to produce farms located in the same region and contaminate crops. Reported prevalence of *E. coli* O157 in beef cattle on rangeland and pasture can vary from 0·9% (82/9122) [11] to 18% (9/50) [12]. However, in the absence of diagnostic or environmental testing it is difficult to identify ranches that are contaminated with *E. coli* O157 because most

\* Author for correspondence: Dr M. Jay-Russell, Western Center for Food Safety, University of California–Davis, 1477 Drew Avenue, Ste. 101, Davis, CA 95618, USA.  
(Email address: mjay@ucdavis.edu)

Table 1. Characteristics of California Central Coast beef ranches enrolled in the study

Ranch ID	County	Grazing area (ha), mean (range)	Drinking water source*	Herd size†				
				Adults	Heifers	Unweaned	Weaned/stockers	Total
A	Monterey	54 (10–162)	Spring, ground or surface	35	13	24	0	72
B	Monterey	287 (93–728)	Ground or surface	47	29	0	0	76
C	San Luis Obispo	162 (20–304)	Ground	37	13	18	0	68
D	Monterey	247 (97–364)	Spring, ground or surface	77	19	32	17	145
E	San Benito	325 (8–405)	Spring or surface	67	9	19	6	101
F	San Luis Obispo	1072 (809–1416)	Spring or ground	48	8	54	15	125
G	San Luis Obispo	19 (0.8–34)	Ground or surface	81	5	42	0	128
H	Monterey	144 (12–275)	Ground or surface	124	34	105	0	263

\* Different combinations of spring, ground or surface sources in each pasture sampled on a particular ranch.

† Size of cattle group.

beef cattle do not exhibit clinical signs of disease when shedding these bacteria in their faeces [13].

It is apparent from the limited reports of *E. coli* O157 occurrence in beef cattle on pasture or rangeland that much remains to be learned about relevant risk factors. If we are to improve the effectiveness of strategies designed to minimize the pre-harvest contamination of leafy green vegetables with *E. coli* O157, and incorporate these strategies into Good Agricultural Practices (GAPs) [14], then it is important to determine management and environmental variables that may reduce or elevate the risk of produce contamination.

Our major objective in this study was to identify environmental and management risk factors for faecal shedding of *E. coli* O157 in rangeland beef cattle that could be a source of microbial contamination in a major leafy green vegetable production region of California. A secondary objective was to determine phylogenetic relationships between *E. coli* O157 strains isolated from cattle faeces and nearby surface water and sediment samples for a network of cow-calf ranches located in California's Central Coast produce production region.

## MATERIALS AND METHOD

### Sampling and confidentiality

This report is part of a larger longitudinal study with repeat sampling in a major coastal produce production area of California [15–17]. Eight cow-calf ranches in three California Central Coast counties (Monterey, San Benito, San Luis Obispo) were enrolled in the study. The demographics of the ranches were typical of the region and are shown in Table 1.

The locations of each property were kept confidential by using an 8-digit alphanumeric code to blind the laboratory to sample location. According to the confidentiality agreement with the ranch owners, maps showing the location of positive samples are not included in this paper.

Ranches were sampled at least quarterly between 16 June 2008 and 26 October 2010. Approximately 30–35 faecal samples were collected from individual animals on each sampling date. Faecal samples were collected per rectum, when feasible or from the interior of freshly deposited faeces (no visible crust) on grazed or cattle loafing areas using a sterile tongue depressor. Additionally, surface water and sediment samples were collected from troughs, streams, ponds, and dip-tanks (insecticide) in the cattle pastures. Water grab samples of up to 290 ml were placed in sterile bottles, and Moore swab samples were collected from streams as described previously [15]. Sediment samples were collected from surface water sources by using a sterile scoop to scrape ~10 g material into a sterile whirlpak bag. Faecal, water and sediment samples were shipped overnight on ice to the laboratory.

### Culture and identification

Samples were cultured for *E. coli* O157 and generic *E. coli* using techniques described previously [15, 18]. Briefly, for *E. coli* O157 isolation, 90 ml tryptose soy broth (TSB, Becton Dickinson, USA) were added to 10 g faeces or sediment samples, or 11 ml 10×TSB were added to 100 ml water grab samples; Moore swabs were added to 250 ml TSB. Samples were enriched in TSB and then incubated overnight at 4 °C. For detection of *E. coli* O157, 20 µl anti-O157

antibody bound to magnetic beads (Invitrogen/Dynal, USA) were added to 1 ml enrichment broth and mixed for 30 min. Fifty microlitres of re-suspended beads were spread on Sorbitol MacConkey agar (Difco Laboratories, USA) with cefixime (0.05 µg/ml, Invitrogen/Dynal) and tellurite (2.5 µg/ml, Invitrogen/Dynal) (CT-SMAC), Rainbow Agar (Biolog, USA) containing novobiocin (20 µg/ml, Sigma-Aldrich) and tellurite (0.8 µg/ml, Invitrogen/Dynal) (NT-Rainbow) [19]. The CT-SMAC and NT-Rainbow plates were then incubated for 24 h at 37 °C. Suspected *E. coli* O157 colonies on CT-SMAC and NT-Rainbow plates were transferred into wells containing reagents for RT-PCR for the presence of the *rfbE* gene for O157 [15]. To determine the concentration of generic *E. coli* in 100 ml water grab samples or 100 ml rinsate from Moore swabs, samples were incubated at 37 °C for 24 h in sealed QuantiTray 2000 with Colilert reagent (IDEXX Laboratories, USA). The most probable number of *E. coli* per 100 ml grab or swab rinsate sample was determined based on the number of fluorescent wells according to the manufacturer's instructions. One millilitre each of tenfold serial dilutions made from 1 ml of a mixture consisting of 10 g sediment and 90 ml TSB were plated on *E. coli*/Coliform Petrifilm (3M Corp., St Paul, USA) and incubated at 37 °C for 24 h. Generic *E. coli* colonies from sediment were recorded as the number of blue colonies with and without gas bubbles.

#### Multi-locus variable number tandem repeat (VNTR) analysis (MLVA)

Phylogenetic relationships between strains were compared using an 11-loci MLVA as described previously [15, 18]. Briefly, amplified fragments were size-fractionated by an ABI 3130 sequencer (Applied Biosystems, USA) and assigned allele numbers. Minimal spanning trees were constructed from MLVA data using BioNumerics v. 6.0 (Sint-Martens-Latem, Belgium). Three outbreak-related *E. coli* O157 isolates were included in the phylogenetic tree for comparison as described previously [15]: a human strain associated with an outbreak in 2006 due to spinach from the California Central Coast ('Spinach'), a human strain associated with an outbreak in 1996 due to consumption of white radish sprouts in Sakai, Japan ('Sakai'), and a strain isolated from Michigan ground beef linked to a multi-state outbreak due to hamburger ('EDL933').

#### Data collection

For each sampling date, climate data were accessed from the University of California Online Statewide Pest Management Program's weather station closest to the sample location [20]. In addition, questionnaires were administered to ranchers to collect information on relevant management practices.


#### Statistical analysis

Herd and animal prevalence were calculated for *E. coli* O157. Mean, standard deviation and range were calculated for generic *E. coli* in water. Climate and management variables were either categorical or continuous (Fig. 1).

Climate data included minimum and maximum air temperature (°C), soil temperature (°C), relative humidity (%), as well as the amount of rain in the past 24 h (mm) and wind speed (m/s). The variable 'season' (dry = 1 April to 30 November or wet = 1 December to 31 March) was calculated from the recorded sample collection date.

Continuous management variables included the number of days that the current herd had spent on the sampled pasture prior to the date on which the sample was collected; the total number of adults, heifers and calves on the sampled pasture (unweaned and weaned) (Table 1); cattle density (total cattle of all age classes/size of sampled pasture in hectares); length of the calving season for cows; number of females purchased in the past 2 years and the number of bulls used in the most recent breeding season. Categorical management variables were source of water (0 = includes ground water, 1 = does not include ground water); whether heifers were 2 or 3 years old when they calve for the first time (0 = 2 years old and 1 = 3 years old); whether or not females were regularly purchased (no/yes); whether or not heifer/calf pair was moved from the calving pasture (no/yes); whether or not cow/calf pair was moved from the calving pasture (no/yes) and source of replacement cattle (0 = market or auction and 1 = private sales).

Firthlogit was used to test univariate associations between climate and management variables and the dichotomous outcome of *E. coli* O157 (no/yes) in cattle faeces and to construct two models: climate variables and management variables ( $P \leq 0.05$ ). A forward-stepping algorithm was used to build a final logistic regression model with farm identification as the group and using variables from the climate and



**Fig. 1.** Risk factors considered for the multivariable logistic regression model of *E. coli* O157 on beef cattle ranches on the California Central Coast.

management models [21]. A logistic regression model was used also to test the association between *E. coli* O157 and generic *E. coli* in cattle faeces ( $P \leq 0.05$ ).

### Software

Laboratory and covariate data were recorded on separate datasheets in Microsoft Excel (Microsoft Corporation, USA) and merged in Microsoft Access (Microsoft Corporation). Merged *E. coli* O157 data were analysed using Stata (StataCorp, USA) and LogXact (version 4.0.4, Cytel Software Corporation, USA).

## RESULTS

### Prevalence of *E. coli* O157

Overall, *E. coli* O157 was present in 2.6% (68/2654) of faecal, 1.5% (3/204) of water and 1.1% (1/93) of sediment samples collected between 16 June 2008 and 26 October 2010 at eight cattle ranches (Table 2). Five

(62.5%) of the eight ranches were positive at least once in the study (Table 3). Proportions of positive faecal samples ranged from 0.0% on ranches A, C and H to 10.1% on ranch D. Three positive surface water and one positive sediment sample were found in creeks or streams on ranches B and G, but all water trough samples were negative (Table 2). The proportion of *E. coli* O157 isolated from faeces was slightly higher than from water and sediment, but the difference was not significant at ranch B ( $P > 0.05$ ).

*E. coli* O157 was not isolated in any sample during 2008 (Table 2), and detection during 2009 and 2010 was limited to only a few sampling dates at ranches B, D, E, F, and G (Table 3). An unusual spike in prevalence occurred at ranch D on 23 September 2009 (Fig. 2) when *E. coli* O157 was isolated from 44/49 (89.8%) of cattle faecal samples; three water samples collected that same date were negative. Of note, this was the only detection of *E. coli* O157 in cattle or water/sediment at ranch D except a single feral pig hunted on the ranch on 21 July 2009.

Table 2. Proportions of *E. coli* O157 isolated from samples from California Central Coast beef ranches between 16 July 2008 and 26 October 2010

Source	Year	No. of samples tested	No. of samples positive (%)	No. of sampling days*	No. of positive sampling days (%)
Faeces	2008	398	0 (0)	7	0 (0.0)
	2009	934	49 (5.3)	19	2 (10.5)
	2010	1322	19 (1.4)	23	4 (17.4)
	All	2654	68 (2.6)	49	6 (12.2)
Water	2008	56	0 (0)	8	0 (0.0)
	2009	90	1 (1.1)	26	1 (3.8)
	2010	58	2 (3.5)	23	2 (8.7)
	All	204	3 (1.5)	57	3 (5.3)
Sediment	2008	30	0 (0)	7	0 (0.0)
	2009	50	1 (2.0)	17	1 (5.9)
	2010	13	0 (0)	7	0 (0.0)
	All	93	1 (1.1)	31	1 (3.2)

\* Number of dates samples were collected through the duration of the study.

Table 3. Proportions of cattle fecal samples positive for *E. coli* O157 from California Central Coast beef ranches between June 16, 2008 to October 26, 2010


Farm ID	No. of samples tested	No. of samples positive (%)	No. of sampling days*	Date of positive samples
A	489	0 (0.0)	10	None
B	480	7 (1.5)	8	2 Sept. 2010
C	200	0 (0.0)	4	None
D	434	44 (10.1)	10	24 Sept. 2009
E	386	6 (1.6)	8	3 Dec. 2009, 25 Mar. 2010
F	271	2 (0.7)	5	14 Oct. 2010
G	256	9 (3.5)	5	24 Apr. 2010
H	138	0 (0.0)	2	None
Total	2654	68 (2.6)		

\* Number of dates samples were collected through the duration of the study.

### MLVA typing

A total of 141 isolates from 71 samples (faeces, water, sediment) represented 36 MLVA types. Cattle faecal *E. coli* O157 isolates from five positive ranches clustered into six groups with no overlap between ranches (Fig. 3). Isolates from the San Benito County ranch (E) comprised two clusters. Faecal isolates from ranch D were all cultured from a single sampling date (23 September 2009) as described above, and these subtypes were not found again in cattle faeces from ranch D or other cow/calf ranches. Nine closely related MLVA types were detected in cattle faecal isolates with 74/89 (83.1%) representing two dominant

types (Fig. 3). The isolate from the feral pig sampled from ranch D 2 months prior to the spike in herd prevalence in September 2009 was identical to the second most dominant MLVA type detected in cattle faeces. Stream water isolates from ranch G were identical or very closely related to cattle faecal samples collected on the same date (23 April 2010). By contrast, stream water and sediment samples collected on ranch B in March 2009 were more closely related to the outbreak-related isolates EDL933 and Sakai than other isolates collected from ranch B. The 2006 spinach-related outbreak strain was not detected in any of the ranch samples, although one isolate collected from water from ranch G was closely related (Fig. 3).



**Fig. 2.** Percent positive *E. coli* O157 isolated from cattle faeces, water and sediment between 16 June 2008 and 26 October 2010 from samples collected on California Central Coast beef ranches.

### Management practices


Mean number of adult cattle grazing in sampled pastures was 60 (range 14–125). The mean (s.d.) cattle density was 6.2 ( $\pm 27.1$ ) head of cattle per hectare. Sampled cattle spent a mean of 59.3 days on the current pasture prior to sampling. Four ranches reported using all three of the following methods to replace females over the 2 years of sampling: home-grown replacements, purchasing from a market and private agreement. On one ranch (ranch D), female replacements were obtained only from the ranch itself. Producers reported a mean of 30 (range 10–50) females introduced to their ranch in the 2 years prior to the sample collection date. Eighty-six percent (6/7) and 71.4% (5/7) of ranches reported introducing cows and bulls from public auctions, respectively. Fifty-seven percent (4/7) of ranches moved cow-calf pairs away from pregnant cows in the herd following calving. Only one ranch reported on one occasion that the sampled pasture had been irrigated 30 days prior to sampling. Pastures were comprised mainly of introduced annual grasses and forbs. Most pastures

were in or near oak woodland or chaparral vegetation types.

### Generic *E. coli*

Water troughs were used on all eight ranches. No *E. coli* O157 was isolated from troughs. In troughs, the mean generic *E. coli* isolated from water,  $2.1 \times 10^3$ , was lower than that from sediment samples,  $8.1 \times 10^6$  ( $P \leq 0.05$ ). However, fewer trough sediment samples ( $n=7$ ) were collected compared to water samples ( $n=70$ ). Mean generic *E. coli* of water from troughs supplied by wells,  $3.7 \times 10^3$  (38 samples), was not significantly different from that of troughs filled using surface water,  $1.4 \times 10^2$  (27 samples) ( $P > 0.05$ ).

For the months in which both water and sediment samples were collected from sources other than troughs, the mean generic *E. coli* in water was higher in streams/creeks in April and August 2009 and for April 2010 ( $P \leq 0.05$ ) compared to other months. The association between *E. coli* O157 positivity and



**Fig. 3** [colour online]. Phylogeny of *E. coli* O157 isolates from cattle ranches on the California Central Coast determined by 11-loci MLVA. A minimal spanning tree was constructed of 36 MLVA types representing the O157 STEC strains isolated from 71 samples. Node size indicates the relative number of isolates of that type; i.e. the smallest size node represents a single strain of that type. The nodes are colour-coded by farm/ranch site code (left legend) and by sample source (right legend). Outbreak-related strains (Sakai, Spinach, EDL933) are included for comparison.

concentration of generic *E. coli* (c.f.u./100 ml) in faeces was not significant ( $P > 0.05$ ).

**Univariate analysis of *E. coli* O157 in faeces**

The only variables with non-significant univariate associations with the outcome ( $P > 0.2$ ) were days on current pasture, cattle density (herd size/size of pasture in hectares), and number of bulls used in the most recent calving (Table 4).

**Separate climate and management models**

Variables in the final climate model were maximum soil temperature, maximum air temperature, maximum relative humidity, wind speed and precipitation (Table 5). The final management model included herd size, percent unweaned cattle, length of calving season, number of replacement females purchased, moved cow/calf pair after calving, and water source (Table 5).

**Multivariate analysis of *E. coli* O157 in faeces**

Variables from the climate and management models were used to build the final model (Table 5).

Exponentiation of the coefficients of the variables in the final logistic regression model for *E. coli* O157 isolated from faecal samples yields the odds. For example, in the present study, increasing the maximum relative humidity by 1% increased the odds of being *E. coli* O157 positive by  $e^{0.26} = 1.29$ . Increasing the herd size by 10 head of cattle increased the odds of being *E. coli* O157 positive by a factor of  $e^{0.08 \times 10} = 2.23$ .

**DISCUSSION**

Our study differs from previous reports of cattle on pasture or rangeland because it is a longitudinal risk factor analysis of *E. coli* O157 conducted on multiple ranches located on the California Central Coast, a region with high concentration of leafy green produce farms. As a result, it addresses a gap in the data and provides information that can be used in environmental assessments of potential risks posed by beef cattle ranches in proximity to fresh leafy green vegetable fields in this region.

**Prevalence**

The low mean animal-level prevalence of *E. coli* O157 in our study is consistent with previous studies of beef herds [22]. The proportion of herds that were positive at least once over the course of the study was much higher than the herd-level prevalence of cow-calf herds in Louisiana, 17.2% (5/29). However, it should be kept in mind that our study involved a smaller number of herds.

Interestingly, a survey of *Salmonella enterica* in a subset of the same samples in this study and a nearby feedlot revealed an extremely low prevalence (1/794, 0.1%) of that pathogen [17].

Prior to this study, data on *E. coli* O157 occurrence in cattle in the California Central Coast came primarily from testing conducted during outbreak investigations. Specifically, three cattle ranches near implicated spinach fields were tested in autumn 2006 in Monterey and San Benito counties following a nationwide outbreak of *E. coli* O157 associated with bagged baby spinach [4, 8, 18]. *E. coli* O157 was isolated from cattle faeces and/or pasture soil on all three ranches, but the ‘outbreak strain’ was only found in samples from a single ranch in San Benito County [4]. Interestingly, the positive cattle faecal samples came from a grass-fed cow-calf herd on rangeland and irrigated pasture, and the percent positive (26/77 faecal

Table 4. *Univariate analysis of factors associated with the presence of E. coli O157 in fecal samples collected on California Central Coast beef ranches between June 16, 2008 and October 26, 2010*

Model	Variable	Level	$\beta$ (coefficient)	95% CI ( $\beta$ )	<i>P</i>	
Climate	Minimum air temperature (°C)		0.21	0.14 to 0.28	<0.01	
	Maximum air temperature (°C)		0.08	0.05 to 0.11	<0.01	
	Minimum soil temperature (°C)		0.31	0.23 to 0.40	<0.01	
	Maximum soil temperature (°C)		0.29	0.21 to 0.36	<0.01	
	Maximum relative humidity (%)		0.42	0.32 to 0.52	<0.01	
	Minimum relative humidity (%)		0.02	0.01 to 0.04	<0.01	
	Precipitation (mm)		-4.15	-8.88 to 0.58	0.09	
	Wind speed (m/s)		0.28	0.04 to 0.51	0.02	
	Season	Dry (April–November)	Reference group			
	Wet (December–March)		-1.92	-2.73 to -1.11	<0.01	
Management	Days on current pasture (day)		-0.002	-0.01 to 0.001	0.21	
	Herd size (head cattle)		0.01	0.01 to 0.02	<0.01	
	Cattle density (herd size/size of pasture in acres)		-0.007	-0.02 to 0.008	0.37	
	No. of females purchased in the past 2 years		-0.17	-0.35 to -0.001	0.05	
	No. of bulls used in the most recent breeding season		-0.02	-0.07 to 0.02	0.27	
	Length of calving season for cows		-0.04	-0.06 to -0.01	0.01	
	Water source	Includes ground water	Reference group			
		Does not include ground water		1.76	1.20 to 2.32	<0.01
	Age of heifer at first calving	2 years old	Reference group			
		3 years old		-3.22	-6.01 to -0.44	0.02
	Regularly purchase females	No	Reference group			
	Yes		-3.79	-6.58 to -1.00	<0.01	

CI, Confidence interval.

Table 5. Climate (only), management (only) and management and climate final multivariable logistic regression models for associations between variables and *E. coli* O157 isolated from faecal samples collected on California Central Coast beef ranches between 16 June 2008 and 26 October 2010

Models	Variable	Level	$\beta$ (coefficient)*	95% CI ( $\beta$ )	P
Climate only	Maximum soil temperature		0.34	0.19 to 0.49	<0.01
	Maximum temperature		0.18	0.10 to 0.26	<0.01
	Maximum relative humidity		0.53	0.43 to 0.64	<0.01
	Wind speed		-1.62	-2.47 to -0.77	<0.01
	Precipitation		-13.47	-20.13 to -6.80	<0.01
	Constant		-60.79	-71.79 to -49.78	<0.01
Management only	Herd size		0.05	0.01 to 0.08	0.01
	Unweaned percent of cattle		0.12	0.05 to 0.18	<0.01
	Length of calving season		-0.20	-0.29 to -0.11	<0.01
	No. of replacement females purchased		0.35	0.10 to 0.60	<0.01
	Moved cow/calf pair†	0	Reference group		
		1	8.20	5.20 to 11.19	<0.01
	Water source‡	0	Reference group		
		1	7.42	4.40 to 10.45	<0.01
	Constant		-19.24	-27.33 to -11.15	<0.01
Climate and management	Maximum 24-h soil temperature (°C)		-1.97	-3.21 to -0.72	<0.01
	Maximum 24-h soil temperature (°C) squared		0.07	0.03 to 0.10	<0.01
	Maximum relative humidity		0.26	0.11 to 0.42	<0.01
	Wind speed		-5.20	-7.39 to -3.00	<0.01
	Herd size		0.08	0.04 to 0.13	<0.01
	Constant		-17.97	-38.16 to 2.21	0.08

CI, Confidence interval.

\* The coefficient values are partial values given that the other coefficients are in the model. Coefficients are interpreted by exponentiation. Increasing the herd size by 10 head of cattle increased the odds of being *E. coli* O157 positive by a factor of  $e^{0.05 \times 10} = 1.65$  or by  $100(e^{0.05 \times 10} - 1) = 65\%$ . Increasing the percentage of unweaned cattle in the herd by 10% increased the odds of being *E. coli* O157 positive by  $e^{0.12 \times 10} = 3.3$ . Increasing the length of the calving season by 10 days decreased the odds of being *E. coli* O157 positive by a factor of  $e^{-0.20 \times 10} = 0.14$ . Increasing the number of females purchased by 1 increased the odds of being *E. coli* O157 positive by a factor of  $e^{0.35} = 1.42$ .

† A very large odds ratio,  $e^{8.2} = 3625$ , was observed for the variable 'moved cow/calf pair to a different pasture after calving'. This can be attributed to the observation that for all *E. coli* O157-positive cases, the cow/calf pair was moved after calving, that is, there were no positive cases in which the cow/calf pair was not moved after calving. This is referred to as separation and can be observed as a zero cell in the  $2 \times 2$  table for *E. coli* O157 and the variable 'moved the cow/calf pair after calving'. Firth-penalized likelihood logistic regression was used to address separation in the data.

‡ A very large odds ratio was also observed for the variable for water source (OR 1675) on adding moved cow/calf pair after calving to the model. There were no positive cases where the cow/calf pair was moved after calving and the water source did or did not include ground water. Despite their extremely large odds ratios, the variables 'moved cow/calf pair' and 'water source' were left in the model because the goal of this model was to identify variables that might be useful in the combined final model with both management and climate variables.

samples, 33.8%) [4, 8] was much higher compared to the combined prevalence (68/2654, 2.6%) found in the survey described here. Culture bias could explain some of the discrepancy, but there were only minor variations in sampling and testing methodologies between the two studies. Alternatively, environmental (e.g. weather, wildlife movement) or management practices specific to the ranch implicated in 2006 may have contributed to the relatively high recovery

of *E. coli* O157 from cow-calf faecal samples following the spinach outbreak.

It is also possible that undetermined factors may cause sudden prevalence increases in cow-calf beef herds. For example, we observed a single peak in herd prevalence (44/49, 89.8%) in September 2009 (Fig. 2) at ranch D involving closely related MLVA types (Fig. 3). The lack of recovery of *E. coli* O157 in water/sediment samples on ranch D suggests limited

environmental dissemination of the bacteria or exposure via cattle drinking-water sources. The detection of an identical *E. coli* O157 MLVA type from a feral pig hunted on ranch D 2 months prior to the spike in cattle herd prevalence is intriguing and could suggest introduction of the pathogen by free-roaming wildlife. Additional research is needed to better characterize livestock–wildlife interactions, and identify risk factors that induce fluctuations in herd prevalence; such information would allow ranchers and growers to potentially reduce dissemination of *E. coli* O157 to nearby produce fields.

We also speculate that the spike in prevalence observed in autumn 2009 (Fig. 2) could be related to reduced forage quality during the driest part of the year. Vegetation in these pastures mostly consists of introduced annual grasses that germinate with the onset of the rainfall season in autumn, grow slowly throughout the cool moist winter, and then exhibit rapid growth and senescence during spring. The vegetation is dry and of lower nutritional value during summer and autumn. Of note, forage production was highly variable throughout the 2.5-year study, with 2008 being about average production, 2009 being very low production, and 2010 being above average production [23–26]. The dry season in the California Central Coast is also the time when the majority of lettuce, spinach, and other leafy green vegetables are grown and harvested in this region.

#### Multivariable model with both climate and management factors

Temperature, season, and precipitation are environmental factors suspected to be associated with the prevalence of *E. coli* O157 in beef cattle [27]. In our study, the odds of being *E. coli* O157 positive increased as the 24 h maximum soil temperature increased from 21 °C to 26.1 °C. The odds of a sample being *E. coli* O157 positive also increased as the maximum relative humidity increased. In an experimental study of epiphytic bacteria at different temperatures (15 °C and 30 °C) and humidity, it was reported that half-lives were shorter at low humidity ( $\leq 50\%$ ), 14 and 3 min, than at high humidity ( $>50\%$ ), 83 and 14 min [28]. Both temperature and humidity are believed to exert their effects by prolonging the survival of *E. coli* O157 in faecal pats on pasture soil or grass. Longer survival increases the period during which it is possible for susceptible cattle to be exposed to these bacteria.

Wind speed was negatively associated with the occurrence of *E. coli* O157 in faecal pats ( $P \leq 0.05$ ) in the present study. The association was the opposite of that for feedlot cattle in Midwestern USA, where wind velocity was positively associated with *E. coli* O157 [29]. It is possible that cattle are more likely to be exposed to airborne bacteria in the typical feedlot because it is a more confined environment with a higher density of animals in close proximity to a higher concentration of *E. coli* O157 present in often dry soil. However, on pasture/rangeland the wind may have had the effect of dispersing the already low concentration of *E. coli* O157 available for exposing cattle, or windier days may lead to drier faecal conditions and reduce survival of this bacterial pathogen.

There was a positive association between herd size (on sampled pasture) and *E. coli* O157, as was reported for UK cattle between the ages of 5 and 28 months (odds ratio 1.16, 95% confidence interval 1.10–1.34) [30]. The mean herd size of ranches participating in this study was similar to the average size of a beef herd in California, 55.5 head of cattle [31]. It is interesting that cattle density was not significantly associated with the outcome in univariate ( $P > 0.2$ ) or multivariate ( $P > 0.05$ ) analysis. This implies that stocking density might not be an important risk factor for cattle on range/pasture given the tendency of cattle to form into a loose spatial group regardless of pasture size, hence, stocking density is not a valid proxy for the spatial density of cattle on rangeland.

#### Separate climate and management models

Maximum air temperature and precipitation were included in the climate model (Table 4) but not in the final model (Table 5). This is in agreement with prior work which found that prevalence was lowest in the winter (low temperature) in a rangeland herd in the Sierra Nevada Mountains in California, with prevalence increasing as the air temperature increased from moderate to high [32]. It is possible that higher temperatures can lead to clustering of cattle as they seek shade and windier locations and as a result susceptible animals are more likely to come into contact with *E. coli* O157 present in faecal pats deposited by other animals in the herd.

Water can transport bacteria from one ranch to another if suitable conditions exist. However, the low prevalence of *E. coli* O157 in water sources and sediment found during this study suggested water

and sediment were not playing a major role in the spread of this pathogen at the time we sampled. In a previous study in the same region, a higher incidence of *E. coli* O157 in the Salinas watershed was observed following heavy rainfall [18], indicating a possible route for transfer of this pathogen to downstream produce operations [33]. The risk from cattle faecal contamination of water sources will ultimately depend on connectivity, slope, proximity and whether or not cattle shedding pathogenic bacteria have direct access to water sources. It has been proposed that bacteria deposited in surface waters settles into the sediment during low flows and becomes re-suspended during high flows [34].

Season was significant in the univariate analysis ( $P \leq 0.2$ ) but not in the final model. It is difficult to compare the effect of season from previous reports because studies were conducted in different climates, some with more extreme temperature variations than are experienced on the California Central Coast. Similar to our findings, there no significant association between season and the prevalence of *E. coli* O157 in beef cattle in Kansas [35]. It is believed that the seasonal effect observed in some studies might be related to other variables such as temperature and precipitation.

In the separate management model (Table 5) the variables of percent unweaned cattle, length of calving season, number of replacement females purchased, moved calf/heifer pair after calving, and water source, were also significantly associated with the outcome. Some previous studies [36] identified lower *E. coli* in unweaned compared to weaned cattle. However, researchers did not identify a difference in *E. coli* O157 prevalence in different age groups of cattle housed on pasture in a Kansas study [22] and over the course of a 45-day preconditioning period, animal-level prevalence of *E. coli* O157 declined from 2.5% to 0.0% in commingled weaned calves originating from 29 different farms [37]. Because most of the faecal samples in this study were collected from rangeland pastures and not individual animals, age and gender were not recorded, thus we were not able to evaluate these covariates.

We also found that moving the cow-calf pair after calving was associated with higher odds of *E. coli* O157. In a previous report, for 2 consecutive years, the prevalence of *E. coli* O157 in faecal samples from cow-calf pairs was higher for cattle recently placed on pasture at 12 weeks post-calving than for the same cattle 5 weeks later [12]. In addition, these calves were not consistently observed to shed *E. coli*

O157 during the same time period that their dams were shedding, suggesting that horizontal transmission of bacteria was important for calf infections. Practices such as moving the cow-calf pair are facilitated by a well-defined (shorter) calving season. In our study, an increase in the length of the calving season for cows had a protective effect. This is contrary to expectations because a shorter calving season is expected to allow for more effective herd management.

Introduction of new stock is another potential management factor that could influence herd prevalence especially if new cattle are sourced from markets or auctions. Colonized newly purchased animals can introduce pathogenic bacteria to previously uninfected herds. Similar to our findings, Synge *et al.* [27] reported that changes in the number of cows in Scottish beef suckler herds were associated with increased odds of *E. coli* O157. It is more likely that there will be more changes in the number of cattle in larger than in smaller herds, with increased risk of the introduction of infected animals, especially where cattle are acquired from multiple sources. Likewise, molecular subtyping of isolates from cattle faeces from the herds in this study showed no movement of strains between these ranches (Fig. 3), which is consistent with closed herd management practices or minimal introduction of new stock at these locations.

In the management-variables-only model (Table 5), the odds of a positive *E. coli* O157 sample were higher if the water source did not include ground water. The prevalence of *E. coli* O157 in water samples in this study, 1.5%, was the same as that reported for ten cow-calf ranches in Kansas [12]. *E. coli* O157 was also isolated from stream samples on cattle ranches in previous reports [22]. A combination of molecular and longitudinal studies with spatial analysis should provide the necessary data to map the direction of movement of *E. coli* O157 between the environment and cattle on ranches. Previous studies investigated the possible role of trough water as a reservoir of pathogenic *E. coli* for cattle herds [10]. Moreover, *E. coli* counts have been reported in other studies to be higher in cattle troughs from which *E. coli* O157 was isolated in previous studies, but this observation could not be tested in our study because *E. coli* O157 was not isolated from cattle troughs.

It is important to note some limitations to our model. Specifically, variables in the management model were affected by separation issues. Separation

or sparseness occurs when there are few to no cases within a category of the variable. When separation is present, inflated standard errors might be observed or confidence limits might not exist [38]. Firthlogit regression was used to address separation issues in the data. Firth's penalized likelihood produces finite parameter estimates and unlike exact logistic regression, it does not result in degenerate outcomes when continuous variables are presented to the model [39]. A simpler solution to the separation issue, involving re-categorization of data into meaningful categories that are less sparse, was also employed in this analysis. The simplest approach to dealing with separation is to omit the variables in question from the analysis. However, with this simple approach the possibility that the omitted variable might be important in predicting the outcome is not taken into consideration and the effects of the remaining variables in the model will not be adjusted for the effect of the omitted variable.

## CONCLUSIONS

Prevalence of *E. coli* O157 was low in the rangeland beef herds sampled in this study, although spikes in herd prevalence were observed at individual ranches. Variables associated with increased *E. coli* O157 on cattle ranches were maximum soil temperature, relative maximum humidity, and herd size on sampled pastures. Molecular epidemiological inferences were that isolates were spatially clustered by ranch with instances of the same or highly related strains being found in both cattle and water/sediment samples. The findings underscore the need for pre-harvest interventions for cattle to prevent environmental dissemination of *E. coli* O157 and reduce the risk of microbial contamination of adjacent fresh produce fields. For example, *E. coli* O157 cattle vaccination represents a promising pre-harvest approach, although questions remain regarding the economic feasibility of vaccinating cattle herds against *E. coli* O157 since the pathogen is a primarily public health, not animal health, concern [40].

## ACKNOWLEDGEMENTS

This research was funded by the U.S. Food and Drug Administration Project No. U01-003-572; by National Research Initiative Competitive Grant nos. 2006-55212-16927 and 2007-35212-18239 from the U.S. Department of Agriculture National Institute

of Food and Agriculture; and the U.S. Department of Agriculture, Agricultural Research Service, CRIS project number 5325-42000-044-00D. This project was also supported in part by an appointment to the Research Participation at the Center for Food Safety and Applied Nutrition administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the U.S. Food and Drug Administration.

Thanks are due to Stefanie Hee and Geri Kavanagh, field support staff, for their efforts, and to Dr Eduardo Vivas for his field advice.

## DECLARATION OF INTEREST

None.

## REFERENCES

1. **Berger CN, et al.** Fresh fruit and vegetables as vehicles for the transmission of human pathogens. *Environmental Microbiology* 2010; **12**: 2385–2397.
2. **GLOBAL G.A.P.** Integrated farm assurance: fruit and vegetables, version 4. Cologne, Germany: GLOBAL G.A.P Secretariat, 2012.
3. **Mandrell RE.** Enteric human pathogens associated with fresh produce: sources, transport, and ecology. In: Fong X, Niemira BA, Doona CJ, Feeherry FE, Gravani RB, eds. *Microbial Safety of Fresh Produce*. Blackwell Publishing: Ames, Iowa, 2009, pp. 5–41.
4. **California Department of Public Health.** Investigation of an *Escherichia coli* O157:H7 outbreak associated with Dole pre-packaged spinach 2007, 2007, pp. 1–51.
5. **California Department of Public Health.** *E. coli* O157:H7 outbreak associated with iceberg lettuce at Taco John's, December 2006, 2008, pp. 1–41.
6. **Gelting RJ, et al.** Irrigation water issues potentially related to the 2006 multistate *E. coli* O157:H7 outbreak associated with spinach. *Agricultural Water Management* 2011; **98**: 1395–1402.
7. **Hilborn ED, et al.** A multistate outbreak of *Escherichia coli* O157:H7 infections associated with consumption of mesclun lettuce. *Archives of Internal Medicine* 1999; **159**: 1758–1764.
8. **Jay MT, et al.** *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle, central California coast. *Emerging and Infectious Diseases* 2007; **13**: 1908–1911.
9. **Chapman PA, et al.** An outbreak of infection due to verocytotoxin-producing *Escherichia coli* O157 in four families: the influence of laboratory methods on the outcome of the investigation. *Epidemiology and Infection* 1997; **119**: 113–119.
10. **LeJeune JT, Besser TE, Hancock DD.** Cattle water troughs as reservoirs of *Escherichia coli* O157. *Applied and Environmental Microbiology* 2001; **67**: 3053–3057.

11. **Renter DG, Sargeant JM, Hungerford LL.** Distribution of *Escherichia coli* O157:H7 within and among cattle operations in pasture-based agricultural areas. *American Journal of Veterinary Research* 2004; **65**: 1367–1376.
12. **Gannon VPJ, et al.** *Escherichia coli* O157:H7 infection in cows and calves in a beef cattle herd in Alberta, Canada. *Epidemiology and Infection* 2002; **129**: 163–172.
13. **Cray Jr. WC, Moon HW.** Experimental infection of calves and adult cattle with *Escherichia coli* O157:H7. *Applied and Environmental Microbiology* 1995; **61**: 1586–1590.
14. **California Leafy Green Marketing Agreement.** 2013. Commodity specific food safety guidelines for the production and harvest of lettuce and leafy greens. (<http://www.caleafygreens.ca.gov/sites/default/files/California%20LGMA%20metrics%2008%2026%2013%20%20Final.pdf>). Accessed 9 September 2013.
15. **Cooley MB, et al.** Development of a robust method for isolation of shiga toxin-positive *Escherichia coli* (STEC) from fecal, plant, soil and water samples from a leafy greens production region in California. *PLoS One* 2013; **8**: e65716.
16. **Benjamin L, et al.** Occurrence of generic *Escherichia coli*, *E. coli* O157 and *Salmonella* spp. in water and sediment from leafy green produce farms and streams on the Central California coast. *International Journal of Food Microbiology* 2013; **165**: 65–76.
17. **Gorski L, et al.** Prevalence, distribution, and diversity of *Salmonella enterica* in a major produce region of California. *Applied and Environmental Microbiology* 2011; **77**: 2734–2748.
18. **Cooley M, et al.** Incidence and tracking of *Escherichia coli* O157:H7 in a major produce production region in California. *PLoS One* 2007; **2**: e1159.
19. **Bosilevac JM, Koohmaraie M.** Prevalence and characterization of non-O157 Shiga toxin-producing *Escherichia coli* isolates from commercial ground beef in the United States. *Applied and Environmental Microbiology* 2011; **77**: 2103–2112.
20. **University of California Agriculture & Natural Resources.** Weather, models & degree-days (<http://www.ipm.ucdavis.edu/WEATHER/index.html>). Accessed 6 June, 2012.
21. **Lofstedt J, Dohoo IR, Duizer G.** Model to predict septicemia in diarrheic calves. *Journal of Veterinary Internal Medicine* 1999; **13**: 81–88.
22. **Sargeant JM, et al.** Results of a longitudinal study of the prevalence of *Escherichia coli* O157:H7 on cow-calf farms. *American Journal of Veterinary Research* 2000; **61**: 1375–1379.
23. **Larsen R.** Droughts: a challenge to ranching in oak woodlands. Oak conservation. (<http://ucanr.edu/blogs/blogcore/postdetail.cfm?postnum=7002>). Accessed 7 March 2014.
24. **San Luis Obispo County.** Weights and measures. Weeding out our pests (Annual Report). San Luis Obispo County, 2009.
25. **San Luis Obispo County.** Weights and measures. Fresh from the field (Annual Report). San Luis Obispo County, 2008.
26. **San Luis Obispo County.** Weights and measures. Reflections (Annual Report). San Luis Obispo County, 2010.
27. **Synge BA, et al.** Factors influencing the shedding of verocytotoxin-producing *Escherichia coli* O157 by beef suckler cows. *Epidemiology and Infection* 2003; **130**: 301–312.
28. **Wathes CM, Howard K, Webster AJ.** The survival of *Escherichia coli* in an aerosol at air temperatures of 15 and 30 degrees C and a range of humidities. *Journal of Hygiene (London)* 1986; **97**: 489–496.
29. **Sargeant JM, et al.** Associations between management, climate, and *Escherichia coli* O157 in the faeces of feedlot cattle in the Midwestern USA. *Preventive Veterinary Medicine* 2004; **66**: 175–206.
30. **Ellis-Iversen J, et al.** Temporal patterns and risk factors for *Escherichia coli* O157 and *Campylobacter* spp. in young cattle. *Journal of Food Protection* 2009; **72**: 490–496.
31. **McBride WD, Matthews K.** The diverse structure and organization of U.S. beef cow-calf farms. U.S. Department of Agriculture, March 2011.
32. **Kondo S, et al.** Longitudinal prevalence and molecular typing of *Escherichia coli* O157: H7 by use of multiple-locus variable-number tandem-repeat analysis and pulsed-field gel electrophoresis in fecal samples collected from a range-based herd of beef cattle in California. *American Journal of Veterinary Research* 2010; **71**: 1339–1347.
33. **Walters SP, Thebo AL, Boehm AB.** Impact of urbanization and agriculture on the occurrence of bacterial pathogens and stx genes in coastal waterbodies of central California. *Water Research* (in press).
34. **Sherer BM, et al.** Indicator bacterial survival in stream sediments. *Journal of Environmental Quality* 1992; **21**: 591–595.
35. **Alam MJ, Zurek L.** Seasonal prevalence of *Escherichia coli* O157:H7 in beef cattle feces. *Journal of Food Protection* 2006; **69**: 3018–3020.
36. **Laegreid WW, Elder RO, Keen JE.** Prevalence of *Escherichia coli* O157:H7 in range beef calves at weaning. *Epidemiology and Infection* 1999; **123**: 291–298.
37. **Dunn JR, et al.** *Escherichia coli* O157:H7 in a cohort of weaned, preconditioned range beef calves. *Journal of Food Protection* 2004; **67**: 2391–2396.
38. **Hosmer DW, Lemeshow S.** *Applied Logistic Regression*. New York: Wiley-Interscience, 2000.
39. **Heinze G, Schemper M.** A solution to the problem of separation in logistic regression. *Statistics in Medicine* 2002; **21**: 2409–2419.
40. **Matthews L, et al.** Predicting the public health benefit of vaccinating cattle against *Escherichia coli* O157. *Proceedings of the National Academy of Sciences USA*. Published online: 16 September 2013. doi:10.1073/pnas.1304978110.