

The influence of binarity on the morpho-kinematics of planetary nebulae

David Jones¹, Amy A. Tyndall², Myfanwy Lloyd² and Miguel Santander-García³

¹European Southern Observatory, Alonso de Córdova 3107, Casilla 19001, Santiago, Chile
email: djones@eso.org

²Jodrell Bank Centre for Astrophysics, University of Manchester, Oxford Road, Manchester, M13 9PL, UK

email: amy.tyndall@postgrad.manchester.ac.uk, myfanwy.lloyd@manchester.ac.uk

³Observatorio Astronómico Nacional, Ap 112, 28803 Alcalá de Henares, Spain
email: m.santander@oan.es

Abstract. The role of central star binarity in the shaping of planetary nebulae (PNe) has been the subject of much debate, with single stars believed to be incapable of producing the most highly collimated morphologies. However, observational support for binary-induced shaping has been sadly lacking. Here, we highlight the results of a continuing programme to spatio-kinematically model the morphologies of all PNe known to contain a close binary central star. Spatio-kinematical modelling is imperative for these objects, as it circumvents the degeneracy between morphology and orientation which can adversely affect determinations of morphology based on imaging alone. Furthermore, spatio-kinematical modelling accurately determines the orientation of the nebular shell, allowing the theoretically predicted perpendicular alignment, between nebular symmetry axis and binary orbital plane, to be tested. To date, every PN subjected to this investigation has displayed the predicted alignment, indicating that binarity has played an important role in the formation and evolution of these nebulae. The further results from this programme will be key, not only in determining whether binary interaction is responsible for shaping the studied PNe, but also in assessing the importance of binarity in the formation and evolution of all PNe in general.

Keywords. binaries: close, ISM: kinematics and dynamics, ISM: jets and outflows, planetary nebulae: general

Central star binarity has long been believed to be associated with the formation of aspherical PNe (de Marco 2009), with an important prediction of all shaping theories being the perpendicular alignment between nebular symmetry axis and binary orbital plane (Mitchell 2007). Therefore, the comparison of nebular and binary orientation provides an obvious test of this ‘binary hypothesis’. Detailed spatio-kinematic modelling is required to accurately determine the morphology and inclination of a given PN due to projection effects in imaging studies. So far, only six PNe with well-studied binary central stars (i.e. with a known binary inclination) have been the subject of detailed spatio-kinematic study – but all have been found to show the expected alignment. The details of the six PNe are listed in Table 1.

A collective analysis of the morphologies and kinematics of the six PNe reveals far more differences than similarities (Jones *et al.* 2011), indicating that understanding the exact shaping mechanism(s) involved will be far from trivial. However, some intriguing avenues for study are thrown up by the sample, all of which are discussed in more detail in Jones *et al.* (2011) and will be further explored in upcoming papers (Jones *et al.*, Santander-García *et al.*, in prep.). Possibly the most interesting discovery is the

Table 1. A table of PNe, with well studied binary central stars, that have been the subject of detailed spatio-kinematic study.

PN	Morphology	References
Abell 41	Open-ended bipolar with equatorial torus	Jones <i>et al.</i> 2010
Abell 63	Barrel-like with bipolar “end-caps”	Mitchell <i>et al.</i> 2007
Abell 65	Double-shelled bipolar	Huckvale <i>et al.</i> 2011
HaTr 4	Elliptical with equatorial torus	Tyndall <i>et al.</i> 2011a,b
NGC 6337	Torus without bipolar lobes, Jet-like corkscrew outflow	García-Díaz <i>et al.</i> 2009
Sp 1	End-on bipolar	Hillwig <i>et al.</i> 2010 Jones <i>et al.</i> 2011

Table 2. The kinematical ages of PNe with binary central stars and their polar outflows.

PN	PN age (yr kpc ⁻¹)	Polar outflow age (yr kpc ⁻¹)	Reference
Abell 63	3500±200	5200±1200	Mitchell <i>et al.</i> 2007
ETHOS 1	900±100	1750±250	Miszalski <i>et al.</i> 2011
The Necklace	1100±100	2350±450	Corradi <i>et al.</i> 2011

presence of extended outflows in three PNe which are older than the central nebula, most likely being formed before the central binary entered the common-envelope phase (see Table 2).

A programme is now underway to derive spatio-kinematic models from all PNe known to contain a post-common-envelope central star, not only to compare the nebular inclinations to those of the central stars but also to further investigate the links between the morpho-kinematic properties of the nebula and the parameters of the central star systems. The results of this investigation will prove invaluable in understanding the processes at work in formation and evolution of these systems.

References

Corradi, R. L. M., Sabin, L., Miszalski, B., Rodríguez-Gil, P., Santander-García, M., Jones, D., Drew, J. E., Mampaso, A., Barlow, M. J., Rubio-Díez, M. M., Casares, J., Viironen, K., Frew, D. J., Giammanco, C., Greimel, R., & Sale, S. E. 2011, *MNRAS*, 410, 1349

de Marco, O. 2009 *PASP*, 121, 316

García-Díaz, M. T., Clark, D. M., López, J. A., Steffen, W., & Richer, M. 2009, *ApJ*, 699, 1633

Hillwig, T. C., Bond, H. E., Afşar, M., & de Marco, O. 2010, *AJ*, 140, 319

Huckvale, L., Prouse, B., Jones, D., Lloyd, M., O’Brien, T. J., & Pollacco, D. 2011, in: A. A. Zijlstra, F. Lykou, I. McDonald & E. Lagadec (eds.), *Spatio-kinematic modelling of the double-shelled PN Abell 65*, Proc. APN5 (ebrary), p. 119

Jones, D., Tyndall, A. A., Huckvale, L., Prouse, B., & Lloyd, M. 2011, in: L. Schmidtbreich, M. Schreiber & C. Tappert (eds.), *Spatio-kinematic modelling: Testing the link between planetary nebulae and close binaries*, ASP Conference Series, p. 179

Jones, D., Lloyd, M., Santander-García, M., López, J. A., Meaburn, J., Mitchell, D. L., O’Brien, T. J., Pollacco, D., & Rubio-Díez, M. M. 2010, *MNRAS*, 408, 2312

Miszalski, B., Corradi, R. L. M., Boffin, H. M. J., Jones, D., Sabin, L., Santander-García, M., Rodríguez-Gil, P., & Rubio-Díez, M. M. 2011, *MNRAS*, 413, 1264

Mitchell, D. L., Pollacco, D., O’Brien, T. J., Bryce, M., López, J. A., Meaburn, J., & Vaytet, N. M. H. 2007, *MNRAS*, 374, 1404

Tyndall, A. A., Jones, D., Lloyd, M., O’Brien, T. J., Pollacco, D. L. & Mitchell, D. L. 2011a in: A. A. Zijlstra, F. Lykou, I. McDonald & E. Lagadec (eds.), *HaTr 4 - A Kinematical Study*, Proc. APN5 (ebrary), p. 121

Tyndall, A. A., Jones, D., Lloyd, M., O’Brien, T. J., & Pollacco, D. L. 2011b *MNRAS*, submitted