

## Original Article

**Cite this article:** Verhagen JGM, Kluiving SJ, and Kars H. The option of Roman canal construction by Drusus in the Vecht river area (the Netherlands): a geoarchaeological approach. *Netherlands Journal of Geosciences*, Volume 101, e4. <https://doi.org/10.1017/njg.2022.2>

Received: 3 June 2021

Revised: 18 December 2021

Accepted: 6 January 2022

### Keywords:

Geoarchaeology; Roman period; Canals; Vecht river; Rhine delta



### Author for correspondence:

Jan G. M. Verhagen,

Email: [jangmverhagen@planet.nl](mailto:jangmverhagen@planet.nl)

<sup>†</sup>Deceased

# The option of Roman canal construction by Drusus in the Vecht river area (the Netherlands): a geoarchaeological approach

Jan G. M. Verhagen<sup>1,2,3</sup> , Sjoerd J. Kluiving<sup>2,3</sup>  and Henk Kars<sup>1,3,†</sup>

<sup>1</sup>Faculty of Science, Cluster Earth and Climate, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands; <sup>2</sup>Faculty of Humanities, Department of Archaeology, VU University Amsterdam, De Boelelaan 1105, 1081 HV Amsterdam, The Netherlands and <sup>3</sup>Research Institute for Culture, Cognition, History and Heritage (CLUE+), VU University Amsterdam, De Boelelaan 1105, 1081 HV Amsterdam, The Netherlands

## Abstract

This paper presents a geoarchaeological study on potential canal subsections present in the Roman-age Vecht branch of the Rhine-Meuse delta (the Netherlands). The first Roman canals in this delta were dug around 12 BC by Drusus, but their location has been the subject of debate since the 16th century, with various hypotheses proposed. Based on actual palaeogeographical knowledge of the Rhine-Meuse delta, the Utrechtse Vecht hypothesis is considered the most plausible. Within the study area, in the northern part of the Vecht system, natural sections of this river may alternate with possible artificial reaches, created at the time of Drusus.

Such artificial canals, being part of an otherwise natural channel belt system, can widen and deepen overtime, eroding all or most of the recognizable features associated with their original construction. As study area was chosen a relatively straight section of the Vecht between two former lakes. Two approaches were used. The first approach centred upon mapping channel morphology and recording sediment stratigraphy of the river deposits through detailed auger coring. Results corroborated the hypothesis of an originally straight feature (landform), confirming that it might have started life as a dug course, but not providing preserved archaeological remains of this stage. The second approach was chronological, whereby a programme of <sup>14</sup>C dating was undertaken to refine the understanding of the origin and development of this reach of the Vecht, allowing earlier chronological investigations to be further contextualised and reassessed. A significant challenge to understand age control and floodplain evolution is the degradation of the top of the clayey peat that was observed below the levee deposits; this degradation is due to the lowering of groundwater levels and causes the end of peat growth to be dated as older than it actually is.

Using new radiocarbon dates we have reconstructed that the Overmeer-Nigtevecht reach of the Vecht between two former lakes started life as a straight channel. We have constrained its age to be closer to the time of Drusus' activities (early Roman age). Although we have not found *in situ* remains of Drusus canal(s), these two new insights make the Vecht option, effectuated by a series of short canals, more likely to be the Drusus canal(s).

## 1. Introduction

The Rhine–Meuse delta has multiple branches, the main ones extending to the North Sea and secondary ones draining into a central lagoon. Most of these channels have a natural origin, but during the last millennia humans increasingly used and adapted the river networks. The time of Roman occupation, beginning c. 12 BC under army commander Drusus, is considered the period with the oldest humanly organised waterworks in the delta. The Romans created a shipping network through the construction of canals (Willems, 1981/1984, 55 ff., 387 ff.; De Kort & Raczynski-Henk, 2014) and by influencing the hydrology of the natural delta channels by groyne construction (Vollgraff, 1938, 1939; Schönfeld, 1940a, 1940b; Willems, 1981/1984, 52 ff); further adaptations included the construction of harbours and other facilities to load and unload cargo (Graafstal, 2002; Blackman, 2008; Morhange et al., 2017; Mirschenz, 2018). The construction of such efficient transport networks and the associated trade opportunities contributed to the successful consolidation of the vast Roman Empire (Ruffing, 2018, 8). Whilst transportation overland was still important, testified by a network of well-maintained roads (Chevallier & Field, 1976; Rathmann, 2004; Klee, 2010, 65 ff., 107ff.; Van Lanen, 2017; Van Lanen & Pierik, 2017), much larger cargoes, both goods and people, could be transported by water (Bechert & Willems, 1995, 24; Eck, 2007, 111; Grewe, 2008, 333; Jansma et al., 2017). Such advantages led the Romans to construct shipping networks made up of canals, natural river channels and sea lanes across the Empire; e.g. several canals were built in and around the

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Netherlands Journal of Geosciences Foundation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Po delta (northern Italy), creating a network that covered a large area immediately behind the coast (Medas, 2013, 2017).

The first Roman canal(s) in the Rhine–Meuse delta, engineered under the leadership of Drusus between 12 and 9 BC, is/are mentioned by two classical authors: Tacitus (in works produced c. 110 AD) and Suetonius (c. 120 AD). Tacitus describes how in 16 AD Germanicus sailed into the Drusus Canal (*fossam, cui Drusianae nomen*) with his ships containing legions and allies (Tacitus, *Annales* II, 8). Following a prosperous journey over lakes and the ocean, he reached the river Ems (Fig. 2). Suetonius reports that Drusus was the first of the Roman commanders to sail the ocean (North Sea) and had canals dug at the other side of the Rhine. These canals were an unprecedented and significant feat of engineering (*fossas novi et immensi operis*); more than 50 years after Drusus, they were still called Drusus Canals (*Drusinae*) (Suetonius, *Vita divi Claudii* I, 2–4). It is striking that the plural ‘canals’ is used here. The classical references indicate that the canal(s) played a role in the Roman attempts to subjugate North Germania and created a navigable connection between the Rhine delta and the Wadden Sea, from where the ships with troops could sail up the North German rivers (the Ems, Weser and Elbe). The use of the new connection avoided the risks of sailing on inhospitable high seas and the logistical problems of transshipment of river- to sea-going vessels and vice versa.

The location of the Drusus Canal(s) has been an issue that has vexed researchers for more than four centuries. For a long time, the route of the river IJssel in the east of the Rhine delta (Fig. 1) was the prime candidate to project Drusus’ canal works along. In the second half of the 16th century, historians and cartographers were convinced that the upper reach of the IJssel (Boven-IJssel in Fig. 1) was the site and the end result of Drusus’ canal digging mentioned in the classic texts. Later, many more hypotheses about the location of the canal(s) of Drusus have been put forward, which are now the subject of inventory and further evaluation (and the broader topic of the PhD thesis project of the first author). An important point here is that for a considerable time researchers thought that towards the end of prehistory the Oude IJssel (Old IJssel) from Doesburg continued northward (Van de Meene, 1979; Teunissen, 1980 and underlying references). New insights into the natural development of the lower reaches of the IJssel (Beneden-IJssel in Fig. 1) have revealed that this Rhine branch is evidently younger than hitherto considered (Makaske *et al.*, 2008, 333) and that it originated in the early Middle Ages (or possibly in the late Roman period) by a natural breakthrough between Doesburg and Deventer (Cohen *et al.*, 2009, 103). The implication of this is that when Drusus arrived in this region, the (Oude) IJssel from Doesburg ran south of the Veluwe to the west merging with the Rhine system at Arnhem (Fig. 2, compare Fig. 1)

Despite the multiple hypotheses put forward, the presence of not a single metre of a Canal of Drusus has yet been corroborated. This stands in sharp contrast to what is known about another Roman canal mentioned by classic sources, constructed under Governor Corbulo in the very west of the delta. Corbulo’s Canal (c. 50 AD) was constructed parallel to the beach barrier complex of South-Holland; since 1989, archaeological evidence for the feature has been identified in dozens of places along a stretch of about 13 km (De Kort & Rackzynski-Henk, 2014; Polak *et al.*, 2019; Hessing & Schrijvers, 2021). This difference might be explained by the fact that in South-Holland much archaeological research was carried out, in response to intensive building activities. A more fundamental explanation may be that the positions of these two waterways within the landscape were basically different.

Whereas Corbulo’s canal was constructed between two roughly equivalent points at the Roman Rhine and the Helinium (Fig. 2), most hypotheses for the Drusus canals assume that they became part of a river network, similar to the Pannerdensch Kanaal in 1707 (Fig. 1; Van de Ven, 2007) and the 16th-century hypothesis of the Boven-IJssel as the Drusus Canal.

In the case of the Drusus Canal(s) post-depositional processes (mainly erosion by widening, deepening and lateral migration) may have resulted in the remains becoming not or hardly recognisable in the subsurface, whereas in the case of the Corbulo Canal, such processes (mainly sedimentation) will not have had a major impact. In ideal cases, when remains of an original canal construction have not been eroded, evidence for revetments and digging (created cut surfaces) are usually found. When little or no *in situ* remains are likely to have been preserved, research will have to focus on indirect indications: e.g. the morphology and depositional chronology of a river section. Furthermore, in the case of the Drusus Canal(s), until this study, no targeted fieldwork had been undertaken, simply because it has been like looking for a needle in a haystack. However, progress in physical geographical and geoarchaeological research in the Rhine–Meuse delta (e.g. Makaske *et al.*, 2008; Bos *et al.*, 2009; Cohen *et al.*, 2009; Erkens, 2009; Van Dinter, 2013) and the coastal area (Vos, 2015) has been used to reassess and partly eliminate the hypotheses about the Drusus Canal(s), from which the Vecht emerged as the most probable Drusus route.

In this study, we first describe the implications of the new palaeogeographical and archaeological insights for our knowledge of the location and function of the Drusus Canal(s), resulting in a new working hypothesis (Section 2). Second, we present the methods and results of our fieldwork (Sections 3 and 4). Finally, we assess whether there are concrete indications of active human intervention by the Romans in the area of the Utrechtse Vecht (Sections 5 and 6).

## 2. Investigating the Drusus Canals using a geoarchaeological perspective

### 2.1. The focus on the Vecht river

Notwithstanding the many hypotheses described in earlier literature, in the actual state of research on the location of the Drusus Canal(s), only three variants remain (denoted as options R1 to R3 below). Each of these would fit well with the new palaeogeographical insights and the location ‘at the opposite side of the Rhine’ mentioned by Suetonius. Figure 2 plots the three remaining possible canal routes of Drusus, they are:

**Option R1**, a route making use of the Utrechtse Vecht. This hypothesis was raised by Van Asch van Wijk (1846) and strengthened by the discovery of the *castellum* (auxiliary fort) near Vechten at the end of the 19th century. This branch of the Rhine is part of the Angstel-Vecht system, a deltaic branch of the Rhine, which evolved by an avulsion of the Rhine near Vechten and Utrecht, between 1000 and 800 BC (Törnqvist, 1993, 149; Bos *et al.*, 2009, 368; Van Dinter *et al.*, 2017).

**Option R2**, a route through the Gelderse Vallei (Fig. 2), a lowland area between the ice-pushed ridges of the Veluwe and the Utrechtse Heuvelrug. The southern part of the valley was originally a peat bog, the central part consisting mainly of cover sands, the northern part peat land again. The area was dissected by some local streams, congregating into the northward draining Eem river. A smaller counterpart stream drained southward towards the Rhine.



**Fig. 1.** The Rhine–Meuse delta in the Netherlands, with annotation of the places and rivers mentioned in the text. The current active river branches are shown in dark blue. The branches that only drain water locally are shown in black. Map background: Vos & de Vries (2013) (v 2.0), situation 2000 AD.

**Option R3**, the valley of the (current) Beneden-IJssel, that lies between the Veluwe ice-pushed ridge at the west side and an area with cover sands and a locally ice-pushed ridge at the east side. In prehistoric times up to early middle ages, streams from these areas carried most of their water to the north (Dortherbeek/Hunnepe, Duurse Beek and Fliert near Deventer; Spek et al., 1996), but some drained south (Berkel and Voorsterbeek near Zutphen) to the Oude IJssel, which in turn, flowed into the Rhine near Arnhem (Fig. 2). Cohen et al. (2009, 104) suggest that it is possible that Drusus has dug a canal through the drainage divide, but to date no traces of such a connection have been found. Traces of such a canal might be eroded by the strongly meandering Beneden-IJssel.

Each of these three options has arguments pro and contra, which are the subject of evaluation (PhD thesis project of the first author). Because of the scope of this study, we will limit ourselves here to a few arguments. Option R1 would have been a (canal) improvement of a natural route, while R2 and R3 would have involved digging across a drainage divide to make a connection between two brook systems, implicating problems with water levels. A practical advantage of option R1 compared to option R3 could have been that it was possible to create a freely flowing waterway, assuming a relationship between the Dam (*moles*) and the Canal(s) of Drusus, both serving the aim to provide a waterborne transport route (Verhagen et al., 2017, 461–463; Fig. 2). From a (military) functional point of view, this is favoured over a combination of the Drusus Dam and an IJssel connection (R3). In the

latter case, the extra water should have back-flooded areas between Westervoort and Doesburg to reach the possible artificial connection across the drainage divide (somewhere between Doesburg and Deventer), connecting the separate brook systems in the IJssel valley. This could have been achieved ‘whether by digging an artificial connection or by creating some kind of portage’, while ‘a difference in water level could be overcome by the construction of a lock, possibly in the form of temporary dams or weirs’ (Polak & Kooistra, 2013, 403–404). This option is still a possible one, but we regard it as less probable. Also the Gelderse Vallei connection (R2) would have needed a portage or a lock. In this area, the palaeogeography of the subsoil is still completely intact, while there is no indication that there has been a through-going watercourse (natural or artificial) from the Rhine delta branch. By this, option R2 has got hardly any support among researchers.

Focusing on the Vecht (R1) option as the most plausible candidate, there are some aspects to be considered in order to research and understand the role of the Vecht connection within the river network better:

- (1) There is the fairly widespread view that the Vecht was navigated by the Romans (see among others Willems, 1981/1984, 58; Kok, 2007, 54; Manten, 2007; Vos et al., 2015, 320). In other studies, the supposed routes of the fleet campaigns of Drusus, Tiberius and Germanicus are drawn on maps via the river Vecht (Lendering & Bosman, 2010, 69,






**Fig. 2.** Remaining options for the navigation route to the north, created under the direction of Drusus. Possible channels are shown in red. Local improvements may have been made in the Rhine section (purple). River network is from early Roman period. Inset: contemporary NW European river network with routes of the fleet trips of Drusus, Tiberius and Germanicus between 12 BC and 16 AD. Palaeogeographical map background: Vos & de Vries (2013) (v 2.0), situation iron age, c. 500 BC.

- 81, 105; Kehne, 2018, 46–49). Willems (1981/1984, 58) and Manten (2007, 63) even take the position that the Drusus' Canal(s) had been constructed in the area of the IJssel, but that the Romans had also used the Vecht as a navigation route. A fundamental question is whether the Vecht could be navigated by the Romans *because* of canal construction in its reaches or *parallel* to canal construction somewhere else. Anyhow, from an operational military point of view, the early *castella* at Vechten and Velsen presuppose a navigable route along the Vecht.
- (2) Some studies in recent decades have discussed the possibility of human intervention (in Roman and later times) by altering the course of parts of the Vecht, without reaching solid conclusions (Van de Meene et al., 1988, 62; Weerts et al., 2002, 70; Van der Velde et al., 2003, 9; Kok, 2007, 54; Manten, 2007) (see also in the discussion section).
  - (3) Among Roman historians, there are several descriptions of a channel branch on the right side of the Rhine delta. According to Pomponius Mela, writing c. 50 AD, this branch flowed through a large lake called Flevum and then into the sea (Mela, *De situ orbis*, III, 24). According to Pliny (23–79 AD), there were three Rhine branches: the Helinium, the Rhine and a northern branch that flowed through lakes (Pliny, *Naturalis*

*Historia* IV, 29). Ptolemy (87–168 AD) mentions three Rhine mouths north of Lugdunum Batavorum (Katwijk aan Zee) (Ptolemy, *Geographia* II, 9, 1); although there are also discrepancies in his geographical coordinates, from the mentioned order along the coast it seems likely that these must be the Oude Rijn, the Oer-IJ and the Vlie. According to Tacitus, Germanicus and his fleet entered the canal named after Drusus and sailed through lakes and the ocean to the Ems (Tacitus, *Annales* II, 8). The mention of lakes and the large lake (*ingens lacus*) Flevum corresponds more to the situation in the Vecht area with its peat-bounded lakes and the Oer-IJ than to the situation of options R2 and R3.

It is not unreasonable to suggest that the Romans were forced to create an artificial route for their flat-bottomed boats in the Angstel-Vecht area, since the river water ran through peat-bounded lakes (see Section 2.2), which in their proximal parts were filled by clastic sediments with a network of small distributaries, while the fluvial activity of the Angstel-Vecht system was strongly reduced during the last few centuries BC (Bos et al., 2009, 368). It is possible that with some human intervention this branch of the river could have been reopened. Hence, we formulate the hypothesis that the Vecht is the most likely location of the 'Drusus





**Fig. 3.** Geological cross-section through the Angstel and Vecht system and adjacent ice-pushed ridge of the Utrechtse Heuvelrug. Transect line is shown in Fig. 4. Legend: 1 = glaciotectionised Early and Middle Pleistocene; 2 = glacio-fluvial outwash sands; 3 = Late Pleistocene cover sands; 4 = Holocene peat (Nieuwkoop Formation); 5 = backbarrier tidal clays (Naaldwijk Formation); 6 = deltaic river channel deposits; 7 = natural levee deposits; 8 = flood basin deposits; 9 = water; 10 = anthropogenic deposits (dike/raising). After: Bos et al. (2009), Figs. 3, 6C and 6D.

Canal(s)'. In this study, a geoarchaeological programme of new field research and associated analyses was undertaken in this area of the river Vecht to test our hypothesis.

## 2.2. Geomorphology of the Angstel and Vecht system

The system of the Angstel and Vecht is located west of the Utrechtse Heuvelrug, the latter area pushed up during the Saalian glaciation. The associated fluvioglacial sandur deposits (Formation of Drente) form the deeper substrate (several m) of the study area, while their top decreases towards the west (Fig. 3). The channel belt cuts through both the Holocene peat and the underlying Pleistocene sand (Weerts et al., 2002; Bos et al., 2009; Bos, 2010). The natural levee and flood basin deposits of the Angstel and Vecht overlie the peat sequence. Bos et al. (2009, 359) have provided a detailed insight into the various river channels, including their migration and crevasse splay development within this system (Fig. 4).

The Vecht and its older sister branch, the Angstel, are part of the Rhine–Meuse delta, which in the Holocene has an extensive history of channel movements, with meander formation, cut offs and channel avulsions (Berendsen & Stouthamer, 2001). The Angstel–Vecht system is positioned at the edge of the active delta. The avulsion which created the Angstel (first phase of the Angstel and Vecht system; Fig. 4, nr. 1), occurred along the prehistoric Rhine near Vechten/Utrecht (Fig. 1), and is dated to c. 900–700 BC (Table 1, E1 and E2; Törnqvist, 1993, 149) and c. 1100–900 BC (Table 1, E4, E7 and E9; Bos et al., 2009, 365). In the second phase, the water discharge had largely shifted to a new course, the Vecht. The onset of this phase has been dated 413–197 BC and 740–233 BC (Table 1, E6 and E10; Bos et al., 2009, 368). These dates are TPQ (*terminus post quem*) dates, which leaves room for an onset up to some time later.


At the bifurcation point of the Rhine and Angstel, the Angstel phase channels have been partly reworked by younger ones during the Roman period and Middle Ages or may be hidden beneath them (Van Dinter et al., 2017, Fig. 6). The Angstel channel created a connection with the northern coastal area, consisting of beach barriers, estuaries with tidal creeks and a coastal plain with a perimarine zone comprising an extensive peat vegetation. Before the avulsion, rainwater and groundwater flow from the ice-pushed

ridge drained via local streams (like the Amstel, Holendrecht and Gein/Gaasp) into the Oer-IJ estuary and the central Netherlands area of lagoonal lakes. The new river channel annexed one of these local peat rivers, creating a connection with the Oer-IJ and the lagoonal lakes (Fig. 2). This area, the development of which has become better known by various studies, also received water from the Rhine from then onwards (Vos et al., 2015, 310; Kluiving & Borger, 2015, 290; De Bont, 2015, 354; De Gans, 2015, 362; Kranendonk et al., 2015, 348ff).

Palaeogeographical reconstruction of the Oer-IJ estuary has shown that it evolved together with the Angstel and Vecht channel belt system (Vos, 2008). Its emergence as a branch of the Rhine delta may have stimulated the development of the Oer-IJ by increased discharges, which may have enlarged the tidal inlet and main stream of the Oer-IJ (Vos et al., 2015, 319). After several centuries as a fully functioning channel belt, the discharge of the system decreased. Around 200 BC, this also may have contributed to the gradual closure of the mouth of the Oer-IJ (Bos et al., 2009, 371), by which the natural drainage of the Oer-IJ was forced in the opposite direction and stalled in the Flevo Lake. The ponding water in the Flevo lake then began to drain northwards to the Wadden Sea; in turn, this may have initiated the opening of a connection between the northern part of the Flevo Lake and the Wadden Sea (Vos et al., 2015, 320; Van Popta et al., 2020, 38).

Nevertheless, it is plausible that the Romans still managed to keep open a short navigable connection between the Oer-IJ and the sea (Borger & Kluiving, 2017, 48). The construction of *castellum* Velsen 1, which had a large harbour with piers (Bosman, 1999, 2016, 30), and then Velsen 2, built in the most important period of actions against the Chaucians operating at sea (Bosman et al., 1998; Bosman, 2016, 69 ff.), make it plausible that there was such a connection between the *castellum* and the sea; however, it is theoretically possible that ships navigated from the *castellum* to the southern North Sea via the Vlie (Vos, 2015, 327).

The Flevo Lake was positioned around the location of the current Markermeer and was called Flevum by the Romans (Pliny, *Naturalis Historia* IV, 29; Mela, *De Chorographia* III, 24). Around 755 AD it is mentioned as 'Almere' and 'stagnum', meaning that tidal currents had (still) little or no influence there at that time (Levison, 1905; Rau, 1968).



**Fig. 4.** Overview of the Angstel and Vecht system, developed as an avulsion of the Rhine, as well as the peat rivers. Legend: 1 = phase 1, discharge via Oud-Aa and Angstel; 2 = phase 2, discharge entirely via the present Vecht. The rectangle near Nigtevecht and Overmeer shows the study area (field research). The orange line shows the position of the cross-section of Fig. 3. Map inset: possibly artificial connections that have developed into river channels. Red ovals = connections between former peat-bounded lakes (sandy and clayey lake fills at B = Breukelense Meer, L = Loenermeer, H = Horstermeer, A = Aetsveldse Meer, M = Muidermeer). Yellow ovals = possibly constructed fairway (lane) or canal within the in late prehistory still water-bearing part of the Aetsveldse Meer. The two avulsion points conform Bos et al. (2009) are indicated in blue. Map background Angstel-Vecht: Bos et al. (2009, 359). Map of other areas: conforms to geomorphological map of the Netherlands and Van Dinter (2017).




A notable phenomenon in the Angstel–Vecht–OerIJ–Flevum river area (Fig. 4) is formed by smaller peat-bounded lakes. Their establishment, dated to between 4700 and 3500 cal. yr. BP, has been related to seepage from the ice-pushed ridges and subsequent erosion of the peat by wind-generated waves (Bos et al., 2009, 365 and 370). When the separating peatland was breached by flood waters, river channel and lake sections got connected and sedimentation was initiated, leading to gradual infilling of the lake with clastic materials (clays, sands) and eventual establishment of a traversing channel section (Bos, 2010). The discharge of river water from the Rhine delta via this system no longer played a significant role after the damming of the Kromme Rijn near Wijk bij Duurstede (Fig. 1) in 1122 AD (Dekker, 1980), which has been confirmed by  $^{14}\text{C}$  dating (Van Dinter et al., 2017, 31).

The channels of the Angstel and Vecht stages of system development are two different ones over most of the study area. North of Breukelen they are rarely amalgamated, thereby allowing two phases in the system to be easily distinguished (Weerts et al., 2002, 67). South of Breukelen the two phases can be distinguished at a point only about 3 km northwest of the centre of Utrecht (Fig. 4; Van Dinter et al., 2017, appendix, 53). In the first phase (up from 1000/800 BC), the water flowed from Breukelen via the Oud-Aa (in the Breukelense Meer; ‘meer’ = lake) to the Loenermeer and then via the Angstel to the Aetsveldse Meer (1 in Fig. 4). As a result of the supply of sediment from the Rhine, the Breukelense Meer and the Loenermeer soon silted up (Bos et al., 2009, 366). Sedimentation in the Aetsveldse Meer developed from the southwest (mouth of the Angstel) to the northeast (outlet at Weesp), forming a branched system of (constantly rejuvenating) crevasse splays in the deposits of the lake (Weerts et al., 2002, 69; Bos et al., 2009, 368; Bos, 2010, 5).

In the second phase, the main body of the water flowed via new tracts of channel near Breukelen and from Loenen through the Horstermeer and Aetsveldse Meer to Muiden, thus creating the present-day Vecht (2 in Fig. 4). Compared to the Angstel, this course is an alternative route between Utrecht and the Flevum/Oer-IJ receiving waterbodies. It has been established by a series of shortcuts between phase-1 channels and peat-bounded lakes (Lake Horstermeer, Aetsveldse Meer; Fig. 4). Bos et al. (2009, 368) mapped and dated two shortcuts and describe them as local avulsions that were naturally triggered. One avulsion originated at the east side of Breukelen, creating a bypass of the Oud-Aa around 413–197 BC (Table 1/Fig. 4, E6). The other avulsion originated near Loenen around 740–233 BC (Table 1/Fig. 4, E10). Note that these are TPQ dates. The two local avulsions need not have occurred simultaneously (Bos et al., 2009, 368). A TAQ dating result is known from the silting-up Angstel near Breukelen (12–327 AD; Table 1/Fig. 4, E8).

### 2.3. Choice of study location and approach

Using the geomorphology (Section 2.2) as a basis, we refocus on the Drusus Canal hypothesis option R1. In the light of the mention of the Canal(s) of Drusus in the classical literature, the aim of our study is to identify potential canal subsections present in the Vecht branch. Among the shortcuts in the area of the peat-bounded lakes (see Fig. 4, inset), the 2-km long one between Overmeer and Nigtevecht (H-A) has the narrowest channel belt width, no recognisable meander cut-offs and the most linear planform. On the onset of this Vecht river reach, we distinguish two possibilities: (a) this section began its life as a canal dug in the time of Drusus (12–9 BC); (b) this section had a natural origin, probably



**Fig. 5.** Three map images of local rivers in the Angstel–Vecht area (for comparison). V1 = section of the Vecht between Overmeer and Nigtevecht; A1 = section of the Angstel between Loenerslout and Abcoude; A2 = section of (a residual channel of) the Angstel between Loenen and Loenerslout in the former Loenermeer. For cross-sections of V1 and A1 see Fig. 3. Legend: 1 = river belt; 2 = natural levee deposits and crevasses (mostly on peat); 3 = flood basin deposits on peat; 4 = peat; 5 = sandy lake fills; 6 = clayey lake fills; 7 = coversands. Source: Bos et al. (2009, 359).

at an earlier moment in the last few centuries BC, conform the TPQ dates of Bos et al. (2009, 368). However, the amount of dating available at present in this area is still limited and in need of further research. Because the Vecht onset mainly can be determined by dating the top of the peat immediately below the natural levee deposits, also the possibility of degradation of the top of the peat has to be taken into account, which was a relevant factor in some earlier studies in similar areas (Makaske et al., 2008; Stouthamer et al., 2008; Pierik et al., 2018, Fig. 4C). Usually, the degradation is the result of the oxidation of organic materials, but sometimes anaerobic bacteria can also cause decomposition (Hoogland et al., 2020, 750). The degradation can have various causes, such as events immediately preceding the covering by the clay, (Roman) canal digging and resulting drainage, medieval peatland reclamation with the digging of ditches and lowering of groundwater levels prior to modern agriculture.

The possibility of human intervention along the course of the river in the Angstel–Vecht system has already been put forward without reaching conclusions (Section 2.1). A possible method of canal construction is that the Romans created a more easterly route by connecting several peat-bounded lakes, partially filled with clastic materials (Fig. 4, inset, connections depicted as red ovals). Locally, it is possible that crevasse splays were already active, but which had not developed into complete avulsions and which were used by the Romans in their canal construction. This leads us to hypothesise that the relatively younger short-cut channels between a number of peat lakes in the Angstel–Vecht system downstream of Breukelen are the result of human intervention by the Romans who dug one or more canals, which subsequently evolved into a river channel.

The Vecht section between Overmeer (exit Horstermeer) and Nigtevecht (entry Aetsveldse Meer) shows a striking planform (Fig. 4, H-A), which despite the meander loops, creates the impression of a relatively straight landform (Fig. 5, V1). The channel belt profile of this 2 km long river section is particularly narrow (Fig. 3), with a short lateral migration path of the meander loops (Bos et al., 2009). It has a channel belt width of approximately 150 m, in contrast to the profiles elsewhere in the Angstel–Vecht system (Fig. 5, A1 and A2) which show widths ranging from 250 to 1100 m.

Based on these considerations, it was decided to carry out field research on the reach of the Vecht between Overmeer and Nigtevecht (= study area). To test our working hypothesis, we



used two approaches. The first approach focused on the geomorphology and sedimentology and the need to understand the detailed morphology and sediment architecture of the selected reach of the Vecht. The second approach was focused on obtaining a more detailed chronology. There are three points of entry to investigate this. The first is that from the moment of (possible) canal construction, this structure starts to show river behaviour and begins to migrate laterally. The timing of the initial phase of migration should be around the moment of construction/origination or some time after that. The second point is that from the commencement of the phase 2 Vecht river, its banks which generally consist of a peat sequence, are covered by clay sediments creating natural levees and in response to this the peat growth can end. The third point is that (after some time) settlements can develop on the natural levees along the river. The oldest phase of site habitation provides a TAQ (*terminus ante quem*) date for the emergence of phase 2.

### 3. Methods

The programme of coring resulted in lithological profiles along 62 short transects (Fig. 6). Once the boundary between intact peat and the channel deposits had been sufficiently mapped out, the most suitable locations were determined for coring along two longer and four medium length transects, in order to investigate the chronology of meander migration. The main transects (AC and HB) were positioned at places where it was estimated that the erosion boundary (Fig. 6, red line) could date from relatively shortly after the assumed canal construction. Coring to date the top of the peat was carried out in the parts of the long transects which lay within the zone of the preserved peat. In total, the combined coring studies led to the drilling of 439 hand-auger cores between November 2015 and October 2018.

Coring was performed using a 7-cm Edelman hand auger (screw-type) and a 3-cm gouge auger. The cores generally reached depths of between 4 and 5 m below ground level, while the distance between individual cores along the transects varied between 5 and 25 m. Along longer transects, a mutual distance between successive cores of 10 m was maintained across the channel fill, whilst in the surrounding peat areas a minimum distance of 50 m was used. The spatial coordinates of the drill holes were recorded using a handheld GPS (Garmin eTrex Vista HCx), with a spatial deviation of about 3 m. Core descriptions were recorded in the Deborah program (release 3), developed by RAAP Archaeological Consultancy, the Netherlands ([www.raap.nl](http://www.raap.nl)).

In order to gain a better insight into the stratigraphy of sediments and possible disturbance by anthropogenic activities such as ploughing, profile descriptions were recorded in small test pits, each 1 m deep at three localities (Fig. 6). Profile descriptions were harmonised with the core descriptions recorded in Deborah. Samples for dating were taken here by hammering in steel boxes.

Field description of the core lithologies is based on the NEN 5104 method (Nederlands Normalisatie-instituut, 1989), which distinguishes: depth, colour, texture, CaCO<sub>3</sub> content, sedimentary structures, the presence of inclusions (organic matter, shells, gravel, anthropogenic materials) and oxidation/reduction phenomena. Grain size of sand was determined with a field reference. Calcium content was determined on all samples using a 10% HCl solution.

In total, 415 sediment samples with the potential to contain organic material suitable for <sup>14</sup>C dating were taken during

fieldwork. Presence of older organic material, which may have been redeposited, is referred to as *reworked*. In the case of a watercourse like the Vecht, migrating within an existing peat area, the possibility of recycling of organic materials must be taken into account. Only primary organic material is suitable for dating (referred to as *local*). Identifying whether organic materials were *local* or *reworked* was performed through palaeobotanical analysis by BIAx (Zaanstad, the Netherlands) prior to <sup>14</sup>C dating, with the aim of selecting as far as possible only primary remains (i.e. locally grown). The most suitable samples for <sup>14</sup>C dating were selected as described in supplementary appendix A (available online appendix in Supplementary material), while the results of the selection steps of all samples are presented in supplementary appendix B (available online appendix in Supplementary material). After analysis of the 415 samples taken, organic remains were selected from 24 samples for dating. The <sup>14</sup>C radiocarbon dating was undertaken using Accelerator Mass Spectrometry Analysis (AMS) and performed by Beta Analytic Inc. (Miami, Florida). To avoid deviations in dating results, samples were pre-treated to remove calcium carbonate and mobile humic acids. All radiocarbon ages reported in this paper, both our own results and the results of others described earlier, were recalibrated in 2021 using the OxCal v4.4.4 calibration curve (Bronk Ramsey, 2021). Recalibrated ages are presented at 1 and 2 sigma probability in Table 1. Ages mentioned in text are all presented at 2 sigma probability.

In order to get an insight into the presence and ages of settlements on the natural levees of the Angstel–Vecht system and possible earlier human activities in the peatland, an inventory of archaeological finds and sites has been compiled for the three periods of prehistory and the Roman Age. Since the start of the Roman period is the crucial time boundary for our working hypothesis, an inventory of medieval and younger finds was not considered useful for this research. The method and table of results of this inventory can be found in supplementary appendix C (available online appendix in Supplementary material).


## 4. Results

### 4.1. Substrate


This section describes the detailed mapping of the researched area using core transects depicted in Fig. 6. Results show core positions through the intact peat, channel belt and infilled peat lakes. Transects AC and HB show the stratigraphy and the results of <sup>14</sup>C dating (Figs. 7 and 8). Four lithological units are recognised:

\***Cover sand:** Moderately silty, moderately fine to extremely fine non-calcareous sand. At the transition with the overlying peat, the sand is moderately humic.


\***Coastal plain margin peat:** Sequence of Holocene wood peat, 2–5 m below sea level. It has a low mineral content at the bottom and is clayey in its upper part. At the top, directly under the clay of the natural levee deposits, a sub-unit Hpd (= Holocene peat degraded) is distinguished (Fig. 7). This unit has a varying composition of layers of clayey peat, peaty clay and humic clay, which can be seen as more or less decomposed peat. The profiles of the test pits revealed a clear unit structure and undisturbed stratigraphy (Figs. 6, 9A/B). The implication of this is that unit Hpd is not the result of mechanical disturbance by anthropogenic activities up from medieval land reclamation, such as deforestation, ploughing and digging of ditches. The latter are easily recognisable when being present. The recognition of unit Hpd is also based on the dating results and therefore it is described further in Section 4.2.1.



**Fig. 6.** Geomorphology of the reach of the river Vecht between the former Horstermeer and Aetsveldse Meer. The red line is the established boundary between the channel belt deposits and the intact Holland peat. Map inset: auger core transects from which samples have been dated using the  $^{14}\text{C}$  method. Prof A to C are the test pits which were used to record stratigraphic profiles (see section 4.2.1).



**Fig. 7.** Transect profile AC illustrating core stratigraphy and <sup>14</sup>C dated sample results (calibrated at 2 sigma). Hpd = the partially degraded top of the Holocene peat (see also discussion in section 5.1).



**Fig. 8.** Transect profile HB with core stratigraphy and <sup>14</sup>C dated sample results (calibrated at 2 sigma). On the right-hand side, the peat has been extracted to a depth of more than 2 meters below ground level, after which the pit has been filled with clay. For legend see Fig. 7.

Locally, the upper part of the peat has been removed by extraction and has been replaced by clay, which contains small fragments of brick/rubble, probably post-medieval in date (Fig. 8, NE part). This latter unit is considered artificial.

\***Natural levee deposits:** Strongly silty to moderately sandy (at the top), poorly to moderately humic clay, overlying the peat sequence.

\***Bedload and channel belt deposits:** Frequently alternating sand and clay layers. The sand layers consist of moderately to strongly silty moderately fine to very fine sand, interpreted as reworked cover sand, redeposited during periods of high fluvial activity. The clay layers of strongly silty to strongly sandy clay, generally inorganic, locally poorly humic with fine plant remains, provide evidence of low energy water conditions. Thin sub-layers are frequently present, which means that a sand layer may include many thin clay layers or vice versa. Locally, a thin peat layer is also

present. The underlying layer of primary cover sand is hard to distinguish from the redeposited sands.

The Vecht channel belt in the research area could be identified by a unit of sand/clay layers of several metre thick (Fig. 6, blue dots), hence by the absence of peat in the core (with core depths of 4–5 m below surface). Further away from the channel belt, the top of the intact peat was reached at depths of 0.4–0.8 m below the surface (Fig. 6, brown dots). At the border of these two terrain units, the (remaining) top of the peat was identified from 0.9 to 5.2 m below the surface (Fig. 6, white dots). The distinction between these border situations and the filling of the former peat-bounded lakes was based on an analysis of the elevation of the top of the remaining peat in the multiple cores ( $n = 432$ ; supplementary appendix D (available online appendix in Supplementary material)). In the former lakes, the lower part of the peat sequence has remained intact (confirming descriptions by Bos *et al.*, 2009).



Only primary meanders are present, that is, the active channel has gradually migrated to its current position and no abandoned meander remains are found. The channel belt shows a relatively straight boundary on the opposite side of each curved boundary which represents a primary meander (Fig. 6). Assuming later meander migration in the activity of the channel belt, these relatively straight edges are interpreted as the oldest sides of the meander belt and the zones to target for investigations whether the channel had a natural (avulsion through peat) or human dug (Drusus canal) origin.

The western boundary near transect AC (Fig. 6) is relatively straight, but on closer inspection not perfectly straight. It has a slightly curved shape, potentially the result of a migration to the left (west) in an initial short period. By the longitudinal component of the river migration, this meander loop has moved about 500 m to the northwest, thus causing this bank to be eroded (approximately in the middle of Fig. 6). The longitudinal migration component was stronger than the transverse component. The latter might be caused by the high erosion resistance of the peat, which can cause sharp bends and straight channels in peatlands (Candel, 2020, ch. 6.4).

#### 4.2. Radiocarbon dating results

After analysis of the 415 samples taken, organic remains were selected from 24 of them for  $^{14}\text{C}$  dating. Dating results are presented which help to determine the end of peat growth and periods of meander migration. The overview of the results is shown in Table 1, including the results of previous studies described in this paper.

##### 4.2.1. End of the peat growth by burial beneath bank sediments

Beyond the Vecht channel belt fills the Holocene peat is present, which has a degraded upper profile (Section 4.1, unit Hpd). On top of the peat in profile B, a 20-cm thick layer of relatively dark humic clay was observed (Fig. 9A). Botanical analysis of this layer yielded some unidentifiable wood fragments, but otherwise no recognisable organic remains. The top of the peat was dated to 818–760 BC (Table 1, sample N18) and to 902–803 BC (Table 1, sample N20). A younger age of 356–57 BC was obtained for sample N14 (Table 1). At these three locations (see Fig. 6), the top of the peat was covered by a dark humic clay layer with a thickness of 21, 19 and 8 cm, respectively (Fig. 9). Therefore it appears that a greater thickness of the layer of humic clay corresponds to an older age of the top of the remaining, underlying peat.

Therefore, the layer of relatively dark humic clay on top of the preserved peat (Fig. 9A) should be interpreted as a former layer of clayey peat, in which plant materials have been completely mineralised, so that only the enclosed clay remains. The same applies to the dark layer of humic clay above the peat in cores 292 and 293. The various dating results show that the degradation of (the top of) the peat has progressed further in the test pits than in core 292. This can be explained by the location of the test pits close to a drainage ditch with slightly lowered water levels, so that the degrading effect is apparently greater than in transect AC, which was located about 30 m away from the ditch. Core 292 also shows some peaty clay layers between the preserved peat layers, that will have been present prior to peat degradation.

It can be deduced that there was continuous peat growth at least until 800/900 BC (Fig. 9B, profile A, N18; Fig. 9C, core 293, N20).

The clay influx in the top of the peat as well as the dark humic clay layer covering it, are interpreted as overbank/floodbasin deposits of the river Angstel. This sediment was deposited on top of the peat, after the Angstel was formed around 900 BC. Our ages for Angstel onset confirm earlier dates in similar contexts (Törnqvist, 1993, 149; Bos et al., 2009, 366, 368), while the dating results are closest to those of Törnqvist (see Fig. 12).

In the study area, about 3 km away from the Angstel, sedimentation was limited and peat growth did not stop completely, confirmed by the three dates on peat between 800 and 100 BC obtained from core 292 (Fig. 7 and 9C). Here the dating results show that the top of the peat remained partly intact. An exception to this is the layer that was noted as clayey peat, of which botanical analysis revealed that it contained abundant wood fragments (core 235, N1; Figs. 6 and 7). This sample was  $^{14}\text{C}$  dated to 239–327 AD. Given its spatial context, this sample is interpreted as a riparian phenomenon, possibly floating woody material deposited on the bank of the river. The main picture from dating results is that a clay influx occurs up from around 900 BC (in line with earlier studies) and that peat growth continues at least until second century BC, but potentially longer because our dates are TPQ dates.

##### 4.2.2. Migration period of the meanders

The coherent trends in radiocarbon dates in each of the main transects of the investigated reach of the Vecht reveals the direction and chronology of migration: in transect AC the migration took place from southwest to northeast (Fig. 7) and in transect HB from northeast to southwest (Fig. 8). Dates from the smaller transects (AX, HC and HE) fit into this reconstruction. This indicates that the river channel along the full length of the studied reach had a broadly synchronous single main phase of migration (Fig. 10) and that all successive phases of meander migration were broadly similar. This is supportive of the idea of a channel that began life as a linear axis, and developed meanders later. The bedload and channel fill deposits can be divided into three chronostratigraphical units (A–C): an oldest unit with an erosive character (A), a middle unit comprising extensive channel aggradation after active migration (B) and the youngest unit indicating that migration had almost stopped (C; Fig. 10).

*Unit A.* The widely varying dates for this initial phase of river migration are older than the known age of Vecht onset (based on the end of peat growth, Section 4.2.1). Moreover, in each of the cores 282 and 332, two samples were dated, the upper sample being older than the lower sample (Fig. 10). This inverted chronology indicates that there is *reworked* organic material in the samples. Unit A thus characterises an erosive phase of fluvial activity, which on the basis of the youngest date (core 332, lower sample) is younger than c. 400 BC. This sample will include *reworked* material and therefore provides a TPQ date, with a possible time gap of several centuries.

*Unit B.* A substantial part of the migration trajectory (about 50%) yielded dates from a relatively short period, the fifth to seventh centuries AD. The dated material was possibly partly *reworked* as well, so the ages may represent a maximal age. Given relatively similar ages of multiple dates, however, we believe that these dates represent channel activity quite accurately. These dates show sedimentation in the inner bend. Due to ongoing meander migration, it is not possible to pinpoint exactly the position of the channels' outer bend edge (migration front). Therefore, reconstruction of the exact channel width over time is not directly possible. It may be possible that the channel remained similar in cross-sectional wet area, while it migrated relatively fast.

**Table 1.** Results of the AMS  $^{14}\text{C}$  dating of samples from the bedload and channel-fill facies of the Vecht and from the top of the peat below the levee deposits. E-nrs. represent earlier published results (last column). N-nrs. show results of the current study. All laboratory ages were recalibrated using the OxCal v4.4.4 calibration curve (Bronk Ramsey, 2021).

nr	Lab. code	$^{14}\text{C}$ -age BP	1 sigma calibrated age	2 sigma calibrated age	Coordinates $x, y$ and surface elevation (m)	Depth (m rel. to sea level)	Sample name/nr.	Dated material	Source (transect core layer)
E1	UtC-01900	2650/50	895–786 BC	923–766 BC	133405/460280/–0.10	–0.53–0.56	Oud Zuilen I-1a	Alnus glut. wood fragm	Törnqvist (1993, 150)
E2	UtC-01901	2620/50	833–768 BC	904–569 BC	133405/460280/–0.10	–0.53–0.56	Oud Zuilen I-1b	Alnus wood fragments	Törnqvist (1993, 150)
E3	UtC-14573	2367/44	515–392 BC	746–369 BC	130221/478555/–1.44	–2.65–2.66	Weesp 1	terr macrofossils	Bos et al. (2009)
E4	UtC-14574	2920/70	1217–1014 BC	1376–922 BC	123751/464139/–1.67	–3.44–3.48	Spengen 2	terr macrofossils	Bos et al. (2009)
E5	UtC-14577	2420/50	731–407 BC	755–398 BC	128141/477478/–1.50	–2.83–2.84	Gein 1	terr macrofossils	Bos et al. (2009)
E6	UtC-14581	2280/50	400–212 BC	413–197 BC	129342/466569/–1.15	–1.87–1.88	Weeresteijn 1	terr macrofossils	Bos et al. (2009)
E7	UtC-14582	2810/50	1045–901 BC	1111–833 BC	129342/466569/–1.15	–3.32–3.33	Weeresteijn 2	terr macrofossils	Bos et al. (2009)
E8	UtC-14583	1877/60	84–236 AD	12–327 AD	126505/465671/–1.07	–1.75–1.76	Portengen 2	terr macrofossils	unpublished
E9	UtC-14584	2870/47	1120–939 BC	1204–919 BC	125711/465662/–1.63	–3.65–3.66	Portengen 3	terr macrofossils	Bos et al. (2009)
E10	UtC-14585	2352/45	513–384 BC	740–233 BC	130257/474979/–1.57	–2.22–2.26	Vecht 1	terr macrofossils	Bos et al. (2009)
E11	GrA 43211	1645/35	381–532 AD	264–539 AD	135413/457670/0.90	–0.55	Utrecht C2	Seeds	Van Dinter et al. (2017)
E12	GrA-50894	880/70	1046–1226 AD	1031–1269 AD	138220/455780/1.50	–1.12	Utrecht C13	Seeds	Van Dinter et al. (2017)
E13	UtC-14937	912/28	1047–1201 AD	1040–1212 AD	135530/457130/1.70	–0.40	Utrecht C102	Phragmites seed	Van Dinter et al. (2017)
N20	BA-512506	2690/30	896–807 BC	902–803 BC	130429/474541/–1.53	–2.43–2.45	Nigtevecht 20	Alnus twigs	this study, AC-293-6
N14	BA-498677	2160/30	350–122 BC	356–57 BC	130471/474567/–1.47	–1.97–2.03	Nigtevecht 14	Alnus seed/catkin	AC-292-6
N3	BA-481823	2240/30	381–210 BC	390–202 BC	130471/474567/–1.47	–2.07–2.23	Nigtevecht 3	Alnus etc seeds	AC-292-8
N4	BA-481824	2500/30	766–551 BC	778–520 BC	130471/474567/–1.47	–2.30–2.37	Nigtevecht 4	Alnus twig	AC-292-10
N1	BA-465924	1780/30	239–327 AD	213–361 AD	130617/474652/–1.26	–2.01–2.06	Nigtevecht 1	Wood fragments	AC-235-5
N2	BA-465925	2830/30	1015–928 BC	1107–904 BC	130617/474652/–1.26	–3.11–3.16	Nigtevecht 2	Bulk s. wood peat	AC-235-9
N19	BA-512505	4000/30	2568–2472 BC	2578–2463 BC	130635/474662/–1.18	–3.97–3.99	Nigtevecht 19	terr macrofossils	AC-238-18
N12	BA-487321	1480/30	568–636 AD	550–644 AD	130692/474693/–0.84	–3.76–3.79	Nigtevecht 12	terr macrofossils	AC-301-17
N13	BA-487322	1590/30	433–536 AD	419–548 AD	130692/474693/–0.84	–5.42–5.49	Nigtevecht 13	Alnus seed/catkin	AC-301-32
N5	BA-481825	930/30	1045–1160 AD	1032–1203 AD	130752/474711/0.65	–1.97–2.10	Nigtevecht 5	Terrestrial seeds	AC-295-11
N22	BA-512508	3490/30	1880–1751 BC	1892–1699 BC	130693/474609/–0.55	–5.26–5.45	Nigtevecht 22	Seeds and tree leave	AX-332-62
N23	BA-512509	2320/30	407–378 BC	459–231 BC	130693/474609/–0.55	–6.30–6.55	Nigtevecht 23	Seeds and tree leave	AX-332-70
N7	BA-481827	380/30	1455–1619 AD	1447–1632 AD	130755/474222/0.36	–1.09–1.24	Nigtevecht 7	Terrestrial seeds	HB-298-8
N11	BA-487320	1520/30	542–596 AD	436–637 AD	130810/474236/–0.89	–2.84–3.04	Nigtevecht 11	Terrestrial seeds	HB-296-13

Table 1. (Continued)

N6	BA-481826	1510/30	549-596 AD	440-640 AD	130828/474242/-0.94	-3.42-3.50	Nigtevecht 6	Wood twig	HB-287-12
N10	BA-487323	1600/30	425-535 AD	416-545 AD	130865/474255/-0.90	-2.20-2.30	Nigtevecht 10	Alnus seed/catkin	HB-285-8
N8	BA-483075	3090/30	1413-1302 BC	1425-1270 BC	130893/474266/-0.73	-4.38-4.48	Nigtevecht 8B	Wood twig	HB-282-17
N9	BA-481829	2530/30	781-570 BC	794-544 BC	130893/474266/-0.73	-4.59-4.73	Nigtevecht 9	terr macrofossils	HB-282-19
N16	BA-498679	1460/30	592-641 AD	564-650 AD	130967/474097/-0.90	-2.84-2.93	Nigtevecht 16	Tree leaves	HC-314-24
N21	BA-512507	1220/30	784-878 AD	687-888 AD	130834/474384/-1.02	-3.67-3.72	Nigtevecht 21	Wood twigs	HE-328-24
N15	BA-498678	2110/30	168-56 BC	339-46 BC	130296/475560/-1.08	-2.98-3.08	Nigtevecht 15	Alnus etc seeds	TW-572-5
N18	BA-498681	2600/30	803-780 BC	818-760 BC	130427/474504/-1.64	-2.03-2.05	Nigtevecht 18	Alnus wood	ProFA-3a
N24	BA-512510	2730/30	901-831 BC	928-810 BC	130427/474504/-1.64	-2.12-2.17	Nigtevecht 24	Alnus wood	ProFA-3d
N17	BA-498680	Contaminated			130795/474368/-1.04	-3.72-3.76	Nigtevecht 17	terr macrofossils	HE-327-11

Alternatively, the active channel was originally wider and narrowed during the migration. In this process, the inner bend filled in relatively fast, while the outer bend migrated at a slower pace (diminishing cross-sectional wet area). Considering its small discharge today, the present-day width of 50–60 m is considered a minimal width for medieval period, while the (minimum) channel belt width of 100 m (Fig. 6) represents the maximum original channel width. Taking into account some narrowing of the Vecht river after abandonment (Fig. 10, N5 and N7), we estimate its width during early medieval migration to have been between 70 and 90 m at maximum. This leaves only small room (about 20 m) for diminishing width of the migrating river in unit B. We therefore think that Unit B was formed under relatively fast migration predominantly, while narrowing must have been very modest.

By the sixth century AD (or possibly somewhat later), the active river likely already experienced 60% of its total migration (Fig. 10). Sample N21 (transect HE, core 328, Fig. 6, cf. Fig. 13) is on the basis of the dating result considered to be within the longitudinal component of river migration and is therefore positioned approximately into the reconstruction of river migration (oval symbol in Fig. 10). The breakthrough of the IJssel in the early Middle Ages in the area between Doesburg and Deventer may have led to a further reduction of water supply to the Neder-Rijn, Kromme Rijn and Vecht (Makaske et al., 2008; Cohen et al., 2009). Van Dinter et al. (2017, 30) reconstruct that the Vecht continued approximately equal discharge in this period, because discharge of the Oude Rijn reduced more quickly than that of the Kromme Rijn (see also Section 5.2).


*Unit C.* In the second millennium AD, meander migration was minimal. The two dates from unit C probably indicate accretion of land in the border part of the river that has decreased in significance. The virtual cessation of river migration can be linked to the closure of the Kromme Rijn at Wijk bij Duurstede in 1122 AD (Dekker, 1980). Various authors accept this year as the break point after which the Kromme Rijn, Oude Rijn and Vecht carried only local water (Berendsen & Stouthamer, 2001; Cohen et al., 2012; Van Dinter et al., 2017, 31).

### 4.3. Analysis of archaeological sites


The results of the inventory of archaeological settlements and isolated finds are shown in Fig. 11. The most spectacular find is the Roman tombstone of Loenersloot, which has probably been transported into the area from elsewhere (Kok, 2008, 43). Most of Mesolithic/Neolithic find locations are situated in the eastern part of the inventoried area, on top of the cover sands below the relatively thin peat layers. Further west, such finds may be hidden below thick peat sequences, as demonstrable near the river Kromme Mijdrecht (Fig. 11). Furthermore, finds from the Bell Beaker period which reflect habitation in the vicinity are known from the Rokin in Amsterdam, found *ex situ* more than 10 m below surface in a tidal-filled channel (Kranendonk et al., 2015).

The spatial relationship of human activity, including settlement, to channel systems can be established on the basis of finds from the Iron Age onwards. Iron Age find spots are located mainly in the first phase of this river branch towards the west (Oud Aa/Angstel). Out of five clear settlements, four are located on the fluvial ridge and one on a crevasse splay in Aetsveldse Meer. The latter one covers a strip of ground more than 500 m long and yielded many different types of finds from the middle to late Iron Age. The discovery of three dugout canoes suggests that people moved through the area by boat. In one such dugout canoe,






**Fig. 9.** a: Annotated photo of wall profile of test pit B. b: column image of the three profile recordings of test pits A, B and C with two dated samples (calibrated at 2 sigma) in profile A (for locations of test pits see Fig. 6). c: column image of cores 292 and 293 (in transect AC, see Fig. 7) with four dated samples. Three lithological units are distinguished, from bottom to top: peat, humic clay (= degraded peat) and clay. The thickness of the layer of degraded peat is indicated in red. Groundwater table near the test pits is 2.15 m below sea level. For lithological legends, see Fig. 7. For data of sample nr's. see Table 1.



**Fig. 10.** Schematic cross-section, showing transversal migration course of the river Vecht in its reach between the former Horstermeer and Aetsveldse Meer. In the trajectory, we distinguish between chronostratigraphical units A, B and C. The trajectory of the transects AC and HB from the red coloured peat border (Fig. 6) up to the current river course has been presented in percentages and combined, with date(s) being displayed in its percentual position. The highest point of the boundary between intact peat and channel fill is taken as zero. The boundary between land and water at the river Vecht is taken as a 100%.



**Fig. 11.** Overview of archaeological sites in the Angstel and Vecht system. All find locations within the range of this map are listed, with the exception of the Oude Rijn channel belt in the far south of the map. 1 = isolated find; 2 = possible settlement (up to 10 finds); 3 = settlement (10 or more finds). Red triangles with a black base refer to finds that were presumably brought in from elsewhere during civil works. For legends of geomorphological zones see Fig. 4. Map background Angstel-Vecht: Bos et al. (2009, 359). Map of other areas: conforms to geomorphological map of the Netherlands and Van Dinter (2017).



**Table 2.** Number of Roman finds by period in the study area of Angstel and Vecht (Fig. 11).

Period	Coins	Other finds	Percentage
Before 12 BC	3	–	11
Early Roman	1	3	15
Middle Roman	9	4	48
Late Roman	7	0	26
Roman indeterminated	–	7	–

found at Nigtevecht in 1987, pottery sherds typologically datable to c. 600 BC were found, showing similarities with pottery from Assendelft (north of the Oer-IJ). Also the stylistic characteristics of pottery found in the Breukelerwaard point to contact with the Oer-IJ area (Kok, 2008, 42).

The biggest difference in the eastern second phase (Vecht) corridor (with the studied reach) compared to the Angstel phase is the complete absence of Iron Age finds along this reach of the Vecht. No Iron Age find spots are known here and in contrast Roman period finds are represented more often, especially between Breukelen and Nigtevecht, although only two of the eight find spots suggest a settlement. Isolated finds from the Roman period for the most part consist of coins found by metal detection. As a consequence, finds from the Roman period may be over-represented in the data. Moreover, after the Middle Ages significant amounts of soil from Utrecht were imported into the Angstel–Vecht area for several purposes (Stouthamer *et al.*, 2008, 23; Kok, 2008, 44). Finds of Roman coins that are assumed to originate from elsewhere are marked in Fig. 11. The relative period distribution of the Roman finds is shown in Table 2 and, taking into account that it mainly concerns coins, it seems to fit within the pattern known from the river area, although the number of finds in the river area is much greater. Because of the limited Roman find material in the Angstel–Vecht area, there should be attached less significance for dating the onset of the Vecht to the Roman finds in the studied reach than to the lack of finds from the Iron Age.

## 5. Discussion

In this section, we will address the question of the character of possible Roman interventions and test our hypothesis, based on the results of  $^{14}\text{C}$  dating within the context of our detailed mapping. We will also discuss the development of the investigated river reach in a broader context, within the Angstel and Vecht channel belt, the natural system of the Rhine–Meuse delta and the Oer-IJ/Vlie estuaries. Understanding this wider landscape context provides insights into how the Romans may have operated in these areas.

### 5.1. Chronological control

What additional insights do the newly acquired chronological data, augmented by previous information, provide us with regarding the palaeogeographical development of the studied part of the Vecht river system? The dates of the most relevant events are plotted together in Fig. 12. Our dating results for the Angstel (phase 1) confirm those of Törnqvist and Bos *et al.* (Section 4.2.1.). In the erosive first stage of Vecht river (phase 2) activity, a general TPQ age has been determined. The youngest sample of *reworked* material from this erosive chronostratigraphical unit A of the channel fill deposits (Fig. 10, core 332) provides a TPQ date of 459–231 BC (Table 1, N23). Our results of migration stage B, dated

to the fifth to seventh century AD (Fig. 10), and final stage C of the Vecht river activity confirm the relative high early medieval discharge of the Vecht river (compared to the Oude Rijn) suggested by Van Dinter *et al.* (2017, 26, 30) and their dating of the final stage of this river, caused by the damming of the Kromme Rijn at Wijk bij Duurstede in 1122 (Fig. 12).


When did peat growth cease completely near the studied Vecht reach (phase 2), in response to the formation of levee deposits by this river? Our dating of the youngest preserved peat at Nigtevecht (Fig. 12, N14, 356–57 BC) differs significantly from the dating by Bos *et al.* (2009, 368) at Nigtevecht (E10, 740–233 BC; the statistical overlap is only 4%). With regard to the origin of the Vecht (phase 2), our date added to those of Bos *et al.* pushed the TPQ date for the onset of Vecht activity from (approximately) 500–200 BC to 350–100 BC. Above the youngest peat (Fig. 9, core 292, N14, 356–57 BC), there is still 8 cm of decomposed clayey peat, while in profile A there is 21 cm of decomposed clayey peat overlying the top of preserved peat, which is from around 800 BC. The new result from our study is the degradation of the peat, as inferred from the dating results and observed lithology, which reinforces the TPQ character of the dating results of Bos *et al.* and our study, with a possibly longer time gap. In the studied reach, there are good indications that peat growth continued until the arrival of the Romans, possibly even longer. With respect to the Loenen avulsion (whether natural or man-made, see Section 2.2/ Fig. 4), it is worth noting that between the avulsion point and the study area, sediments were firstly deposited in the Horstermeer before they could be spread over the peat as natural levee deposits along the studied reach of the Vecht (Bos *et al.*, 2009, 368). Our data did not provide sufficient information about possible separate episodes of avulsions (whether or not natural) near Breukelen and Loenen.

Dating results of unit B (Fig. 10) indicate that by 400 AD the Vecht river existed and was actively depositing Rhine-clay across the peatland (Fig. 6/ Fig. 13). The sample with abundant wood fragments (Fig. 7, core 235; Table 1, N1),  $^{14}\text{C}$  dated to 213–361 AD and interpreted as floating woody material deposited on the levee by the river Vecht, is an indication that levee formation was already going on in the third/fourth century AD. Also relevant is a dating from the Oud-Aa channel fill after it was abandoned as a part of the Angstel phase 1 (Table 1, E8, 12–327 AD). With this, there is some limited room (400 to max. 500 years) of past peat formation now lost in the peat-degradation hiatus.

The results of the inventory of archaeological find locations show that along the Vecht (phase 2) there are no Iron Age remains, but there are some of Roman date. This is in great contrast to the Angstel (phase 1), where five clear Iron Age settlements are present (section 4.3). The spatial distribution of finds along the Vecht in the Roman period is more in line with the general picture of the area several tens of kilometres north of the Roman *limes*, which suggests that during the first and second centuries AD there were only sparse native settlements in this area. The finds suggest potential occupation on the natural levees of the Vecht at Overmeer during the Roman period (second century AD), but the limited amount of finds indicates that this interpretation must be treated with caution. More significance for dating the onset of the Vecht should be attached to the lack of finds from the Iron Age.

It appears that the initiation of this reach of the Vecht lies between c. 200 BC and c. 250 AD. These dates do not discount its potential origin as a Roman canal built by Drusus but do not provide sufficient evidence either. Considering the decomposition of the top of the peat, the chronological limits together with





**Fig. 12.** Ages of the most relevant geomorphological, historical and archaeological events in the Angstel–Vecht area. Legend: 1 = calibrated <sup>14</sup>C age of earlier results, with 1 and 2 sigma probability; 2 = idem, results of this study; 3 = time period composed of several ages; 4 = habitation period, based on archaeological data (sources: supplementary Appendix C and Bos et al. 2009, 370); 5 = historical event, mentioned in text. All radiocarbon ages mentioned in this study were recalibrated using the OxCal v4.4.4 calibration curve (Bronk Ramsey, 2021).

geomorphological, archaeological and historical arguments make the possibility of canal construction in the Roman period more likely.


### 5.2. Channel geometry and its evolution

The mapping shows a narrow channel belt (c. 150 m wide) with a meander shape with low sinuosity, through which within the borders of the channel belt one straight axis extending 2 km in length can be drawn for the purpose of analysis (Fig. 13). The western boundary between channel deposits and preserved peat (near transect AC) has a slightly curved shape, interpreted here as a migration to the left (west) during an initial short period. Alternatively, it could be that a hypothetical Roman canal was not constructed in a perfectly straight line. Other lowland Roman canals – the c. 30-km long Corbulo Canal (De Kort & Rackzynski-Henk, 2014, 56; Polak et al., 2019; Hessing & Schrijvers, 2021; Fig. 2) and the Fossa Augusta, Fossa Flavia and Fossa Claudia, which covered together a distance of c. 150 km in the Po delta in Italy (Uggeri, 1997; Laurence, 1999, 117; Medas, 2013, 2017) – also consist of rather straight sections that were connected with slight bends, although also was made use of existing tidal channels or lagoonal lakes. The Canal of Corbulo was dug through the peat directly behind a beach barrier, so that the morphology of the edge of this landform has influenced the shape of the canal, resulting in straight sections connected with slight bends. Assuming Drusus canals in the Vecht

area, they were dug to help military campaigns in an offensive period as a series of short-cuts to make transportation of troops and supplies easier. This differs from, for example, the longer canals of Corbulo and in the Po delta, which were established to address long-term infrastructure needs such as the transportation of goods between different areas.

‘Identification of the construction phase of a (Roman) canal can be difficult, depending on the post-depositional processes that affect the state of preservation of the canal’ (Salomon et al., 2014, 3). During our investigation, we could not identify any direct evidence of a canal, such as revetments or created cut surfaces, known from Roman canals which usually have a width of 12–15 m (De Kort & Rackzynski-Henk, 2014, Table 1). The channel belt boundary identified in transects AC and AX (Fig. 13) could date from a relatively short timeframe after the assumed canal construction. Based on our oldest age results of channel belt sedimentation (Fig. 10, unit B), we know that the initiation of this reach of the river or canal must in any case be before approximately 400 AD. However, the initial period of two-way widening (to 50–70 m) and vertical deepening of the channel before starting to migrate, and the sample N1 (Fig. 7, core 235; 213–361 AD) taken within the levee deposits, are indications that levee formation was already going on in the third/fourth century AD (Section 5.1).

The narrow width of the Vecht channel belt between Overmeer and Nigtevecht is mainly related to the limited impact (low sinuosity) of the river’s meandering activity (Bos et al., 2009, 361-D, ch.



**Fig. 13.** Geomorphology of the investigated reach of the Vecht with the location of a potential Roman canal, which is also the axis of the successive medieval meanders between former Horstermeer and Aetsveldse Meer. The arrows indicate the migration direction of the river and meander loops. The red dashed line is the estimated position of the river around 5th–6th century AD. See also Fig. 6. In the former Horstermeer, the map shows a LiDAR image with a corresponding residual channel.

5.3.3). Meander activity is much lower in peaty areas (due to the higher erosion resistance) than where a river flows through a clay- and sand-filled lake (Berendsen & Stouthamer, 2001; Bridge, 2003; Makaske & Weerts, 2005; Candel, 2020). It could be questioned whether this reach of the Vecht is a 'straight river', which is defined as a slightly winding river with low sinuosity due to a high erosion resistance of the encasing substrate (Berendsen, 2005, 272). Mostly the centre line through the successive windings has at least a slightly curved shape, but in the studied reach of the Vecht the initial stage was very straight, also for peaty rivers, so we think it is an exceptional one (Fig. 13). Note that, based on the migration of the meanders, it can be estimated that the distance from the axis to the outer bends of the meanders, which is approximately 130 m (6.5%) today, was only approximately 70 m (3.5%) around the fifth to sixth century AD (Fig. 13). In a natural situation, the main structure (centre line through the meander loops) usually has a (slightly) curved shape, for example, perimarine crevasses along the Rhine (Van Dinter, 2013), but in the case of V1 it seems to be a really straight line, which raises the hypothesis of canal construction (Fig. 5, V1; cf. Fig. 13).

Apart from the notable channel belt geometry and the difficulties in dating the origin of the investigated reach of the Vecht, the question is what the landscape looked like through which the canal was dug. A small peat river navigable for canoes might have existed here, similar to examples described by Candel (2020 and underlying references), hardly depositing inorganic sediments over the peat, but this is not that likely. The western side of Horstermeer and Aetsveldse Meer where the Angstel established its course earlier, would have accommodated such streams, whereas the more eastern side appears a stream-less fen-peatland through which runoff was more diffuse (Van Loon, 2010). If such diffuse courses existed, the Romans could have converted them into a navigable route for flat-bottomed boats, but why would they have aimed for it? Why didn't they use the Angstel river a bit further west? Because that was blocked/closed off? At the time of arrival of Drusus, if the Angstel river could only be used by canoes and not by Roman flat-bottomed boats, some kind of human intervention was necessary.

To discuss the hypothesis of a dug canal, it is also relevant to consider the expenditure of time and resources. Digging a new route through a peat-forming fenland might have taken less effort than dredging and keeping open winding older river courses. The latter were prone to infilling, with problems in digging like heavy vegetation and slumping of clayey shores by seepage, being less stable than peaty shores. This is a plausible assumption for general human use of wetland landscapes and may have been one of the reasons for creating a new (more eastern) and navigable route in addition to the Oud-Aa and Angstel river course. This also fits in with the abandonment of the crevasse splays and the Nifterleek in the Aetsveldse Meer and the timing of abandonment of the Oud-Aa (Section 5.1, Fig. 12). Another example in which a canal was dug through peat next to a natural course is the Corbulo Canal (Hessing, 1990, 342; Jansen, 2011, 22; De Kort, 2013, 235).

The discharge of the Vecht near Nigtevecht in the Roman and early medieval period can be estimated as analogous to the estimates for the Kromme Rijn, Oude Rijn and Vecht near Utrecht (Van Dinter et al., 2017, 25 ff.). A reconstruction of the discharge of these three rivers has been derived from channel geometry, that is, cross-sectional wet area of these rivers near Utrecht, assuming an equal energy slope for these rivers near Utrecht.

For proper estimation and comparison, it is necessary to use the available data of the dimensions of the active Vecht channel. The


present-day Vecht open river near Nigtevecht still has a width of 50–60 m, while the river cannot have been wider than 100 m, the minimum width of the channel belt (Fig. 13). Taking into account some narrowing of the Vecht river before and after abandonment (Section 4.2.2), we estimate its original width (in case of a possible dug canal: after maximum two-side widening, deepening and having started meandering) to have been between 70 and 90 m at maximum. Based on Fig. 10, we estimate its depth (measured from the top of the peat, Fig. 10) to be about 4 m, which gives a cross-sectional wet area of about 200 m<sup>2</sup>. This is higher than Van Dinter's estimate near Utrecht (100 m<sup>2</sup>). A plausible explanation for the difference is that in the peatland area near Nigtevecht the gradient (energy slope) will be lower than in a river delta area like near Utrecht, resulting in a lower ratio of the discharge versus the cross-sectional wet area.

Reconstruction of discharge near Utrecht through time, from the beginning of the Angstel avulsion to the damming of the Kromme Rijn by Van Dinter et al. (2017, 28), is shown in Fig. 14. Using our estimate near Nigtevecht (section 4.2.2) and taking into account the lower energy slope, we supplemented this with an approximate reconstruction of the discharge of the Vecht near Nigtevecht (Fig. 14). The red curve shows the result when using the same energy slope as near Utrecht, as a maximal discharge. Given the fact that this discharge is higher than that of the Kromme Rijn (its feeding channel) just after 500 AD, this is likely an overestimation. The blue curve, based on a lower energy slope in the area near Nigtevecht, fits better with the available data. The blue reconstruction, with only slightly diminishing Vecht discharge up from fifth century AD (based on dates from unit B; Fig. 10), supports the suggestion by Van Dinter et al. (2017, 30) that in this period a larger share of the water from the Kromme Rijn was discharged via the Vecht than via the Oude Rijn. Alternatively, given that our unit B dates are TPQ ages, it cannot be excluded that the meander migration took place some centuries later. In that case, it might be correlated with an increased meandering of the Vecht near Utrecht in the second half of the 10th century and the 11th century, described by Van Dinter et al. (2017, 27), although they present no enhanced discharge in their reconstruction (Fig. 14). We also reconstructed the initial stage (of the Vecht phase 2) having two possible origins (a = natural or b = human-induced). In case of a natural origin, we expect the initiation to have taken somewhat more time, and therefore Vecht activity starts earlier in our reconstruction.

### 5.3. The Romans and the Vecht

The starting point for our research was the Vecht as the best candidate of multiple hypotheses for the location of the Drusus Canal(s). Amongst modern authors, there was already a widely shared opinion that the Vecht was already navigated by the Romans (see introduction). But this needs not necessarily to be explained as the Romans navigating the Vecht as a natural river. The Vecht may as well have been navigated by the Romans because of canal construction in its reaches. It is striking that none of the previous authors has made a direct association with the Canals of Drusus. Some authors even took the position that the Drusus' Canal(s) had been constructed in the area of the IJssel, but that the Romans had also used the Vecht as a navigation route (Willems, 1981/1984, 58; Manten, 2007, 63). This raises the question as to whether the Romans used more than one shipping route to the north, a suggestion that might be understood from the use of the plural '*fossae*' by Suetonius (*Vita divi Claudii* I, 2–4). However,





**Fig. 14.** Estimated discharge proportions based on cross-sectional wet area of Kromme Rijn, Oude Rijn and Angstel/Vecht near Utrecht (black) and near Nigtevecht (red/blue) as a function of time. The blue graphic has been corrected for the lower energy slope in this area. a = variant on the basis of a natural origin of the phase 2 channel belt; b = variant on the basis of an artificial intervention by the Romans; c = estimate with uncertainty range of maximum discharge. Source: Van Dinter *et al.* (2017), Fig. 10 (black), supplemented with the results of this study (red/blue).

the Romans needed only one shipping route, so why would they make big efforts to construct a second one?

It seems more plausible to explain the plural as a series of canals connected in succession in the area of the Vecht and maybe the Flevo Lake and the Vlie; together, they may have formed a single navigable route, e.g. in the places indicated with red ovals in Fig. 4. Among these R1 canals, there also can have been a shortcut in the area of the Flevo Lake and the Vlie (Huisman, 1995, 191; see Fig. 2). The entire navigable route could have been created by Drusus with a series of passages and short-cuts of river bends from Vechten to the great lakes and the land of the Frisians (Halbertsma, 1982, 46).

Finally, there must have been an easily navigable river to the north, because in the middle of the first century AD, several classical authors mention a right branch of the Rhine that flowed through the Flevo Lake; before that time, only the Rhine and the Waal were mentioned as river branches (Section 2.1). Ptolemy (87–168? AD) even mentions three Rhine mouths north of the Helinium (Meuse–Waal estuary) and Lugdunum Batavorum (Katwijk aan Zee), of which the most northern is explicitly called Flevum (Vlie). (Ptolemy, *Geographia* II, 9, 12).

The descriptions by the classical authors fit well with a series of canals. Through the input of extra water from the Rhine, they would have developed into a naturalised, freely flowing channel connection between the Rhine and (eventually) the Wadden Sea; the energy of the river meant that it did not have to be kept open by other means such as dredging. A sufficient influx of river water into the Vecht depended primarily on the character of the bifurcation of the Oude Rijn and Angstel/Vecht channel belts (Fig. 2/4). Whether or not the Romans also made channel adjustments at this bifurcation to ensure sufficient water inflow into the Vecht (e.g. shortcutting bends or constructing a groyne, like at Herwen, Fig. 2), is hard to determine, as old river phases have been eroded by or may be hidden below younger ones (see Van Dinter *et al.*, 2017, ch. 4.1.1.).

The *castellum* of Vechten, named Fectio, founded c. 5 BC (Zandstra & Polak, 2012, 243, 249; Polak, 2014, 75, 93) has been assumed to have been built near the bifurcation of the Rhine and Vecht (Polak & Wynia, 1991; Bechert & Willems, 1995, 81; Van der Tuuk & Cruysheer, 2014, 106). It had a military logistical function in the fleet expeditions of Tiberius and Germanicus to northern Germany (Zandstra & Polak, 2012; Polak, 2014). Van Dinter (2013, 20) reconstructs the bifurcation point of the Rhine and Vecht during the early first century AD in the city of Utrecht, presumably adjacent to the Roman fort (Traiectum) at Utrecht. This *castellum* was built c. 40 AD (Broer & De Bruijn, 1997; Zandstra & Polak, 2012; Polak & Kooistra, 2013; Polak, 2014). At that time, the strategic route via the Vecht to the north had become less important to the Romans and their efforts were focused on guarding the Rhine *limes* as a river transport corridor and supplying Britannia via the Oude Rijn (Graafstal, 2009, 186; Breeze *et al.*, 2018, 67). The older *castellum* of Vechten was situated at the outer side of the Oudwulvenbroek meander (Fig. 4), which was abandoned in the early Roman period as the result of a meander cut-off c. 1.5 km upstream of the *castellum* (Van Dinter *et al.*, 2017, appendix, 42; Van Dinter *et al.*, 2013, 20 and underlying references). But what if this was not the result of natural processes, but of an intervention by the Romans (the first of a series of successive cut-off canals by the Romans, including the studied Vecht reach near Nigtevecht), and if the cut-off point is the place where, according to Tacitus, Germanicus entered a canal named after his father Drusus?

Since, according to Suetonius, Drusus had dug canals at the northern side of the Rhine, it is almost obvious that he (also) made adjustments to the system of the Angstel and Vecht. Although Suetonius speaks about an enormous work (*immensi operis*), Drusus may have achieved this by digging a number of relatively short canals, so that in a short time he established a navigable connection to the Wadden Sea. A second Roman waterway via the

Beneden-IJssel area is not plausible, because locks would have been required there and the Romans only needed one waterway towards the north. The Dam and the Canals of Drusus can be considered to be parts of one coherent design. This makes it an excellent example of Roman shipping infrastructure, which must have been devised at a high central political level, like the projects which undoubtedly were undertaken elsewhere in the Roman Empire. If the focus of further investigations, such projects can yield important information if undertaken using a combined geoarchaeological and historical approach.

This study has revealed the potential course of a canal system dug by Drusus, which demands further interdisciplinary geoarchaeological research. Elements of this research can be:

- \*refining the dating results of Vecht development and end of peat growth by investigation in long profile trenches at suitable locations, especially where top of the peat may be better preserved (under dikes and former buildings);
- \*continued study to refine dating of the abandonment of the Oud-Aa near Breukelen;
- \*research with a morphological approach at other possible Roman canal locations in the Vecht area (e.g. near the cut-off point northeast of the *castellum* of Vechten).

## 6. Conclusions

This study has undertaken a detailed geoarchaeological investigation of the reach of the Utrechtse Vecht between Overmeer and Nigtevecht, which has a strikingly straight shape. The working hypothesis of this investigation was that the Romans created a navigable route of canals for flat-bottomed boats in this area by connecting a number of peat-bounded lakes in the Angstel–Vecht, which subsequently developed into a naturalised river course. The detailed mapping of the studied Vecht channel between Overmeer and Nigtevecht shows an overall very straight main channel planform over a length of 2 km with very low sinuosity. The results indicate that the successive meanders of the studied reach of the Vecht channel formed quite steadily in the same way during the same time period. This is supportive of the idea of a channel that began along a straight axis, but became meandering later.

Our top of the peat dates and the inferred first clay influx in the peat confirm the onset of the phase 1 Angstel around 900 BC. Our dating of the younger Vecht branch onset (phase 2) does not contradict earlier dating results by Bos et al. (2009), although our dating results are slightly younger. However, in our study visual inspection in test pits has revealed peat degradation, that must be taken into account when interpreting chronologies. By this we know that the end dates of peat growth (both for the earlier and our datings) tend to be interpreted as older than they actually are, so that Roman age becomes more probable. We also confirmed that the abandonment of the Angstel–Vecht system took place around 1100 AD.

Three chronostratigraphic units of river development with migration are distinguished in the channel deposits by dating organic remains from the channel belt: an active and erosive first unit, starting some shorter or longer time after about 200 BC; a second unit with extensive channel fill deposition after and during active river migration from about 400–900 AD; a final unit associated with the damming of the Kromme Rijn in 1122 AD (and fossilisation). Dates from fills of the Oud-Aa (phase 1) channel and woody material deposited on the levee of the Vecht indicate

that river activity must have started well before 250 AD. Based on our dating evidence and that of previous studies, the onset of the Vecht river (the Loenen–Nigtevecht–Muiden branch) may lie between 200 BC and 250 AD, which makes a Roman age more likely. Combined with our results on the initial straight shape of the studied reach Overmeer–Nigtevecht, this gives more room (time) for the possibility of canal digging activities at the time of Drusus. This study has therefore made the potential course of a series of canals dug by Drusus in this area more likely. Confirming this course demands further interdisciplinary research.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/njg.2022.2>

**Acknowledgements.** This study was part of the PhD project of the first author on Roman waterworks in the Rhine–Meuse delta. The authors are grateful to Feike Miedema, Jurgen de Kramer and Eckhart Heunks for their cooperation in the fieldwork, to Nico Willemse and Laura Boukje Stelwagen (RAAP, NL) for their help with the core profile presentation, to Anton Cruysheer and Stijn Heeren for their help to collect data of archaeological finds and to Henk van Haaster (Biax) for the botanical analysis of the samples. We especially want to thank Kim Cohen (Utrecht University) for his valuable feedback on interpretation and wording of our palaeogeographical results. Furthermore, we are grateful to Kees Kasse (VU University), Peter Vos (TNO) and Jasper Candel (Wageningen University) for discussions on different aspects of this study. Andy Howard (Landscape Research & Management, UK) is thanked for reviewing the English grammar of this paper. Our thanks are also due to Rien Polak, Kim Cohen and one anonymous reviewer that have improved the quality of our paper. We are grateful to the province of Gelderland and the VU University Amsterdam for making this study possible. Last but not least, we want to thank Nico Roymans for his stimulating feedback on the analyses, interpretations and conclusions of this study.

## References

- Bechert, T. & Willems, W.J.H., 1995. Die römische Reichsgrenze zwischen Mosel und Nordseeküste. Theiss-Verlag (Stuttgart): 120 pp.
- Berendsen, H.J.A., 2005. Fysisch-geografisch onderzoek: thema's en methoden. Van Gorcum (Assen): 319 pp.
- Berendsen, H.J.A. & Stouthamer, E., 2001. Palaeogeographic development of the Rhine-Meuse delta, The Netherlands. Van Gorcum (Assen): 130 pp.
- Blackman, D.J., 2008. Sea transport, part 2: Harbours. In: Oleson, J.P. (ed.): The Oxford Handbook of Engineering and Technology in the Classical World. Oxford University Press (Oxford): 638–670.
- Borger, G.J. & Kluiving, S.J., 2017. The wet heart of the Netherlands. In Kluiving, S.J., Kootker, L. & Hermans, R. (eds): Interdisciplinarity between humanities and science; a Festschrift in honour of Prof. Dr. Henk Kars. Sidestone Press (Leiden): (Clues 2): 37–54.
- Bos, I.J., Feiken, H., Bunnik, F.P.M. & Schokker, J. 2009. Influence of organics and clastic lake fills on distributary channel processes in the distal Rhine-Meuse delta (the Netherlands). Palaeogeography, Palaeoclimatology, Palaeoecology 284: 355–374. <https://doi.org/10.1016/j.palaeo.2009.10.017>
- Bos, I.J., 2010. Architecture and facies distribution of organic-clastic lake fills in the fluviodeltaic Rhine-Meuse system, the Netherlands. Journal of Sedimentary Research 80: 1–19. <https://doi.org/10.2110/jsr.2010.035>
- Bosman, A.V.A.J., 1999. Velsen 1 and Velsbroek B6, the latest excavations, results and interpretation from features and finds. In: N. Gudea (ed.): Roman Frontier Studies XVII. Proceedings of the 17th International Congress of Roman Frontier Studies:303–308.
- Bosman, A.V.A.J., 2016. Rome aan de Noordzee. Burgers en barbaren te Velsen. Sidestone Press (Leiden): 136 pp.
- Bosman, A.V.A.J., Bosman, W.J. & de Weerd, M.D., 1998. De vroeg-Romeinse basis Velsen 2. Het onderzoek in het tracé van de drinkwatertransportleiding in de Zuiderscheg in september 1997. Instituut voor Pre- en Protohistorie (Amsterdam): 57 pp.

- Breeze, D.J., Jilek, S., Graafstal, E.P., Willems, W.J.H. & S Bödecker, S.**, 2018. *Frontiers of the Roman Empire. The lower German limes*. Sidestone Press (Leiden): 143 pp.
- Bridge, J.S.**, 2003. *Rivers and floodplains. Forms, processes and sedimentary record*. Blackwell Publishing (Oxford).
- Broer, C.J.C. & de Bruijn, M.W.J.**, 1997. *Antonia, Wiltenburg, Traiectum. De kennis van het Romeinse verleden van Utrecht door de eeuwen heen*. Jaarboek Oud-Utrecht 1997: 97–124 (Utrecht).
- Bronk Ramsey, C.**, 2021. *OxCal 4.4.4*. (Oxford).
- Candel, J.H.J.**, 2020. *Ahead of the curve. Channel pattern formation of low-energy rivers*. PhD thesis Wageningen University (Wageningen). <https://doi.org/10.18174/506616>
- Chevallier, R. & Field, N.H.**, 1976. *Roman Roads*. Batsford (London): 272 pp.
- Cohen, K.M., Stouthamer, E., Hoek, W.Z., Berendsen, H.J.A. & Kempen, H.F.J.**, 2009. *Zand in Banen - ZanddiepteKaarten van het Rivierengebied en het IJsseldal in de provincies Gelderland en Overijssel (3<sup>rd</sup> ed.)*. Provincie Gelderland (Arnhem): 133 pp.
- Cohen, K.M., Stouthamer E., Pierik, H.J. & Geurts, A.H.**, 2012. *Rhine-Meuse Delta Studies' Digital Basemap for Delta Evolution and Palaeogeography*. Dept. Physical Geography, Utrecht University (Utrecht); Data Archiving and Networked Services (DANS). <https://doi.org/10.17026/dans-x7g-sjtw>
- de Bont, C.**, 2015. *Digging the river: the historical geography of the Amstel area (800–1275 AD)*. *Netherlands Journal of Geosciences* **94**: 353–360. <https://doi.org/10.1017/njg.2014.42>
- de Gans, W.**, 2015. *The geology of the Amstel river in Amsterdam (Netherlands): man versus nature*. *Netherlands Journal of Geosciences* **94**: 361–373. <https://doi.org/10.1017/njg.2014.41>
- de Kort, J.W.**, 2013. *Het kanaal van Corbulo. Onderzoek naar een Romeinse waterweg in de gemeente Leidschendam-Voorburg tussen 1989 en 2010*. Westerheem 62: 233–243.
- de Kort, J.W. & Rackzynski-Henk, Y.**, 2014. *The Fossa Corbulonis between the Rhine and Meuse estuaries in the Western Netherlands*. *Water History* **6**: 51–71. <https://doi.org/10.1007/s12685-014-0097-3>
- Dekker, C.**, 1980. *De dam bij Wijk*. In: *Scrinium et scriptura. Opstellen betreffende de Nederlandse geschiedenis aangeboden aan Prof. Dr. J.L. van der Gouw, bij zijn afscheid als buitengewoon hoogleraar in de archiefwetenschap en in de palaeografie van de veertiende tot de zeventiende eeuw aan de Universiteit van Amsterdam*. Erven Van der Kamp (Groningen): 248–266.
- Eck, W.**, 2007. *Der Rhein als Lebensader der Städte des römischen Germaniens*. In: C. Ohlig (ed.): *Von der cura aquarum bis zur EU-Wasserrahmenrichtlinie – Fünf Jahre DWHG. Schriften der Deutschen Wasserhistorischen Gesellschaft*, Band 11/1: 106–126.
- Erkens, G.**, 2009. *Sediment dynamics in the Rhine catchment. Quantification of fluvial response to climate change and human impact*. *Netherlands Geographical Studies* 388, Utrecht.
- Graafstal, E.P.**, 2002. *Logistiek, communicatie en watermanagement, over de uitrusting van de Romeinse rijksgrens in Nederland*. *Westerheem* 51: 2–27.
- Graafstal, E.P.**, 2009. *River Frontiers or fortified corridors?* In: Hodgson, N., Bidwell, P. & Schachtmann, N. (eds): *Proceedings of the XXI International Congress of Roman Frontier Studies (Limes Congress) held at Newcastle upon Tyne in August 2009*. *Roman Frontier Studies* 2009: 186–193.
- Grewe, K.**, 2008. *Tunnels and Canals*. In: Oleson, J.P. (ed.): *The Oxford handbook of engineering and technology in the classical world*. Oxford University Press (Oxford): 319–336.
- Halbertsma, H.**, 1982. *Friesland's oudheid*. Dissertatie Rijksuniversiteit Groningen (Groningen).
- Hessing, W.A.M.**, 1990. *Leidschendam: Rietvink*. In: Hessing, W.A.M. (red.). *Archeologische kroniek van Holland over 1989, II Zuid-Holland*. (Holland), vol. 22(6): 342–343.
- Hessing, W.A.M. & Schrijvers, R.S.** 2021. *Archeologische waarnemingen van het kanaal van Corbulo tussen Leiden en Rijswijk. Bureaustudie ten behoeve van de waardering van het kanaal van Corbulo als onderdeel van het nominatiedossier UNESCO Werelderfgoed*. Amersfoort (Vestigia rapport V1721).
- Hoogland, F., Roelandse, A.S., de La Loma González, B., Waterloo, M.J., Mooij, P.W., Verhagen, S.A. & Velstra, J.**, 2020. *Investigating the effectiveness of drain infiltration to minimize peat oxidation in agricultural fields in Flevoland, the Netherlands*. *Proceedings IAHS* 382: 747–753.
- Huisman, K.**, 1995. *De Drususgrachten: een nieuwe hypothese*. *Westerheem* **44**: 188–194.
- Jansen, B.**, 2011. *Plangebied Vlietwijk gemeente Voorschoten. Archeologisch vooronderzoek: een bureau- en inventariserend veldonderzoek*. RAAP rapport nr. 2485 (Weesp).
- Jansma, E., van Lanen, R. & Pierik, H.J.**, 2017. *Travelling through a river delta: a landscape-archaeological reconstruction of river development and long-distance connections in the Netherlands during the first millennium AD*. *Medieval Settlement Research* **32**: 35–39.
- Kehne, P.**, 2018. *Zur Erforschung der Germanicusfeldzüge, zu den Ursachen für die Unmöglichkeit ihrer Rekonstruktion und zu den Problemen des Germanicus-Bildes*. In: Burmeister, S. & Ortisi, S. (eds): *Phantom Germanicus. Spurensuche zwischen historischer Überlieferung und archäologischem Befund*. *Materialhefte zur Ur- und Frühgeschichte Niedersachsens*, vol. 53: 31–94.
- Klee, M.**, 2010. *Lebensadern des Imperiums. Strassen in römischen Weltreich*. Theiss Verlag (Stuttgart).
- Kluyving S.J. & Borger, G.J.**, 2015. *Fluvial History of the Vecht-Amstel area in the Holocene: Interdisciplinary collaborations, controversies and progress*. *Netherlands Journal of Geosciences* **94**: 289–293. <https://doi.org/10.1017/njg.2015.31>
- Kok, R.S.**, 2007. *Wat de bodem vertelt over de vroegste bewoningsgeschiedenis van de Vechtstreek en de rol die de amateur-archeoloog daarbij kan spelen*. *Tijdschrift Historische Kring Breukelen* **22**(2): 49–62.
- Kok, R.S.**, 2008. *De IJzertijd en Romeinse Tijd van de Vechtstreek*. Jaarverslag AWN-afdeling Naerdincklant **2008**: 34–54.
- Kranendonk, P., Kluyving, S.J. & Troelstra, S.R.**, 2015. *Chrono- and archaeostratigraphy and development of the river Amstel: results of the North/South underground line excavations, Amsterdam, the Netherlands*. *Netherlands Journal of Geosciences* **94**: 333–352. <https://doi.org/10.1017/njg.2014.38>
- Laurence, R.**, 1999. *The roads of Roman Italy: Mobility and cultural change*. Abingdon. <https://doi.org/10.4324/9780203062418>
- Lendering, J. & Bosman, A.**, 2010. *De rand van het Rijk. De Romeinen en de Lage Landen*. Athenaeum-Polak & Van Genneep (Amsterdam): 303 pp.
- Levison, W.**, 1905. *Vita Bonifatii auctore Willibaldo, Vitae Sancti Bonifatii Archiepiscopi Moguntini, Scriptores rerum Germanicarum in usum scholarum ex Monumentis Germaniae Historicis separatim editi*. Hannover/Leipzig.
- Makaske, B. & Weerts, H.J.T.**, 2005. *Muddy lateral accretion and low stream power in a sub-recent confined channel belt, Rhine-Meuse delta, central Netherlands*. *Sedimentary Geology* **52**(3), 651–668.
- Makaske, B., Maas, G.J. & van Smeerdijk, D.G.**, 2008. *The age and origin of the Gelderse IJssel*. *Netherlands Journal of Geosciences – Geologie en Mijnbouw* **87**: 323–337.
- Manten, A.A.**, 2007. *Hebben de Romeinen ingegrepen in de loop van de Vecht bij Breukelen?* *Tijdschrift Historische Kring Breukelen* **22**: 63–66.
- Medas, S.**, 2013. *La navigazione interna lungo l'arco fluvio-lagunare dell'Alto Adriatico in età antica tra Ravenna, Altino e Aquileia*. In: Bonifacio, A. & Caniato, G. (eds): *Barche tradizionale della laguna Veneta*, Venezia: 107–129.
- Medas, S.**, 2017. *La navigazione lungo le idrovie Padano in epoca Romana*. In: Cantoni, G. & Capurso, A. (eds): *On the Road, via Emilia 187 AC – 2017* (Parma): 146–161.
- Mirschenz, M.**, 2018. *Römische Häfen am Rhein. Einde Herausforderung für die Forschung*. In: C. Eger (ed.): *Warenwege – Warenflüsse. Handel, Logistik und Transport am Römischen Niederrhein*. (Xantener Berichte 32). Nünnerich-Asmus Verlag (Oppenheim am Rhein): 247–269.
- Morhange, C., Marriner, N., Bony, G., Flaux, C., Gaième, M. & M. Kouka**, 2017. *Geoaerchaeology of ancient harbours in lagoonal contexts: an introduction*. In: Franconi, T.V. (ed.): *Fluvial Landscapes in the Roman World*. (Journal of Roman Archaeology Supplementary Series no. 104). (Portsmouth): 97–110.
- Nederlands Normalisatie-instituut**, 1989. *Nederlandse Norm NEN 5104, Classificatie van onverharde grondmonsters*. Nederlands Normalisatieinstituut (Delft).
- Pierik, H.J., Stouthamer, E., Schuring, T. & Cohen, K.M.**, 2018. *Human-caused avulsion in the Rhine-Meuse delta before historic embankment (The Netherlands)*. *Geology* **46**: 1–4. <https://doi.org/10.1130/G45188.1>



- Polak, M.**, 2014. An early Roman naval base at Vechten (prov. Utrecht/NL): Facts and fiction. In: Nickel, C., Röder, M. & Scholz, M. (eds): *Honesta Missione*, Festschrift für Barbara Pferdehirt. (Monographien des Römisch-Germanischen Zentralmuseums, band 100). (Mainz): 69–98.
- Polak, M. & Wynia, S.L.**, 1991. The Roman forts at Vechten, a survey of the excavations 1829–1989. *Oudheidkundige Mededelingen uit het Rijksmuseum van Oudheden* 71: 125–156.
- Polak, M. & Kooistra, L.I.**, 2013. A sustainable frontier? The establishment of the Roman frontier in the Rhine delta. Part 1: From the end of the Iron Age to the death of Tiberius (ca. 50 BC – AD 37). *Jahrbuch des Römisch-Germanischen Zentralmuseums* 60 (Mainz): 355–458.
- Polak, M., Bödecker, S., Berger, L., Zandstra, M.J.M. & Leene, T.**, 2019. Frontiers of the Roman Empire - The Lower German Limes. Nomination file for inscription on the UNESCO World Heritage List. Part I – Nomination file. Bonn.
- Rathmann, M.**, 2004. Die Reichsstrassen der Germania Inferior. *Bonner Jahrbücher* 204: 1–45.
- Rau, R.**, 1968. Briefe des Bonifatius, Willibalds Leben des Bonifatius nebst einigen zeitgenössischen Dokumenten. Ausgewählte Quellen zur deutschen Geschichte des Mittelalters, Vol. IVb. Wissenschaftliche Buchgesellschaft (Darmstadt).
- Ruffing, K.**, 2018. Wirtschaft und Handel in der römischen Kaiserzeit. In: C. Eger (ed.): *Warenwege – Warenflüsse. Handel, Logistik und Transport am Römischen Niederrhein*. (Xantener Berichte 32). Nünnerich-Asmus Verlag (Oppenheim am Rhein): 3–23.
- Salomon, F., Purdue, L., Goiran, J.P. & Berger, J.F.**, 2014. Introduction to the special issue: Roman canals studies – main research aims. *Water History* 6: 1–9. <https://doi.org/10.1007/s12685-014-0101-y>
- Schönfeld, J.F.**, 1940a. Waar lag – ten tijde van de Romeinen – het splitsingspunt van den Rijn? *De Ingenieur* 55: 223–230.
- Schönfeld, J.F.**, 1940b. Is het ten tijde der Romeinen bestaande splitsingspunt van den Rijn gevonden? *Tijdschrift van het Koninklijk Aardrijkskundig Genootschap*, tweede serie 57: 549–570.
- Spek, T., Zeiler, F.D. & W. Raap.** 1996. Van de Hunnepe tot de Zee, de geschiedenis van het Waterschap Salland. *IJsselakademie* (Kampen).
- Stouthamer, E., Berendsen, H.J.A., Peeters, J. & Bouman, M.T.I.J.**, 2008. Toelichting Bodemkaart Veengebieden provincie Utrecht, schaal 1:25.000. Utrecht.
- Teunissen, D.**, 1980. Enkele gebeurtenissen uit het leven van de Romeinse veldheer Drusus in het licht van enige nieuwe geologische gegevens uit oostelijk Gelderland. *Westerheem* 29: 321–334.
- Törnqvist, T.E.**, 1993. Fluvial sedimentary geology and chronology of the Holocene Rhine-Meuse delta, the Netherlands. *Netherlands Geographical Studies* 166, PhD Thesis Utrecht University (Utrecht).
- Uggeri, G.**, 1997. I canali navigabili dell'antico delta Padano. In: Quilici Gigli, S. (ed.): *Uomo, acqua e paesaggio – Atti dell'Incontro sul tema 'Irregimentazione delle acque e trasformazione del paesaggio antico'*, S. maria Capua Vetere, 22–23 novembre 1996. (Roma): 55–60.
- van Asch van Wijck, M.M.A.J.**, 1846. *Geschiedkundige beschouwing van het oude handelsverkeer der stad Utrecht, van de vroegste tijden af tot aan de veertiende eeuw* (vol. 2). Utrecht.
- van de Meene, E.A.**, 1979. Het ontstaan van de Gelderse IJssel. *Koninklijk Nederlands Aardrijkskundig Genootschap – Geografisch Tijdschrift*, *Nieuwe Reeks* 13: 202–210.
- van de Meene, E.A., van Meerkerk, M. & van der Staay, J.**, 1988. Toelichting bij de geologische kaart van Nederland 1:50.000, Blad Utrecht Oost (31-O). *Rijks Geologische Dienst* (Haarlem).
- van de Ven, G.**, 2007. Verdeel en beheers! 300 jaar Pannerdensch Kanaal. *Veen Magazines* (Diemen).
- van der Tuuk, L. & Cruysheer, A.** 2014. De Utrechtse Vecht, levensader in de vroege middeleeuwen. *Jaarboekje 2013 van het oudheidkundig genootschap Niftarlake*, 102–151.
- van der Velde, H.M., Bloo, S.B.C., van Beurden, L., Brinkhuizen, D.C., van Dijk, J., van Dinter, M., Esser, E., Schreijer, E. & Spanjer, M.**, 2003. Archeologisch onderzoek in het kader van de verbreding van de A2. Aanvullend Archeologisch Onderzoek naar een vindplaats uit de ijzertijd en de Hollandse waterlinie. ADC-rapport 167 (Bunschoten). <https://doi.org/10.17026/dans-292-k8za>
- van Dinter, M.**, 2013. The Roman Limes in the Netherlands: how a delta landscape determined the location of the military structures. *Netherlands Journal of Geosciences - Geologie en Mijnbouw* 92 (1), 11–32.
- van Dinter, M.**, 2017. Living along the Limes. Landscape and settlement in the Lower Rhine Delta during Roman and Early Medieval times. *Utrecht Studies in Earth Sciences*, vol. 135, dissertation University of Utrecht (Utrecht).
- van Dinter, M., Cohen, K.M., Hoek, W.Z., Stouthamer, E., Jansma, E. & Middelkoop, H.**, 2017. Late Holocene lowland fluvial archives and geoarchaeology: Utrecht's case study of Rhine river abandonment under Roman and Medieval settlement. 20-yr FLAG special issue on Fluvial Archives, *Quaternary Science Reviews* 166: 1–39. <https://doi.org/10.1016/j.quascirev.2016.12.003>
- van Lanen, R.J.**, 2017. Changing ways. Patterns of connectivity, habitation and persistence in Northwest European lowlands during the first millennium AD. Dissertation University of Utrecht (Utrecht Studies in Earth Sciences 137).
- van Lanen, R.J. & Pierik, H.J.**, 2017. Calculating connectivity patterns in delta landscapes: modelling Roman and early medieval route networks and their stability in dynamic lowlands, *Quaternary International* 501: 393–412. <http://doi.org/10.1016/j.quaint.2017.03.009>
- van Loon, A.H.**, 2010. Unravelling hydrological mechanisms behind fen deterioration in order to design restoration strategies. De hydrologische oorzaken van de achteruitgang van laagveengebieden. PhD thesis, University of Utrecht (Utrecht).
- van Popta, Y., K.M. Cohen, P.C. Vos & Spek, T.**, 2020. Reconstructing eroded medieval landscapes of the Noordoostpolder (Zuyder Zee area, The Netherlands): An interdisciplinary palaeogeographical take on the historic landscape development between AD 1100 and 1400. *Landscape History*, 41(2): 27–56. <https://doi.org/10.1080/01433768.2020.1835180>
- Verhagen, J.G.M., Kluiving, S.J., Anker, E., van Leeuwen, L. & Prins, M.A.**, 2017. Geoarchaeological prospection for Roman waterworks near the late Holocene Rhine-Waal delta bifurcation, the Netherlands. *Catena* 149: 460–473. <https://doi.org/10.1016/j.catena.2016.03.027>
- Vollgraff, C.W.**, 1938. De dijk van Drusus. *Mededeelingen der koninklijke Nederlandsche akademie van wetenschappen afd. letterkunde, Nieuwe Reeks* 1: 555–577.
- Vollgraff, C.W.**, 1939. De moles van Drusus. *Mededeelingen der koninklijke Nederlandsche akademie van wetenschappen afd. letterkunde, Nieuwe Reeks* 2: 141–143.
- Vos, P.C.**, 2008. The geological development of the Oer-IJ area. In Kok, M.S.M. (ed.): *The homcoming of religious practice: an analysis of offering sites in the wet low-lying parts of the landscape in the Oer-IJ area (2500 BC–AD 450)*. University of Amsterdam (Amsterdam): 81–94.
- Vos, P.C.**, 2015. Origin of the Dutch coastal landscape. Long-term landscape evolution of the Netherlands during the Holocene, described and visualized in national, regional and local palaeogeographical map series. Dissertation university of Utrecht. *Barkhuis* (Groningen). <https://doi.org/10.2307/j.ctt2204s8d>
- Vos, P. & de Vries, S.**, 2013. Tweede generatie palaeogeografische kaarten van Nederland (versie 2.0). Utrecht: Deltares (on 7 Oct 2018 downloaded from [www.archeologiein nederland.nl](http://www.archeologiein nederland.nl)).
- Vos, P.C., de Koning, J. & van Eerden, R.**, 2015. Landscape history of the Oer-IJ tidal system, Noord-Holland (the Netherlands). *Netherlands Journal of Geosciences* 94: 295–332. <https://doi.org/10.1017/njg.2015.27>
- Weerts, H.J.T., Cleveringa, P. & Gouw, M.J.P.**, 2002. De Vecht/Angstel, een riviersysteem in het veen. *Grondboor en Hamer* 56(3/4): 66–71.
- Willems, W.J.H.**, 1981/1984. Romans and Batavians, a regional study in the Dutch eastern river area. *Berichten van de Rijksdienst voor het Oudheidkundig Bodemonderzoek* 31: 7–201.
- Willems, W.J.H.**, 1981/1984. Romans and Batavians, a regional study in the Dutch eastern river area. *Berichten van de Rijksdienst voor het Oudheidkundig Bodemonderzoek* 34: 42–331 (202–491). Dissertation University of Amsterdam 1986 (Amsterdam). <https://doi.org/10.17026/dans-zky-yx4w> and <https://doi.org/10.17026/dans-z4j-53sp>
- Zandstra, M.J.M. & Polak, M.**, 2012. De Romeinse versterkingen in Vechten-Fectio. Het archeologisch onderzoek in 1946–1947. *Auxiliaria* 11 (Nijmegen).