

COMPLEX-HARMONIC MEIER'S THEOREM

SHINJI YAMASHITA

1. Fatou's theorem is true for a bounded complex-valued harmonic function in the disk $D: |z| < 1$. One asks naturally: "Is Meier's topological analogue of Fatou's theorem (simply, "*MF* theorem"; [14, p. 330, Theorem 6], cf. [10, p. 154, Theorem 8.9]) true for a bounded complex-valued harmonic function in D ?" We shall give the affirmative answer to this question. Furthermore, the horocyclic *MF* theorem [2, p. 14, Theorem 5] in the complex-harmonic form will be proved in parallel.

For recent various discussions on Plessner's and Meier's theorems we consult [1~7, 11, 12, 15~18].

2. In the rest of this note we denote by $\delta(\zeta_0, \rho)$ the open disk $|z - \zeta_0| < \rho$ in the z -plane.

LEMMA 1. *Let a function $g(\zeta)$ be complex-valued and harmonic (simply, "complex-harmonic") in $\delta(\zeta_0, \rho)$ and $|g(\zeta)| < 1$ for $\zeta \in \delta(\zeta_0, \rho)$. Then we have*

$$(1) \quad |g(\zeta) - g(\zeta_0)| \leq (8/\pi) \arctan(|\zeta - \zeta_0|/\rho)$$

for $\zeta \in \delta(\zeta_0, \rho)$ (*Schwarz's lemma*).

Proof. Let $w = (\zeta - \zeta_0)/\rho$ and consider the function

$$G(w) = \{g(\rho w + \zeta_0) - g(\zeta_0)\}/2$$

in $D: |w| < 1$. Then $G(0) = 0$ and $|G(w)| < 1$ in D , so that we may apply the ready Schwarz lemma [13, p. 101, Lemma] to the complex-harmonic G in D . The inequality [13, p. 101, (3)]

$$|G(w)| \leq (4/\pi) \arctan |w|$$

for $w \in D$ proves (1).

Q.E.D.

The reader should know the definition of cluster set, chordal cluster set and angular cluster set [10, pp. 1, 72 and 73].

Received July 10, 1970.

LEMMA 2. *Let a function $f(z)$ be complex-harmonic in D with $|f(z)| < 1$ for $z \in D$. Assume that*

$$(2) \quad C_X(f, 1) \neq C_D(f, 1),$$

where X is a chord of the unit circle passing through the point $z = 1$. Then there exists an angle Δ at $z = 1$ (i.e., the interior of a triangle lying in D except for one vertex $z = 1$) such that

$$(3) \quad C_\Delta(f, 1) \neq C_D(f, 1).$$

Proof. Choose a point $P \in C_D(f, 1) - C_X(f, 1)$ and let

$$0 < 2\varepsilon < \text{dis} \{P, C_X(f, 1)\}.$$

By (2) such a point P does exist and further we can find a rectilinear segment $X_1 \subset X$ terminating at $z = 1$ such that

$$(4) \quad \overline{f(X_1)} \cap \delta(P, \varepsilon) = \phi \text{ (empty)}$$

by the very definition of $C_X(f, 1)$. Let φ be the directed angle, $|\varphi| < \pi/2$, made by X and the radius of D at $z = 1$ and suppose without loss of generality that $0 \leq \varphi < \pi/2$. Set

$$r(z_o) = |1 - z_o| \sin(\pi/4 - \varphi/2), \quad z_o \in X_1$$

and choose a constant μ such that

$$(5) \quad 0 < \mu < \tan(\pi\varepsilon/16).$$

Then $\tan(\pi\varepsilon/16) < 1 < \pi/2$ because of $\varepsilon < 1$ and for any point $z \in \delta(z_o, \mu r(z_o))$ ($z_o \in X_1$) we have

$$(6) \quad |f(z) - f(z_o)| \leq (8/\pi) \arctan \{ \mu r(z_o) / r(z_o) \} < \varepsilon/2$$

by (1) of Lemma 1 and (5) if z_o is so near to $z = 1$ that $\delta(z_o, r(z_o)) \subset D$. Now, as $X_1 \ni z_o \rightarrow 1$, the disks $\delta(z_o, \mu r(z_o))$ sweep an angle Δ at $z = 1$, so that by (4) and (6) we have

$$(7) \quad \overline{f(\Delta)} \cap \delta(P, \varepsilon/4) = \phi.$$

Now that (7) means $P \notin \overline{f(\Delta)}$ we have

$$P \in C_D(f, 1) - C_\Delta(f, 1),$$

which proves (3). Q.E.D.

For the terminology, “right horocycle”, “right horocyclic cluster set”,

“right horocyclic angle”, etc. we refer to [2, pp. 4–6].

LEMMA 3. *Let a function $f(z)$ be complex-harmonic in D with $|f(z)| < 1$ for $z \in D$. Assume that*

$$(8) \quad C_{h(1)}(f, 1) \neq C_D(f, 1),$$

where $h(1) = h_r^+(1)$ is a right horocycle at $z = 1$. Then there exists a right horocyclic angle $H(1) = H_{r_1, r_2, r_3}^+(1)$ at $z = 1$ such that

$$(9) \quad C_{H(1)}(f, 1) \neq C_D(f, 1).$$

Proof. We use a different method from Bagemihl’s [2, p. 14, Lemma 3]. By (8) we can find a point $P \in C_D(f, 1) - C_{h(1)}(f, 1)$ and we then set

$$0 < 2\varepsilon < \text{dis} \{P, C_{h(1)}(f, 1)\}.$$

By the definition of $C_{h(1)}(f, 1)$ we obtain a subarc α of $h(1)$ terminating at $z = 1$ such that

$$(10) \quad \overline{f(\alpha)} \cap \delta(P, \varepsilon) = \phi.$$

We consider next the map

$$z = \chi(\zeta) = (\zeta - 1)/(\zeta + 1)$$

from the half plane $\text{Re} \zeta > 0$ onto D . The initial point of $h(1)$ lies on the real axis, which we denote by x , $|x| < 1$. Then the image L_x of $h(1)$ by the map χ^{-1} is the half line

$$L_x = \{\zeta; \text{Re} \zeta = (1 + x)/(1 - x) \text{ and } \text{Im} \zeta \leq 0\}.$$

Let β be the image of α by χ^{-1} and let

$$(11) \quad 0 < \mu < \tan(\pi\varepsilon/16).$$

Let $0 < \rho < (1 + x)/(1 - x)$ and consider the composed function $F(\zeta) = f \circ \chi(\zeta)$ in the disk $\delta(\zeta_o, \rho)$, where $\zeta_o \in \beta$. Then for $\zeta \in \delta(\zeta_o, \mu\rho) \subset \delta(\zeta_o, \rho)$, we have

$$(12) \quad |F(\zeta) - F(\zeta_o)| \leq (8/\pi) \text{arc tan}(\mu\rho/\rho) < \varepsilon/2$$

by (1) of Lemma 1 combined with (11). Now, as $\beta \ni \zeta_o \rightarrow \infty$ (i.e., $\alpha \ni \chi(\zeta_o) \rightarrow 1$) the disks $\delta(\zeta_o, \mu\rho)$ sweep a strip of width $2\mu\rho$ whose image by χ contains a right horocyclic angle $H(1) = H_{r_1, r_2, r_3}^+(1)$ at $z = 1$. By (10) and (12) we have

$$\overline{f(H(1))} \cap \delta(P, \varepsilon/4) = \phi,$$

so that we have (9).

Q.E.D.

Remark. Lemma 3 is true if the word “right” is replaced by “left” where it is.

3. A point $e^{i\theta}$ of the circle is a Meier point (horocyclic Meier point, resp.) of a complex-harmonic function $f(z)$ in D if $C_D(f, e^{i\theta})$ is a proper subset of the Riemann sphere and if every chordal cluster set (every right or left horocyclic cluster set, resp.) of f at $e^{i\theta}$ coincides with $C_D(f, e^{i\theta})$ [10, p. 153], [2, p. 6].

By means of Lemmas 2 and 3, and Collingwood’s maximality theorem ([8, p. 1241, Theorem 4], [9, p. 8, Theorem 4]; [10, p. 80, Theorem 4.10]) or its ready generalization from (Stolz) angles to horocyclic angles we have the following two theorems.

THEOREM 1. *Let a function $f(z)$ be bounded, complex-valued and harmonic in the disk $|z| < 1$. Then all points of the circle $\Gamma: |z| = 1$ are, except perhaps for a set of first Baire category on Γ [10, p. 75], Meier points of f .*

THEOREM 2. *Let a function $f(z)$ be bounded, complex-valued and harmonic in the disk $|z| < 1$. Then all points of the circle $\Gamma: |z| = 1$ are, except perhaps for a set of first Baire category on Γ , horocyclic Meier points of f .*

4. As a concluding remark we note that further generalizations of Theorems 1 and 2 are possible (cf. [15]). Let Ω and Ω' be domains in the z -plane and in the ζ -plane respectively. A complex-valued function $f(z)$ in Ω is called K -quasi-conformal harmonic (simply, “ $KQCH$ ”) in Ω provided that $f(z)$ is of the composed form $f(z) = g \circ Q(z)$, where $\zeta = Q(z)$ is a K -quasi-conformal homeomorphism ($K \geq 1$) from Ω onto Ω' and $g(\zeta)$ is complex-harmonic in Ω'^{**} . The key lemma for the proof of MF or horocyclic MF theorem of $KQCH$ functions in D is, of course, an analogue of the Schwarz lemma:

LEMMA 1^{b18}. *Let a function $f(z)$ be $KQCH$ and $|f(z)| < 1$ in the disk $\delta(z_0, q)$. Then for $z \in \delta(z_0, q)$ we have*

$$(13) \quad |f(z) - f(z_0)| \leq (8/\pi) \arctan(4q^{-1/K} |z - z_0|^{1/K}).$$

Proof. We may consider $f = g \circ T$, where T is a K -quasi-conformal self-homeomorphism of $\delta(z_0, q)$ with the additional property that $z_0 = T(z_0)$, and g is complex-harmonic in $\delta(z_0, q)$. Furthermore, we know about T that [15, p. 323, line 2 from below]

^{*}) A domain Ω' may depend upon f .

$$|T(z) - T(z_0)| \leq 4q^{1-(1/K)}|z - z_0|^{1/K}, \quad z \in \delta(z_0, q),$$

an inequality due to A. Mori, so that combining this with Lemma 1 of section 2 we obtain (13).

REFERENCES

- [1] F. Bagemihl, Meier points of holomorphic functions, *Math. Ann.* **155** (1964), 422–424.
- [2] F. Bagemihl, Horocyclic boundary properties of meromorphic functions, *Ann. Acad. Sci. Fenn. Ser. A-I. Math.* No. **385** (1966), 18 pp.
- [3] F. Bagemihl, Chordal limits of holomorphic functions at Plessner points, *J. Sci. Hiroshima Univ. Ser. A-I.* **30** (1966), 109–115.
- [4] F. Bagemihl, On the sharpness of Meier's analogue of Fatou's theorem, *Israel J. Math.* **4** (1966), 230–232.
- [5] F. Bagemihl, Generalizations of two theorems of Meier concerning boundary behavior of meromorphic functions, *Publ. Math. Debrecen*, **14** (1967), 53–55.
- [6] F. Bagemihl, An improvement on Plessner's theorem, *Math. Z.* **110** (1969), 177–179.
- [7] F. Bagemihl, Meier points and horocyclic Meier points of continuous functions, *Ann. Acad. Sci. Fenn. Ser. A-I. Math.* No. **461** (1970), 7 pp.
- [8] E.F. Collingwood, Cluster sets of arbitrary functions, *Proc. Nat. Acad. Sci. U.S.A.* **46** (1960), 1236–1242.
- [9] E.F. Collingwood, Cluster set theorems for arbitrary functions with applications to function theory, *Ann. Acad. Sci. Fenn. Ser. A-I. Math.* No. **336/8** (1963), 15 pp.
- [10] E.F. Collingwood and A.J. Lohwater, *The theory of cluster sets*, Cambridge (1966).
- [11] S. Dragosh, Horocyclic boundary behavior of meromorphic functions, *J. Analyse Math.* **22** (1969), 37–48.
- [12] S. Dragosh, Horocyclic cluster sets of functions defined in the unit disc, *Nagoya Math. J.* **35** (1969), 53–82.
- [13] E. Heinz, On one-to-one harmonic mappings, *Pacific J. Math.* **9** (1959), 101–105.
- [14] K.E. Meier, Ueber die Randwerte der meromorphen Funktionen, *Math. Ann.* **142** (1961), 328–344.
- [15] S. Yamashita, Quasi-conformal extension of Meier's theorem, *Proc. Japan Acad.* **46** (1970), 323–325.
- [16] S. Yamashita, Cluster sets of algebroid functions, *Tôhoku Math. J.* **22** (1970), 273–289.
- [17] S. Yamashita, Boundary behaviour of functions harmonic in the unit ball, *Proc. Japan Acad.* **46** (1970), No. **6**, 490–493.
- [18] S. Yamashita, On Fatou- and Plessner-type theorems, *Proc. Japan Acad.* **46** (1970), No. **6**, 494–495.

*Mathematical Institute
Tôhoku University
Sendai 980, Japan*