



The Equivariant Cohomology Rings of Peterson Varieties in All Lie Types

Megumi Harada, Tatsuya Horiguchi, and Mikiya Masuda

Abstract. Let G be a complex semisimple linear algebraic group and let Pet be the associated *Peterson variety* in the flag variety G/B . The main theorem of this note gives an efficient presentation of the equivariant cohomology ring $H_S^*(\text{Pet})$ of the Peterson variety as a quotient of a polynomial ring by an ideal J generated by quadratic polynomials, in the spirit of the Borel presentation of the cohomology of the flag variety. Here the group $S \cong \mathbb{C}^*$ is a certain subgroup of a maximal torus T of G . Our description of the ideal J uses the Cartan matrix and is uniform across Lie types. In our arguments we use the Monk formula and Giambelli formula for the equivariant cohomology rings of Peterson varieties for all Lie types, as obtained in the work of Drellich. Our result generalizes a previous theorem of Fukukawa–Harada–Masuda, which was only for Lie type A .

1 Introduction

The main result of this paper is an explicit and efficient presentation of the equivariant cohomology ring of Peterson varieties in all Lie types in terms of generators and relations, in the spirit of the well-known Borel presentation of the cohomology of the flag variety. We briefly recall the setting of our results. The *Peterson variety* has been much studied due to its relation, for example, to the quantum cohomology of the flag variety [11, 13]. Thus it is natural to study their topology, *e.g.*, the structure of their (equivariant) cohomology rings. In Lie type A the Peterson variety Pet can be easily described in a concrete manner. Specifically, it is defined to be the following subvariety of the full flag variety $\text{Flags}(\mathbb{C}^n)$:

$$\text{Pet} := \{V_\bullet \mid NV_i \subseteq V_{i+1} \text{ for all } i = 1, \dots, n-1\},$$

where V_\bullet denotes a nested sequence $0 \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq V_{n-1} \subseteq V_n = \mathbb{C}^n$ of subspaces of \mathbb{C}^n and $\dim_{\mathbb{C}} V_i = i$ for all i and $N: \mathbb{C}^n \rightarrow \mathbb{C}^n$ denotes a principal nilpotent operator. For a general complex semisimple linear algebraic group G , there is also a concrete Lie-theoretic description of the Peterson variety as a subvariety of its flag variety G/B , which we recall in Section 2 below.

In addition, there is a natural subgroup $S \cong \mathbb{C}^*$ (which is a subgroup of the maximal torus T of G) which acts on Pet (for details see Section 2). The inclusion of Pet into

Received by the editors June 5, 2014.

Published electronically December 9, 2014.

AMS subject classification: 55N91, 14N15.

Keywords: equivariant cohomology, Peterson varieties, flag varieties, Monk's formula, Giambelli's formula.

G/B induces a natural ring homomorphism

$$H_T^*(G/B) \rightarrow H_S^*(\text{Pet}).$$

The purpose of this manuscript is to give an efficient presentation of the equivariant cohomology ring $H_S^*(\text{Pet})$ in all Lie types, using the recent work of Drellich [4] which gives a Monk formula and a Giambelli formula for Peterson varieties in all Lie types. In particular, we are able to obtain a uniform description of the relevant ideal J , valid for all Lie types, using the Cartan matrix associated to the Lie algebra \mathfrak{g} of G . In particular, our analysis shows that the ideal J is generated by quadratics.

Our proof uses Hilbert series and regular sequences, in a similar spirit to previous work of Fukukawa and the first and third authors [6], which was in turn motivated by the work of [5, 7], which computes the graph cohomology of the GKM graphs of the flag varieties of classical type and of G_2 .

Brion and Carrell give a different description of the equivariant cohomology rings of Peterson (and other nilpotent Hessenberg) varieties as the coordinate rings of certain affine curves [3, Theorem 3] using techniques from algebraic geometry. In a related direction, by extending the ideas of the present manuscript, we give in [1] a uniform description (with explicit generators and relations) of the equivariant cohomology rings of arbitrary regular nilpotent Hessenberg varieties—a class of varieties that includes the Peterson variety and the full flag variety—in Lie type A .

This paper is organized as follows. We briefly recall the necessary background in Section 2. We derive the relevant quadratic relations in Section 3. In particular, a key computation is contained in Lemma 3.3. The main theorem is proven in Section 4.

2 Background on the Peterson Variety

In this section we record some facts about Peterson varieties that we require in this manuscript.

Let G be a complex semisimple linear algebraic group of rank n . We fix B a Borel subgroup and T a maximal torus of G such that $T \subseteq B \subseteq G$. These choices then determine the following data:

- a set of simple roots $\Delta = \{\alpha_1, \dots, \alpha_n\}$,
- the associated Weyl group W ,
- the associated Lie algebras $\mathfrak{t} \subseteq \mathfrak{b} \subseteq \mathfrak{g}$,
- root spaces $\mathfrak{g}_\alpha \subseteq \mathfrak{g}$ for each root α .

Definition Let E_α be a basis element of the root space \mathfrak{g}_α and let $N_0 = \sum_{\alpha \in \Delta} E_\alpha$, a regular nilpotent operator. In this setting we may define the *Peterson variety (associated with \mathfrak{g})* as

$$\text{Pet} := \left\{ gB \in G/B \mid \text{Ad}(g^{-1})(N_0) \in \mathfrak{b} \oplus \bigoplus_{\alpha \in -\Delta} \mathfrak{g}_\alpha \right\}.$$

As is well known, the maximal torus T acts on G/B by left multiplication. This action does not in general preserve the Peterson variety. However, by using the homomorphism $\phi: T \rightarrow (\mathbb{C}^*)^n$ defined by $t \mapsto (\alpha_1(t), \dots, \alpha_n(t))$ and defining S to be

the connected component of the identity in

$$\phi^{-1}(\{(c, c, \dots, c) \mid c \in \mathbb{C}^*\}),$$

it can be seen that the restriction of the T -action on G/B to the subgroup S does preserve Pet [8, Lemma 5.1].

Next recall that the T -fixed points of G/B are in bijective correspondence with the Weyl group W of G . Moreover, since the S -fixed points Pet^S of the Peterson variety satisfy the relation

$$\text{Pet}^S = \text{Pet} \cap (G/B)^T,$$

we may view Pet^S as a subset of the Weyl group W . Indeed, the fixed point set Pet^S can be described concretely as follows. For a subset K of the set Δ simple roots, let W_K denote the parabolic subgroup generated by K and let w_K denote the longest element of W_K . Then it is known [8, Proposition 5.8] that

$$\text{Pet}^S = \{w_K \mid K \subseteq \Delta\}.$$

Here and below we always use complex coefficients \mathbb{C} for our cohomology rings and hence omit it from our notation. Let $\alpha_i: T \rightarrow \mathbb{C}^*$ be a homomorphism which thus determines a complex 1-dimensional representation of T . Let $ET \times_T \mathbb{C} \rightarrow BT$ be the corresponding complex line bundle, and by slight abuse of notation we let $\alpha_i \in H^2(BT)$ also denote the corresponding first Chern class. With this notation in place we have

$$H^*(BT) = \mathbb{C}[\alpha_1, \dots, \alpha_n].$$

Consider the 1-dimensional representation of the diagonal subgroup

$$\{(c, c, \dots, c) : c \in \mathbb{C}^*\} \subseteq (\mathbb{C}^*)^n$$

obtained via the projection $(c, c, \dots, c) \rightarrow c$. Composing with the restriction to S of the above homomorphism ϕ , we obtain a 1-dimensional representation of S and an associated line bundle $ES \times_S \mathbb{C} \rightarrow BS$ with first Chern class denoted $t \in H^2(BS)$. With this notation in place we have

$$H^*(BS) = \mathbb{C}[t].$$

Next we recall that the inclusion homomorphism $S \hookrightarrow T$ induces a homomorphism $\pi: H^*(BT) \rightarrow H^*(BS)$, and from the definition of ϕ we obtain

$$(2.1) \quad \pi(\alpha_i) = t \quad (i = 1, 2, \dots, n).$$

We now consider the commutative diagram

$$(2.2) \quad \begin{array}{ccc} H_T^*(G/B) & \longrightarrow & \bigoplus_{w \in (G/B)^T = W} H_T^*(w) \\ \rho \downarrow & & \downarrow \pi \\ H_S^*(\text{Pet}) & \longrightarrow & \bigoplus_{w \in \text{Pet}^S \subseteq W} H_S^*(w), \end{array}$$

where all the maps are induced from inclusions of subgroups or inclusions of subspaces. As is well known, the odd cohomology $H^{\text{odd}}(G/B)$ of G/B vanishes. The same holds for the Peterson variety, *i.e.*, $H^{\text{odd}}(\text{Pet}) = 0$ [12]. Thus we obtain that

both horizontal maps in (2.2) are injective, and we may identify $H_T^*(G/B)$ (respectively $H_S^*(\text{Pet})$) with its image under these maps. For $w \in (G/B)^T \cong W$ (respectively $w \in \text{Pet}^S \subseteq W$) and $f \in H_T^*(G/B)$ (respectively $f \in H_S^*(\text{Pet})$) we will denote by $f(w)$ the restriction of f to the w -th factor $H_T^*(w) = H^*(BT) = \mathbb{C}[\alpha_1, \dots, \alpha_n]$ (resp. $H_S^*(w) = H^*(BS) = \mathbb{C}[t]$) in the direct products on the right hand sides of (2.2).

For $v \in W$, we let σ_v denote the corresponding equivariant Schubert class in $H_T^*(G/B)$, and let p_v denote its image $\rho(\sigma_v)$ in $H_S^*(\text{Pet})$. We call p_v a *Peterson Schubert class* (associated to v). Let s_i be the simple reflection corresponding to a simple root α_i . The vertices of the Dynkin diagram corresponding to the set of simple roots $\Delta = \{\alpha_1, \dots, \alpha_n\}$ is in 1-1 correspondence with Δ . Here and below, we assume to be fixed an ordering of the simple roots as given in [4, Figure 1] (which in turn agrees with the standard ordering in [9, p. 58]). With respect to this ordering, given any subset $K = \{\alpha_{a_1}, \alpha_{a_2}, \dots, \alpha_{a_k}\}$ of the simple roots with $a_1 < a_2 < \dots < a_k$, we define an element v_K of W by the formula

$$v_K := s_{a_1}s_{a_2}\cdots s_{a_k}.$$

The Peterson Schubert classes p_{v_K} corresponding to the Weyl group elements v_K defined above satisfy the following property.

Proposition 2.1 (Theorem 3.5 in [4]) *The Peterson Schubert classes $\{p_{v_K} \mid K \subseteq \Delta\}$ form a $\mathbb{C}[t]$ -module basis for $H_S^*(\text{Pet})$.*

It follows from Proposition 2.1 that the ρ in (2.2) is surjective. It is well known that the equivariant Schubert classes σ_{s_i} generate $H_T^*(G/B)$ as a $\mathbb{C}[\alpha_1, \dots, \alpha_n]$ -algebra. From the surjectivity of the homomorphism ρ we immediately obtain the following.

Proposition 2.2 *The Peterson Schubert classes p_{s_i} ($i = 1, 2, \dots, n$) generate $H_S^*(\text{Pet})$ as a $\mathbb{C}[t]$ -algebra.*

Since the odd cohomology $H^{\text{odd}}(\text{Pet})$ of the Peterson variety vanishes, we know that as a $\mathbb{C}[t]$ -module the equivariant cohomology $H_S^*(\text{Pet})$ is isomorphic to $\mathbb{C}[t] \otimes H^*(\text{Pet})$. It is known [3, Theorem 3] that

$$(2.3) \quad \begin{aligned} F(H^*(\text{Pet}), s) &= (1 + s^2)^n, \\ F(H_S^*(\text{Pet}), s) &= \frac{(1 + s^2)^n}{1 - s^2}, \end{aligned}$$

where the left hand sides denotes the Hilbert series of the graded rings $H^*(\text{Pet})$ and $H_S^*(\text{Pet})$ with respect to the variable s (of degree 1).

Fix an integer i with $1 \leq i \leq n$ and a subset $K \subseteq \Delta$. From Proposition 2.1 it follows that the product $p_{s_i} \cdot p_{v_K}$ can be written uniquely as a $\mathbb{C}[t]$ -linear combination of the p_{v_J} (for $J \subseteq \Delta$). Monk's formula gives a concrete computation of the coefficients in this linear combination.

Theorem 2.3 (Monk’s formula for Peterson varieties for all Lie types [4, Theorem 4.2]) *The Peterson Schubert classes satisfy the following relation:*

$$p_{s_i} \cdot p_{v_K} = p_{s_i}(w_K) \cdot p_{v_K} + \sum_{\substack{J \supset K \\ |J|=|K|+1}} c_{i,K}^J \cdot p_{v_J}$$

where the coefficients $c_{i,K}^J$ are nonnegative rational numbers. More specifically, we have

$$c_{i,K}^J = (p_{s_i}(w_J) - p_{s_i}(w_K)) \cdot \frac{p_{v_K}(w_J)}{p_{v_J}(w_J)}.$$

Next we recall Giambelli’s formula. From Proposition 2.2 it follows that each module generator p_{v_K} can be expressed as a polynomial (with $\mathbb{C}[t]$ coefficients) in the (ring) generators p_{s_i} . Giambelli’s formula gives a concrete expression for this polynomial as follows.

Theorem 2.4 (Giambelli’s formula for Peterson varieties for all Lie types [4, Theorem 5.5]) *Suppose K is a subset of the simple roots Δ . Assume that the Dynkin diagram corresponding to the subset K is connected. Then*

$$\frac{|K|!}{|\mathcal{R}(v_K)|} \cdot p_{v_K} = \prod_{\alpha_i \in K} p_{s_i},$$

where $|\mathcal{R}(v_K)|$ denotes the number of distinct reduced-word expressions for v_K .

Remark (cf. Theorem 5.3 in [4]) The connectedness requirement in the above theorem is not serious, in the following sense. Suppose $J, K \subseteq \Delta$ are two subsets of Δ such that their corresponding Dynkin diagrams are connected. Suppose, however, that $J \cup K$ has a corresponding Dynkin diagram that is not connected. Then $p_{v_{J \cup K}}$ is simply the product of p_{v_J} and p_{v_K} , i.e.,

$$p_{v_{J \cup K}} = p_{v_J} \cdot p_{v_K}.$$

3 Quadratic Relations Satisfied by the Peterson Schubert Classes p_{s_i}

In this section, we derive certain quadratic relations satisfied by the cohomology-degree-2 Peterson Schubert classes p_{s_i} ($i = 1, 2, \dots, n$) by using Monk’s formula (Theorem 2.3), Giambelli’s formula (Theorem 2.4), and Billey’s formula (recalled below). We will then show in the next section that these relations are sufficient to determine the equivariant cohomology ring $H_S^*(\text{Pet})$ of the Peterson variety.

Theorem 3.1 (Billey’s formula [2, Theorem 4]) *Let $w \in W$ and fix a reduced word decomposition $w = s_{b_1} s_{b_2} \cdots s_{b_m}$ of w . Set $r(i, w) := s_{b_1} s_{b_2} \cdots s_{b_{i-1}}(\alpha_{b_i})$. For an equivariant Schubert class σ_v for $v \in W$ we have the following:*

$$\sigma_v(w) = \sum_{\substack{\text{reduced words} \\ v = s_{b_{j_1}} s_{b_{j_2}} \cdots s_{b_{j_\ell}}}} \prod_{i=1}^{\ell} r(j_i, w).$$

We begin with some elementary computations involving Peterson Schubert classes. First, from Monk's formula (Theorem 2.3) applied to the case $K = \{\alpha_i\}$ and $v_K = s_i$ we obtain

$$(3.1) \quad p_{s_i}^2 = p_{s_i}(s_i) \cdot p_{s_i} + \sum_{j \neq i} c_i^j \cdot p_{v_{\{\alpha_i, \alpha_j\}}}$$

where

$$(3.2) \quad c_i^j = (p_{s_i}(w_{\{\alpha_i, \alpha_j\}}) - p_{s_i}(s_i)) \cdot \frac{p_{s_i}(w_{\{\alpha_i, \alpha_j\}})}{p_{v_{\{\alpha_i, \alpha_j\}}}(w_{\{\alpha_i, \alpha_j\}})}$$

More specifically, since Theorem 3.1 implies that $\sigma_{s_i}(s_i) = \alpha_i$, from (2.1) we conclude that

$$(3.3) \quad p_{s_i}(s_i) = t.$$

We record the following.

Lemma 3.2 In (3.1), if s_i and s_j commute, then $c_i^j = 0$.

Proof Since s_i and s_j commute, we have $w_{\{\alpha_i, \alpha_j\}} = s_i s_j$. Moreover, from Theorem 3.1 we can compute

$$\sigma_{s_i}(w_{\{\alpha_i, \alpha_j\}}) = \sigma_{s_i}(s_i s_j) = \alpha_i.$$

From (2.1) we get $p_{s_i}(w_{\{\alpha_i, \alpha_j\}}) = t$. Then the equations (3.2) and (3.3) imply $c_i^j = 0$, as desired. ■

In the case when s_i and s_j do not commute, the Dynkin diagram corresponding to the subset $K = \{\alpha_i, \alpha_j\}$ is connected, so Giambelli's formula (Theorem 2.4) yields

$$(3.4) \quad p_{v_{\{\alpha_i, \alpha_j\}}} = \frac{1}{2} p_{s_i} p_{s_j}.$$

In this case, the coefficient appearing in (3.1) can be expressed in terms of the Cartan matrix.

Lemma 3.3 In (3.1), if s_i and s_j do not commute, then

$$c_i^j = -\langle \alpha_i, \alpha_j \rangle$$

where $\langle \alpha_i, \alpha_j \rangle$ denotes the Cartan integer.

Proof From (3.2), (3.3), and (3.4), we can compute

$$(3.5) \quad c_i^j = \frac{2(p_{s_i}(w_{\{\alpha_i, \alpha_j\}}) - t)}{p_{s_j}(w_{\{\alpha_i, \alpha_j\}})}$$

so it suffices to compute $p_{s_i}(w_{\{\alpha_i, \alpha_j\}})$ and $p_{s_j}(w_{\{\alpha_i, \alpha_j\}})$. In what follows we use the notation

$$a_{ij} := \langle \alpha_i, \alpha_j \rangle, \quad i \neq j, \quad a := a_{ij} a_{ji}.$$

With this notation in place, note that by definition of the Cartan integers we have that the action of the simple reflections s_j on the simple roots α_j can be expressed as

$$(3.6) \quad s_j(\alpha_i) = \begin{cases} \alpha_i - a_{ij}\alpha_j & (i \neq j), \\ -\alpha_i & (i = j). \end{cases}$$

In order to prove the lemma, we consider each of the possible cases.

(i) In the case when the Dynkin diagram corresponding to $\{\alpha_i, \alpha_j\}$ is of the form $\overset{\circ}{i} \text{---} \overset{\circ}{j}$, the order of $s_i s_j$ is 3, so we have

$$w_{\{\alpha_i, \alpha_j\}} = s_i s_j s_i = s_j s_i s_j.$$

Using Theorem 3.1 and (3.6) in this case, we can compute that

$$\begin{aligned} \sigma_{s_i}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_i}(s_i s_j s_i) = \alpha_i + s_i s_j(\alpha_i) = a\alpha_i - a_{ij}\alpha_j, \\ \sigma_{s_j}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_j}(s_j s_i s_j) = \alpha_j + s_j s_i(\alpha_j) = a\alpha_j - a_{ji}\alpha_i. \end{aligned}$$

Then (2.1) implies

$$p_{s_i}(w_{\{\alpha_i, \alpha_j\}}) = (a - a_{ij})t, \quad p_{s_j}(w_{\{\alpha_i, \alpha_j\}}) = (a - a_{ji})t.$$

Finally (3.5) yields

$$(3.7) \quad c_i^j = \frac{2(a - a_{ij} - 1)}{(a - a_{ji})},$$

and substituting $a = a_{ij}a_{ji}$, $a_{ij} = -1$ we obtain $c_i^j = -a_{ij}$ as desired.

(ii) In the case $\overset{\circ}{i} \text{---} \overset{\circ}{j}$, the order of $s_i s_j$ is 4, so we have

$$w_{\{\alpha_i, \alpha_j\}} = s_i s_j s_i s_j = s_j s_i s_j s_i.$$

Using this together with Theorem 3.1 and (3.6) we have

$$\begin{aligned} \sigma_{s_i}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_i}(s_i s_j s_i s_j) = \alpha_i + s_i s_j(\alpha_i) = a\alpha_i - a_{ij}\alpha_j, \\ \sigma_{s_j}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_j}(s_j s_i s_j s_i) = \alpha_j + s_j s_i(\alpha_j) = a\alpha_j - a_{ji}\alpha_i, \end{aligned}$$

which is the same as case (i) above. Thus (3.7) also holds in this case, and since $a = a_{ij}a_{ji} = 2$ we obtain $c_i^j = -a_{ij}$ as required.

(iii) Finally, in the case $\overset{\circ}{i} \text{---} \overset{\circ}{j}$, the element $s_i s_j$ has order 6 and thus

$$w_{\{\alpha_i, \alpha_j\}} = s_i s_j s_i s_j s_i s_j = s_j s_i s_j s_i s_j s_i.$$

In this case we have $a = 3$, so Theorem 3.1 and (3.6) yield that

$$\begin{aligned} \sigma_{s_i}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_i}(s_i s_j s_i s_j s_i s_j) = \alpha_i + s_i s_j(\alpha_i) + (s_i s_j)^2(\alpha_i) = 4\alpha_i - 2a_{ij}\alpha_j, \\ \sigma_{s_j}(w_{\{\alpha_i, \alpha_j\}}) &= \sigma_{s_j}(s_j s_i s_j s_i s_j s_i) = \alpha_j + s_j s_i(\alpha_j) + (s_j s_i)^2(\alpha_j) = 4\alpha_j - 2a_{ji}\alpha_i. \end{aligned}$$

Then from (2.1) we compute

$$p_{s_i}(w_{\{\alpha_i, \alpha_j\}}) = (4 - 2a_{ij})t, \quad p_{s_j}(w_{\{\alpha_i, \alpha_j\}}) = (4 - 2a_{ji})t.$$

Equation (3.5) then implies

$$c_i^j = \frac{2(3 - 2a_{ij})}{4 - 2a_{ji}},$$

and finally, using that $a_{ij}a_{ji} = 3$ we get that $c_i^j = -a_{ij}$ as desired.

This completes the proof of the lemma. ■

From the above considerations we obtain the following proposition.

Proposition 3.4 *In the equivariant cohomology ring $H_S^*(\text{Pet})$ of the Peterson variety, the following quadratic relations are satisfied:*

$$\sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle p_{s_i} p_{s_j} - 2t p_{s_i} = 0, \quad 1 \leq i \leq n.$$

Proof If s_i and s_j commute, then $\langle \alpha_i, \alpha_j \rangle = 0$, so by Lemma 3.2 the conclusion of Lemma 3.3 holds in this case. From this and (3.4) we see that (3.1) can be expressed as

$$p_{s_i}^2 = t \cdot p_{s_i} - \frac{1}{2} \sum_{j \neq i} \langle \alpha_i, \alpha_j \rangle p_{s_i} p_{s_j}.$$

Since $\langle \alpha_i, \alpha_i \rangle = 2$ for any i , the above equation can be re-written to be of the form given in the statement of the proposition. ■

4 The Main Theorem

Let $(\langle \alpha_i, \alpha_j \rangle)_{1 \leq i, j \leq n}$ be the Cartan matrix associated to a rank n semisimple Lie algebra \mathfrak{g} . Using the coefficients in the Cartan matrix, we define an ideal J in the polynomial ring $\mathbb{C}[x_1, \dots, x_n, t]$ as follows:

$$J := \left(\sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle x_i x_j - 2t x_i \mid 1 \leq i \leq n \right).$$

From Proposition 2.2 and Proposition 3.4 it then follows that the map sending x_i to p_{s_i} defines a surjective $\mathbb{C}[t]$ -algebra homomorphism

$$(4.1) \quad \varphi: \mathbb{C}[x_1, \dots, x_n, t]/J \twoheadrightarrow H_S^*(\text{Pet}).$$

Here $H^*(BS) = \mathbb{C}[t]$ and Pet denotes the Peterson variety associated to the Lie algebra \mathfrak{g} . Since $H^{\text{odd}}(\text{Pet}) = 0$, as a $H^*(BS)$ -module we have

$$H_S^*(\text{Pet}) \cong H^*(BS) \otimes H^*(\text{Pet}).$$

Defining the ideal \check{J} as

$$(4.2) \quad \check{J} = \left(\sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle x_i x_j \mid 1 \leq i \leq n \right),$$

we then also have a surjective ring homomorphism

$$(4.3) \quad \check{\varphi}: \mathbb{C}[x_1, \dots, x_n]/\check{J} \twoheadrightarrow H^*(\text{Pet}).$$

The following is the main theorem of this paper.

Theorem 4.1 *The maps φ and $\check{\varphi}$ of (4.1) and (4.3) are isomorphisms.*

In order to prove Theorem 4.1 we use the theory of regular sequences. For reference, we briefly recall the definition and a key property of regular sequences (cf. [6]).

Definition Let R be a graded commutative algebra over \mathbb{C} and let R_+ denote the positive-degree elements in R . Then a homogeneous sequence $\theta_1, \dots, \theta_r \in R_+$ is a *regular sequence* if θ_k is a nonzero-divisor in the quotient ring $R/(\theta_1, \dots, \theta_{k-1})$ for every $1 \leq k \leq r$. This is equivalent to saying that $\theta_1, \dots, \theta_r$ is algebraically independent over \mathbb{C} and R is a free $\mathbb{C}[\theta_1, \dots, \theta_r]$ -module.

It is a well-known fact (see for instance [14, p. 35]) that a homogeneous sequence $\theta_1, \dots, \theta_r \in R_+$ is a regular sequence if and only if

$$(4.4) \quad F(R/(\theta_1, \dots, \theta_r), s) = F(R, s) \prod_{k=1}^r (1 - s^{\deg \theta_k}),$$

where $F(R/(\theta_1, \dots, \theta_r), s)$ and $F(R, s)$ denote the Hilbert series of the graded rings $R/(\theta_1, \dots, \theta_r)$ and R , respectively.

The following proposition gives a convenient characterization of regular sequences.

Proposition 4.2 ([6, Proposition 5.1]) *A sequence of positive-degree homogeneous elements $\theta_1, \dots, \theta_r$ in the polynomial ring $\mathbb{C}[z_1, \dots, z_r]$ is a regular sequence if and only if the solution set in \mathbb{C}^r of the equations $\theta_1 = 0, \dots, \theta_r = 0$ consists only of the origin $\{0\}$.*

We can now prove our main theorem.

Proof of Theorem 4.1 We first claim that if $\check{\varphi}$ is an isomorphism then it follows that φ is an isomorphism. To see this, suppose that $\check{\varphi}$ is an isomorphism. Then the sequence

$$\begin{aligned} \theta_i &:= \sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle x_i x_j - 2tx_i \quad \text{for } 1 \leq i \leq n, \\ \theta_{n+1} &:= t \end{aligned}$$

in $\mathbb{C}[x_1, \dots, x_n, t]$ is regular, where $\deg(x_i) = \deg(t) = 2$. Indeed,

$$\begin{aligned} &F(\mathbb{C}[x_1, \dots, x_n, t]/(\theta_1, \dots, \theta_n, \theta_{n+1}), s) \\ &= F(\mathbb{C}[x_1, \dots, x_n]/J, s) \\ &= (1 + s^2)^n \\ &= \frac{1}{(1 - s^2)^{n+1}} \cdot (1 - s^4)^n (1 - s^2) \\ &= F(\mathbb{C}[x_1, \dots, x_n, t], s) \prod_{i=1}^{n+1} (1 - s^{\deg \theta_i}), \end{aligned}$$

so this follows from (4.4). Note that a subsequence $\theta_1, \dots, \theta_n$ of a regular sequence $\theta_1, \dots, \theta_{n+1}$ is again a regular sequence, so from (4.4) and (2.3) we obtain

$$\begin{aligned} F(\mathbb{C}[x_1, \dots, x_n, t]/J, s) &= F(\mathbb{C}[x_1, \dots, x_n, t]/(\theta_1, \dots, \theta_n), s) \\ &= \frac{1}{(1 - s^2)^{n+1}} \prod_{i=1}^n (1 - s^{\deg \theta_i}) \end{aligned}$$

$$\begin{aligned} &= \frac{(1 + s^2)^n}{1 - s^2} \\ &= F(H_S^*(\text{Pet}), s), \end{aligned}$$

from which it follows that φ is an isomorphism.

Thus it suffices to check that $\check{\varphi}$ is an isomorphism. We already know that $\check{\varphi}$ is surjective, and from equation (2.3) we know that $F(H^*(\text{Pet}), s) = (1 + s^2)^n$. Thus, in order to show that $\check{\varphi}$ is injective it suffices to show that

$$(4.5) \quad F(\mathbb{C}[x_1, \dots, x_n]/\check{J}, s) = (1 + s^2)^n.$$

Note that by (4.4), the equality (4.5) is equivalent to saying that $\sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle x_i x_j$ ($1 \leq i \leq n$) is a regular sequence. Furthermore, by Proposition 4.2, in order to prove (4.5) it in turn suffices to show that the zero set of the collection of quadratic equations

$$(4.6) \quad \sum_{j=1}^n \langle \alpha_i, \alpha_j \rangle x_i x_j = 0, \quad 1 \leq i \leq n,$$

given by the generators of the ideal \check{J} of (4.2) is $\{0\}$, i.e., the equations (4.6) have only the trivial solution.

Suppose, in order to derive a contradiction, that (4.6) has a nontrivial solution (b_1, \dots, b_n) . In particular, setting $I = \{i \mid b_i \neq 0\}$, we have $I \neq \emptyset$ and so, since $b_i \neq 0$ for $i \in I$, we obtain from (4.6) that

$$\sum_{j \in I} \langle \alpha_i, \alpha_j \rangle b_j = 0, \quad i \in I.$$

Since $(\langle \alpha_i, \alpha_j \rangle)_{i, j \in I}$ is a $|I| \times |I|$ square matrix which is again the Cartan matrix of a semisimple Lie algebra, it must be positive definite [10, Section 2.4] and in particular nonsingular. Thus the b_i must be 0 for $i \in I$, contradicting the assumption on I . Thus (4.6) has only the trivial solution, as desired. ■

Remark Theorem 4.1 is a generalization to all Lie types of the computation given in [6]. Indeed, the generators of the ideal given in [6] are the same as those given above, up to a scalar factor of $1/2$.

Remark In fact, Theorem 4.1 holds also with \mathbb{Q} coefficients. Indeed, since both φ and $\check{\varphi}$ can be defined over \mathbb{Z} , if the maps become isomorphisms upon tensoring with \mathbb{C} then they are also isomorphisms upon tensoring with \mathbb{Q} .

Acknowledgements The first author is supported in part by an NSERC Discovery Grant (Individual), an Ontario Early Researcher Award, a Canada Research Chair (Tier 2) Award, and a Japan Society for the Promotion of Science Invitation Fellowship for Research in Japan (Fellowship ID L-13517). The first author additionally thanks the Osaka City University Advanced Mathematics Institute for its hospitality, which made this collaboration possible. The third author is partially supported by a JSPS Grant-in-Aid for Scientific Research 25400095.

References

- [1] H. Abe, M. Harada, T. Horiguchi, and M. Masuda, *The equivariant cohomology rings of regular nilpotent Hessenberg varieties in Lie type A*. In preparation; research announcement at [arxiv:1411.3065](https://arxiv.org/abs/1411.3065).
- [2] S. Billey, *Kostant polynomials and the cohomology of G/B* . *Duke Math. J.* **96**(1999), 205–224. <http://dx.doi.org/10.1215/S0012-7094-99-09606-0>
- [3] M. Brion and J. Carrell, *The equivariant cohomology ring of regular varieties*. *Michigan Math. J.* **52**(2004), 189–203. <http://dx.doi.org/10.1307/mmj/1080837743>
- [4] E. Drellich, *Monk's rule and Giambelli's formula for Peterson varieties of all Lie types*. [arxiv:1311.3014](https://arxiv.org/abs/1311.3014).
- [5] Y. Fukukawa, *The graph cohomology ring of the GKM graph of a flag manifold of type G_2* . [arxiv:1207.5229](https://arxiv.org/abs/1207.5229).
- [6] Y. Fukukawa, M. Harada, and M. Masuda, *The equivariant cohomology rings of Peterson varieties*. [arxiv:1310.8643](https://arxiv.org/abs/1310.8643). *J. Math. Soc. Japan*, to appear.
- [7] Y. Fukukawa, H. Ishida, and M. Masuda, *The cohomology ring of the GKM graph of a flag manifold of classical type*. [arxiv:1104.1832](https://arxiv.org/abs/1104.1832). *Kyoto J. Math.*, to appear. <http://dx.doi.org/10.1215/21562261-2693478>
- [8] M. Harada and J. Tymoczko, *Poset pinball, GKM-compatible subspaces, and Hessenberg varieties*. [arxiv:1007.2750](https://arxiv.org/abs/1007.2750).
- [9] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*. Graduate Texts in Math. **9**, Springer-Verlag, New York–Berlin, 1978.
- [10] J. E. Humphreys, *Reflection Groups and Coxeter Groups*. Cambridge Stud. Adv. Math. **29**, Cambridge University Press, Cambridge, 1990.
- [11] B. Kostant, *Flag manifold quantum cohomology, the Toda lattice, and the representation with highest weight ρ* . *Selecta Math. (N.S.)* **2**(1996), 43–91. <http://dx.doi.org/10.1007/BF01587939>
- [12] M. Precup, *Affine pavings of Hessenberg varieties for semisimple groups*. *Selecta Math.* **19**(2013), 903–922. <http://dx.doi.org/10.1007/s00029-012-0109-z>
- [13] K. Rietsch, *Totally positive Toeplitz matrices and quantum cohomology of partial flag varieties*. *J. Amer. Math. Soc.* **16**(2003), 363–392 (electronic). <http://dx.doi.org/10.1090/S0894-0347-02-00412-5>
- [14] R. P. Stanley, *Combinatorics and Commutative Algebra*. Second edition, Birkhäuser, Boston, 1996.

Department of Mathematics and Statistics, McMaster University, 1280 Main Street West, Hamilton, ON L8S 4K1

e-mail: megumi.harada@math.mcmaster.ca

Department of Mathematics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan

e-mail: d13sar0z06@ex.media.osaka-cu.ac.jp masuda@sci.osaka-cu.ac.jp