

# INFRARED SPECTRA OF T TAU STARS AND RELATED OBJECTS

G. I. SHANIN and V. S. SHEVCHENKO

*Astronomical Institute of the Uzbekian Academy of Science*

and

A. G. SHCHERBAKOV

*Crimean Astrophysical Observatory, Crimea, U.S.S.R.*

**Abstract.** Four T Tau stars and related objects (RY Tau, T Tau, AB Aur and V1057 Cyg) have been included in our spectroscopic programme since 1973. The present paper is concerned with the spectroscopic observations made at the Crimea with the single-stage image tube S1. Tentative atomic line identifications are given for programme stars. Ca II and O I emission line equivalent widths and profiles are presented for RY Tau, T Tau and AB Aur. The  $\lambda 10830 \text{ \AA}$  line of neutral helium has shown P Cyg-type features for T Tau and V 1057 Cyg.

## 1. RY Tau, T Tau and AB Aur

### 1.1. LINE IDENTIFICATIONS

In October 1972, October–December 1973, and June–July 1974 we obtained several spectra of programme stars in the  $0.8\text{--}1.1 \mu$  region with the 50-in. Crimean Observatory reflector and the single-stage image tube. Instrumentation has been described by Vitrichenko *et al.* (1974a, 1974b). The equipment yielded reciprocal linear dispersions of  $168$  and  $48 \text{ \AA mm}^{-1}$  with resolutions of spectrograms of about  $7$  and  $2 \text{ \AA}$ , respectively.

Tentative atomic line identifications were performed for RY Tau, T Tau and AB Aur. Primary sources of wavelengths were the *Revised Multiplet Table* (Moore, 1945) and the compilation of solar lines by Swensson *et al.* (1970). Identifications in the  $\lambda\lambda 8200\text{--}8600 \text{ \AA}$  region were obtained from  $48 \text{ \AA mm}^{-1}$  spectrograms; at  $\lambda\lambda 8000\text{--}11000 \text{ \AA}$  features were located using a  $168 \text{ \AA mm}^{-1}$  dispersion. Results are listed in Table I.

We found some absorption features at  $\lambda\lambda 10422$  and  $10440 \text{ \AA}$  and a clear emission line at  $\lambda 10568 \text{ \AA}$  for T Tau which have not been identified. Unfortunately, we have no identifications in the  $\lambda\lambda 9000\text{--}11000 \text{ \AA}$  region for many other very weak (on  $168 \text{ \AA mm}^{-1}$ ) absorption features of T Tau.

### 1.2. EMISSION LINES

The  $\lambda 10830 \text{ \AA}$  line of He I is certainly present in the spectrum of T Tau, obtained with  $168 \text{ \AA mm}^{-1}$  dispersion during the night of 1973, October 26. The profile of  $\lambda 10830$  showed P Cyg-type features with the equivalent width of the emission component of about  $3.8 \text{ \AA}$ .

In the spectrum of AB Aur obtained on October 7, 1972, no emission feature at  $\lambda 10830 \text{ \AA}$  was found with the same dispersion while broad emission in the hydrogen P 7 line ( $\lambda 10049 \text{ \AA}$ ) is clearly present. Also, the spectrum of AB Aur from 1973, Oc-

TABLE I  
Line identifications of T Tau stars<sup>a</sup>

Wavelength (Å)	Element	Mult. No	Star		
			RY Tau	T Tau	AB Aur
1	2	3	4	5	6
8374	H	11	a	a	a
8387	Fe I	60	?	a	-
8413	H	10	a	a	?
8437	H	10	a	a	?
8446	O I	4	e	e	e
8460	?	-	e	e	e
8467)	H	10	a	a	?
8468)	Fe I	60	a	a	n
8482	Fe I	999	a	a	n
8498)	Ca II	2	e	e	e
8502)	H	10	-	-	-
8514	Fe I	60	?	a	n
8542)	Ca II	2	e	e	e
8545)	H	10	-	-	-
8612	Fe I?	-	-	a	-
8662)	Ca II	2	-	e	-
8665)	H	9	-	-	-
8688	Fe I	60	-	a	-
8728	Si I?	79	-	a	-
9998)	Ti I	149	-	a	-
10001)	Si I	64	-	a	-
10003)	Ti I	193	-	a	-
10011)	Ti I	193	-	a	-
10012)	Ti I	193	-	a	-
10020)	Si I	41	-	a	-
10025)	Si I	64	-	a	-
10049	H	8	-	a	e
10422	?	-	-	a	-
10440	?	-	-	a	-
10568	?	-	-	e	-
10827	Si I?	5	-	a	-
10830	He I	1	-	e	n

<sup>a</sup> a - absorption; e - emission; n - no line; (-) - no line measurements.

tober 6 has shown strong emission lines of Ca II and O I (see Tables I and II). It is interesting that AB Aur showed no emission on the 78 Å mm<sup>-1</sup> spectrograms taken by Wallerstein (1971) during January 1970.

A microphotometer tracing of the Ca II triplet region in the T Tau spectrum is given in Figure 1. It is interesting that the Ca II central line intensities are of the same height. The complex structure of emission features in RY Tau, T Tau and AB Aur can be seen in line profiles obtained from 48 Å mm<sup>-1</sup> spectrograms (Figure 2).

Table II presents typical emission lines in these stars. The successive columns give: names of stars, dates, spectrogram numbers, dispersion, equivalent widths of the λ 8498 Å, Ca II and relative equivalent widths of lines 1<sub>λ</sub>. No corrections were made to

TABLE II  
Emission line equivalent widths of T Tau stars<sup>a</sup>

Star	Date	No sp.	D Å mm <sup>-1</sup>	W <sub>λ</sub> Ca II 8498 (Å)	I <sub>λ</sub>			O I 8446
					Ca II 8498	Ca II 8542	Ca II 8662	
1	2	3	4	5	6	7	8	9
RY Tau	Oct. 2, 73	617	48	17.4	1.00	1.17	b	0.12
T Tau	Oct. 1, 73	611	48	5.2	1.00	1.06	b	0.61
T Tau	Oct. 2, 73	616	48	4.5	1.00	1.26	b	0.62
T Tau	Oct.26, 73	646	168	4.6	1.00	0.86	0.82	0.62
T Tau	Dec.25, 73	702	168	4.6	1.00	0.51	0.70	0.61
AB Aur	Oct. 6, 73	626	48	1.8	1.00	0.92	b	0.27

<sup>a</sup> All equivalent widths have errors of the order of 20 %

<sup>b</sup> We have no observations of this line.




Fig. 1. Microphotometer tracing of the infrared Ca II triplet region of the T Tau spectrum. Original dispersion is 168 Å mm<sup>-1</sup>.




Fig. 2. Emission line profiles of Ca II and O I for RY Tau, T Tau and AB Aur obtained from  $48 \text{ \AA mm}^{-1}$  spectrograms.

the equivalent widths  $I_{\lambda}$  for the continuum energy distribution. Since we found that the hydrogen absorption lines are very weak, their contribution to the equivalent widths of the Ca II triplet lines have not been taken into account. Figure 1 and Table II indicate that observed relative intensities of lines in the Ca II triplet differ from their theoretical optically thin values 1, 9, and 5. The greatest differences take place for T Tau. A very interesting feature of the T Tau Ca II triplet is the variability of the relative line equivalent widths on a long time scale. An analysis of Table II indicates also that the triplet ratios observed in the spectrum of T Tau are not due to self-absorption.

Apart from internal variations of Ca II lines, in the same spectra of T Tau no variability of the ratio

$$W_{\lambda 8446, \text{O I}} / W_{\lambda 8498, \text{Ca II}}$$

has been observed. However, insufficient data are available at present to allow certain confirmation of the Ca II lines variations.

### 2. V 1057 Cygni

In June and July 1974, we obtained at a dispersion of  $48 \text{ \AA mm}^{-1}$  seven spectra of V 1057 Cyg in the Ca II triplet and  $\lambda 10830 \text{ \AA}$  regions. The profile of the  $\lambda 10830 \text{ \AA}$  line of neutral helium in the V 1057 Cyg spectrum is compared with the P Cyg profile (taken in July, 1974) in Figure 3. Shell features of He I in V 1057 Cyg are shown. The line has a P Cyg-type absorption component displaced about  $450 \text{ km s}^{-1}$  shortward. (According to Herbig and Harlan (1971),  $H\alpha$  has the same displacement of about  $420 \text{ km s}^{-1}$ ). In contrast to P Cyg, the He I absorption of V 1057 Cyg is very broad, which must be indicative of an enormous velocity gradient. Grasdalen (1973) reports that the equivalent width of the  $\lambda 10830 \text{ \AA}$  He I line was of the order of  $6 \text{ \AA}$  in May




Fig. 3. Comparison between observed line profiles for V 1057 Cyg and P Cyg (see text). Original dispersion is  $168 \text{ \AA mm}^{-1}$ .

1972, which is in close agreement with our estimated average value of  $7.2 \pm 0.5 \text{ \AA}$  for June 1974.

Details of the  $\lambda 10830 \text{ \AA}$  region which appear in all spectrograms of V 1057 Cyg are represented in Figure 3 by filled circles. Two features ( $\lambda\lambda 10817$  and  $10822 \text{ \AA}$ ) we interpret as emission lines of multiplet no. 118 of Cr I, which shows clear emission features without appreciable displacement at  $\lambda\lambda 10648$ ,  $10667$  and  $10672 \text{ \AA}$ . Herbig and Harlan (1971) reported emission features in the infrared Ca II triplet in 1971, but no emission has been found for 1974 and only absorption features of hydrogen and calcium lines have been detected. The  $\lambda 8446 \text{ \AA}$  line of O I is also in absorption.

Other absorption lines in the spectra of V 1057 Cyg observed in the  $\lambda\lambda 10580\text{--}10900 \text{ \AA}$  region are all lines of the carbon multiplet no. 1 plus four lines of the silicon multiplet no. 5. The possibility that the  $\lambda 10827 \text{ \AA}$  Si I absorption line contributes about 30% to the  $\lambda 10830 \text{ \AA}$  helium line is very great. The presence of C I and Si I lines and their intensities suggest that V 1057 Cyg may now have a spectral type of about F 5. It is interesting that the absorption spectrum of V 1057 Cyg in the  $\lambda\lambda 10580\text{--}10900 \text{ \AA}$  region is very similar to the spectrum of  $\gamma$  Geminorum (Cepheid, F 7–G 3 Ib) in the same region (Shanin and Shcherbakov, 1974).

### Acknowledgements

We are indebted to R. E. Gershberg and V. P. Grinin for useful discussions. We wish to thank also Tatjana Nikulina for helping with the preparation of the figures.

### References

- Grasdalen, G. L.: 1973, *Astrophys. J.* **182**, 781.  
 Herbig, G. H. and Harlan, E. A.: 1971, *IAU Info. Bull. Var. Stars* No. 543.  
 Moore, C. E.: 1945, *Revised Multiplet Table*, Princeton Univ. Obs. Contr., No. 20.  
 Shanin, G. I. and Shcherbakov, A. G.: 1974, *Bull. Krymsk. Astrofiz. Obs.*, in press.  
 Swensson, J. W., Benedict, W. S., Delbouille, L., and Roland, G.: 1970, *The Solar Spectrum from  $\lambda 7498$  to  $\lambda 12016$* , *Mem. Soc. Roy. Sci. Liège*, Special vol. No. 5.  
 Vitrichenko, E. A., Shanin, G. I., Volkov, V. U., Shevchenko, V. S., and Shcherbakov, A. G.: 1974a, *Astron. Zh. U.S.S.R.* **51**, 866.  
 Vitrichenko, E. A., Shanin, G. I., Shevchenko, V. S., and Shcherbakov, A. G.: 1974b, *New Techniques in Astronomy*, Moscow, in press.  
 Wallerstein, G.: 1971, *Publ. Astron. Soc. Pacific* **83**, 77.