

# CNS SPECTRUMS<sup>®</sup>

The International Journal of Neuropsychiatric Medicine

## Molecular Genetics: Part One

*M.T. Pato, H. Nicolini, C.N. Pato, guest editors*

### Genetics of Schizophrenia: Current Findings and Issues

*J. L. Kennedy*

### Identification of a Highly Homogenous Population for Genetic Study of Psychiatric Disorders

*K. Schindler*

### Genetic Anticipation in Portuguese Families With Bipolar Mood Disorder

*A. Macedo*

### Understanding the Genetic Basis of Obsessive-Compulsive Disorder

*H. Nicolini*

### The Genetics of Attention- Deficit/Hyperactivity Disorder

*R. Ram*

### Identification of Attention-Deficit/ Hyperactivity Disorder Pedigrees and Substance Use Disorder Pedigrees Through an ADHD Proband Sample

*A. Bauer, R. Ram*

### DNA Microarrays for Polymorphism Detection and Genotyping: Utility in the Understanding of Complex Neuropsychiatric Diseases

*P. Sklar*

**Photo Essay** This issue describes new methods and approaches which may help unravel the genetic contribution to complex neuropsychiatric disorders. **Articles Inside.**

An **MBL** Journal  
COMMUNICATIONS  
**CME 3**

# More physicians are diagnosing Alzheimer's disease.....



\*The most common adverse events leading to discontinuation in clinical trials with ARICEPT® (donepezil HCl) were nausea, diarrhea, and vomiting. Clinical studies of ARICEPT® have shown no increase, relative to placebo, in the incidence of either peptic ulcer disease or gastrointestinal bleeding. Nevertheless, cholinesterase inhibitors may be expected to increase gastric acid secretion. Therefore, patients (especially those at increased risk for developing ulcers — eg, history of ulcer disease, receiving concurrent nonsteroidal anti-inflammatory drugs) should be monitored closely for gastrointestinal bleeding. In clinical trials, syncopal episodes have been reported in association with the use of ARICEPT® (2% vs 1% for placebo).

# That's why they're prescribing ARICEPT® (donepezil HCl)

## CLINICALLY PROVEN TO ENHANCE COGNITIVE FUNCTION

With over 700,000 patient starts, ARICEPT® is the world's most-prescribed therapy for the treatment of mild to moderate Alzheimer's disease. Remember ARICEPT® for these important benefits:

- **Once-daily dosing**
- **No titration required**
- **Excellent safety profile**
- **Well-tolerated therapy\***

ONCE-A-DAY  
**ARICEPT®**  
(donepezil HCl)  
5-MG AND 10-MG TABLETS

**THERAPY TO REMEMBER™**

*Please see brief summary of prescribing information  
on the last page of this advertisement.*



**ONCE-A-DAY**  
**ARICEPT®** (donepezil HCl)  
**THERAPY TO REMEMBER™** 5-MG AND 10-MG TABLETS

**ARICEPT® (Donepezil Hydrochloride Tablets)**

**Brief Summary**—see package insert for full prescribing information. **INDICATIONS AND USAGE** ARICEPT® is indicated for the treatment of mild to moderate dementia of the Alzheimer's type. **CONTRAINDICATIONS** ARICEPT® is contraindicated in patients with known hypersensitivity to donepezil hydrochloride or to piperidine derivatives. **WARNINGS Anesthesia:** ARICEPT®, as a cholinesterase inhibitor, is likely to exaggerate succinylcholine-type muscle relaxation during anesthesia. **Cardiovascular Conditions:** Because of their pharmacological action, cholinesterase inhibitors may have vagotonic effects on heart rate (eg, bradycardia). The potential for this action may be particularly important to patients with "sick sinus syndrome" or other supraventricular cardiac conduction conditions. Syncopal episodes have been reported in association with the use of ARICEPT®. **Gastrointestinal Conditions:** Through their primary action, cholinesterase inhibitors may be expected to increase gastric acid secretion due to increased cholinergic activity. Therefore, patients should be monitored closely for symptoms of active or occult gastrointestinal bleeding, especially those at increased risk for developing ulcers, eg, those with a history of ulcer disease or those receiving concurrent nonsteroidal anti-inflammatory drugs (NSAIDs). Clinical studies of ARICEPT® have shown no increase, relative to placebo, in the incidence of either peptic ulcer disease or gastrointestinal bleeding. ARICEPT®, as a predictable consequence of its pharmacological properties, has been shown to produce diarrhea, nausea, and vomiting. These effects, when they occur, appear more frequently with the 10 mg/day dose than with the 5 mg/day dose. In most cases, these effects have been mild and transient, sometimes lasting one to three weeks, and have resolved during continued use of ARICEPT®. **Genitourinary:** Although not observed in clinical trials of ARICEPT®, cholinomimetics may cause bladder outflow obstruction. **Neurological Conditions:** Seizures: Cholinomimetics are believed to have some potential to cause generalized convulsions. However, seizure activity also may be a manifestation of Alzheimer's Disease. **Pulmonary Conditions:** Because of their cholinomimetic actions, cholinesterase inhibitors should be prescribed with care to patients with a history of asthma or obstructive pulmonary disease. **PRECAUTIONS Drug-Drug Interactions Drugs Highly Bound to Plasma Proteins:** Drug displacement studies have been performed *in vitro* between this highly bound drug (96%) and other drugs such as furosemide, digoxin, and warfarin. ARICEPT® at concentrations of 0.3-10 µg/mL did not affect the binding of furosemide (5 µg/mL), digoxin (2 ng/mL), and warfarin (3 µg/mL) to human albumin. Similarly, the binding of ARICEPT® to human albumin was not affected by furosemide, digoxin and warfarin. **Effect of ARICEPT® on the Metabolism of Other Drugs:** No *in vivo* clinical trials have investigated the effect of ARICEPT® on the clearance of drugs metabolized by CYP 3A4 (eg, cisapride, terfenadine) or by CYP 2D6 (eg, imipramine). However, *in vitro* studies show a low rate of binding to these enzymes (mean K<sub>d</sub> about 50-130 µM), that, given the therapeutic plasma concentrations of donepezil (164 nM), indicates little likelihood of interference. Whether ARICEPT® has any potential for enzyme induction is not known. Formal pharmacokinetic studies evaluated the potential of ARICEPT® for interaction with theophylline, cimetidine, warfarin and digoxin. No significant effects on the pharmacokinetics of these drugs were observed. **Effect of Other Drugs on the Metabolism of ARICEPT®:** Ketoconazole and quinidine, inhibitors of CYP450, 3A4 and 2D6, respectively, inhibit donepezil metabolism *in vitro*. Whether there is a clinical effect of these inhibitors is not known. Inducers of CYP 2D6 and CYP 3A4 (eg, phenytoin, carbamazepine, dexamethasone, rifampin, and phenobarbital) could increase the rate of elimination of ARICEPT®. Formal pharmacokinetic studies demonstrated that the metabolism of ARICEPT® is not significantly affected by concurrent administration of digoxin or cimetidine. **Use with Anticholinergics:** Because of their mechanism of action, cholinesterase inhibitors have the potential to interfere with the activity of anticholinergic medications. **Use with Cholinomimetics and Other Cholinesterase Inhibitors:** A synergistic effect may be expected when cholinesterase inhibitors are given concurrently with succinylcholine, similar neuromuscular blocking agents or cholinergic agonists such as bethanechol. **Carcinogenesis, Mutagenesis, Impairment of Fertility** Carcinogenicity studies of donepezil have not been completed. Donepezil was not mutagenic in the Ames reverse mutation assay in bacteria. In the chromosome aberration test in cultures of Chinese hamster lung (CHL) cells, some clastogenic effects were observed. Donepezil was not clastogenic in the *in vivo* mouse micronucleus test. Donepezil had no effect on fertility in rats at doses up to 10 mg/kg/day (approximately 8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis). **Pregnancy** **Pregnancy Category C.** Teratology studies conducted in pregnant rats at doses up to 16 mg/kg/day (approximately 13 times the maximum recommended human dose on a mg/m<sup>2</sup> basis) and in

**Table 2. Adverse Events Reported in Controlled Clinical Trials in at Least 2% of Patients Receiving ARICEPT® and at a Higher Frequency Than Placebo-treated Patients**

Body System/Adverse Event	Placebo (n=355)	ARICEPT® (n=747)
<b>Percent of Patients With Any Adverse Event</b>	<b>72</b>	<b>74</b>
<b>Body as a Whole</b>		
Headache	9	10
Pain, Various Locations	8	9
Accident	7	7
Fatigue	3	5
<b>Cardiovascular System</b>		
Syncope	1	2
<b>Digestive System</b>		
Nausea	6	11
Diarrhea	5	10
Vomiting	3	5
Anorexia	2	4
<b>Hemic and Lymphatic System</b>		
Echymosis	3	4
<b>Metabolic and Nutritional Systems</b>		
Weight Decrease	1	3
<b>Musculoskeletal System</b>		
Muscle Cramps	2	6
Arthritis	1	2
<b>Nervous System</b>		
Insomnia	6	9
Dizziness	6	8
Depression	<1	3
Abnormal Dreams	0	3
Somnolence	<1	2
<b>Urogenital System</b>		
Frequent Urination	1	2

age. **Other Adverse Events Observed During Clinical Trials** ARICEPT® has been administered to over 1700 individuals during clinical trials worldwide. Approximately 1200 of these patients have been treated for at least 3 months and more than 1000 patients have been treated for at least 6 months. Controlled and uncontrolled trials in the United States included approximately 900 patients. In regards to the highest dose of 10 mg/day, this population includes 650 patients treated for 3 months, 475 patients treated for 6 months and 116 patients treated for over 1 year. The range of patient exposure is from 1 to 1214 days. Treatment emergent signs and symptoms that occurred during 3 controlled clinical trials and two open-label trials in the United States were recorded as adverse events by the clinical investigators using terminology of their own choosing. To provide an overall estimate of the proportion of individuals having similar types of events, the events were grouped into a smaller number of standardized categories using a modified COSTART dictionary and event frequencies were calculated across all studies. These categories are used in the listing below. The frequencies represent the proportion of 900 patients from these trials who experienced that event while receiving ARICEPT®. All adverse events occurring at least twice are included, except for those already listed in Tables 1 or 2. COSTART terms too general to be informative, or events less likely to be drug caused, are classified by body system and listed using the following definitions: **frequent adverse events**—those occurring in at least 1/100 patients; **infrequent adverse events**—those occurring in 1/100 to 1/1000 patients. These adverse events are not necessarily related to ARICEPT® treatment and in most cases were observed at a similar frequency in placebo-treated patients in the controlled studies. No important additional adverse events were seen in studies conducted outside the United States. **Body as a Whole:** *Frequent:* influenza, chest pain, toothache; *Infrequent:* fever, edema face, periorbital edema, hernia hiatal, abscess, cellulitis, chills, generalized coldness, head fullness, listlessness. **Cardiovascular System:** *Frequent:* hypertension, vasodilation, atrial fibrillation, hot flashes, hypotension; *Infrequent:* angina pectoris, postural hypotension, myocardial infarction, AV block (first degree), congestive heart failure, arteritis, bradycardia, peripheral vascular disease, supraventricular tachycardia, deep vein thrombosis. **Digestive System:** *Frequent:* fecal incontinence, gastrointestinal bleeding, bloating, epigastric pain; *Infrequent:* eructation, gingivitis, increased appetite, flatulence, periodontal abscess, cholelithiasis, diverticulitis, drooling, dry mouth, fever sore, gastritis, irritable colon, tongue edema, epigastric distress, gastroenteritis, increased transaminases, hemorrhoids, ileus, increased thirst, jaundice, melena, polydipsia, duodenal ulcer, stomach ache. **Endocrine System:** *Infrequent:* diabetes mellitus, goiter. **Hemic and Lymphatic System: *Infrequent:* anemia, thrombocytopenia, thrombocytopenia, eosinophilia, erythrocytopenia. **Metabolic and Nutritional Disorders:** *Frequent:* dehydration; *Infrequent:* gout, hypokalemia, increased creatine kinase, hyperglycemia, weight increase, increased lactate dehydrogenase. **Musculoskeletal System:** *Frequent:* bone fracture; *Infrequent:* muscle weakness, muscle fasciculation. **Nervous System:** *Frequent:* delusions, tremor, irritability, paresthesia, aggression, vertigo, ataxia, increased libido, restlessness, abnormal crying, nervousness, aphasia; *Infrequent:* cerebrovascular accident, intracranial hemorrhage, transient ischemic attack, emotional lability, neuralgia, coldness (localized), muscle spasm, dysphoria, gait abnormality, hypertonia, hypokinesia, neurodermatitis, numbness (localized), paranoia, dysarthria, dysphasia, hostility, decreased libido, melancholia, emotional withdrawal, nystagmus, pacing. **Respiratory System:** *Frequent:* dyspnea, sore throat, bronchitis; *Infrequent:* epistaxis, postnasal drip, pneumonia, hyperventilation, pulmonary congestion, wheezing, hypoxia, pharyngitis, pleurisy, pulmonary collapse, sleep apnea, snoring. **Skin and Appendages:** *Frequent:* pruritus; diaphoresis, urticaria; *Infrequent:* dermatitis, erythema, skin discoloration, hyperkeratosis, alopecia, fungal dermatitis, herpes zoster, hirsutism, skin striae, night sweats, skin ulcer. **Special Senses:** *Frequent:* cataract, eye irritation, vision blurred; *Infrequent:* dry eyes, glaucoma, earache, tinnitus, blepharitis, decreased hearing, retinal hemorrhage, otitis externa, otitis media, bad taste, conjunctival hemorrhage, ear buzzing, motion sickness, spots before eyes. **Urogenital System:** *Frequent:* urinary incontinence, nocturia; *Infrequent:* dysuria, hematuria, urinary urgency, metrorrhagia, cystitis, enuresis, prostate hypertrophy, pyelonephritis, inability to empty bladder, breast fibroadenosis, fibrocystic breast, mastitis, pyuria, renal failure, vaginitis. **Postintroduction Reports** Voluntary reports of adverse events temporally associated with ARICEPT® that have been received since market introduction that are not listed above, and that there is inadequate data to determine the causal relationship with the drug include the following: abdominal pain, agitation, cholecystitis, confusion, convulsions, hallucinations, heart block, hemolytic anemia, hyponatremia, pancreatitis, and rash. **OVERDOSAGE** **Because strategies for the management of overdose are continually evolving, it is advisable to contact a Poison Control Center to determine the latest recommendations for the management of an overdose of any drug.** As in any case of overdose, general supportive measures should be utilized. Overdosage with cholinesterase inhibitors can result in cholinergic crisis characterized by severe nausea, vomiting, salivation, sweating, bradycardia, hypotension, respiratory depression, collapse and convulsions. Increasing muscle weakness is a possibility and may result in death if respiratory muscles are involved. Tertiary anticholinergics such as atropine may be used as an antidote for ARICEPT® overdose. Intravenous atropine sulfate titrated to effect is recommended: an initial dose of 1.0 to 2.0 mg IV with subsequent doses based upon clinical response. Atypical responses in blood pressure and heart rate have been reported with other cholinomimetics when co-administered with quaternary anticholinergics such as glycopyrrolate. It is not known whether ARICEPT® and/or its metabolites can be removed by dialysis (hemodialysis, peritoneal dialysis, or hemofiltration). Dose-related signs of toxicity in animals included reduced spontaneous movement, prone position, staggering gait, lacrimation, clonic convulsions, depressed respiration, salivation, miosis, tremors, fasciculation and lower body surface temperature. **DOSAGE AND ADMINISTRATION** The dosages of ARICEPT® shown to be effective in controlled clinical trials are 5 mg and 10 mg administered once per day. Controlled clinical trials indicate that the 10 mg dose, with a one week titration, is likely to be associated with a higher incidence of cholinergic adverse events than the 5 mg dose. Because steady state is not achieved for 15 days and because the incidence of such effects may be influenced by the rate of dose escalation, treatment with a dose of 10 mg should not be contemplated until patients have been on a daily dose of 5 mg for 4 to 6 weeks. Whether or not to employ a dose of 10 mg is a matter of prescriber and patient preference. ARICEPT® should be taken in the evening, just prior to retiring, and may be taken with or without food.**

Revised September, 1998

**Table 1. Comparison of Rates of Adverse Events in Patients Titrated to 10 mg/day Over 1 and 6 Weeks**

Adverse Event	No titration		One-week titration	Six-week titration
	Placebo (n=315)	5 mg/day (n=311)	10 mg/day (n=315)	10 mg/day (n=269)
Nausea	6%	5%	19%	6%
Diarrhea	5%	8%	15%	9%
Insomnia	6%	6%	14%	6%
Fatigue	3%	4%	8%	3%
Vomiting	3%	3%	8%	5%
Muscle Cramps	2%	6%	8%	3%
Anorexia	2%	3%	7%	3%

pregnant rabbits at doses up to 10 mg/kg/day (approximately 16 times the maximum recommended human dose on a mg/m<sup>2</sup> basis) did not disclose any evidence for a teratogenic potential of donepezil. However, in a study in which pregnant rats were given up to 10 mg/kg/day (approximately 8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis) from day 17 of gestation through day 20 postpartum, there was a slight increase in still births and a slight decrease in pup survival through day 4 postpartum at this dose; the next lower dose tested was 3 mg/kg/day. There are no adequate or well-controlled studies in pregnant women. ARICEPT® should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. **Nursing Mothers** It is not known whether donepezil is excreted in human breast milk. ARICEPT® has no indication for use in nursing mothers. **Pediatric Use** There are no adequate and well-controlled trials to document the safety and efficacy of ARICEPT® in any illness occurring in children. **ADVERSE REACTIONS Adverse Events Leading to Discontinuation** The rates of discontinuation from controlled clinical trials of ARICEPT® due to adverse events for the ARICEPT® 5 mg/day treatment groups were comparable to those of placebo-treatment groups at approximately 5%. The rate of discontinuation of patients who received 7-day escalations from 5 mg/day to 10 mg/day, was higher at 13%. The most common adverse events leading to discontinuation, defined as those occurring in at least 2% of patients and at twice the incidence seen in placebo patients were nausea (1% [5 mg] and 3% [10 mg] vs 1% [placebo]), diarrhea (<1% [5 mg] and 3% [10 mg] vs 0% [placebo]), and vomiting (<1% [5 mg] and 2% [10 mg] vs <1% [placebo]). **Most Frequent Adverse Clinical Events Seen in Association with the Use of ARICEPT®** The most common adverse events, defined as those occurring at a frequency of at least 5% in patients receiving 10 mg/day and twice the placebo rate, are largely predicted by ARICEPT®'s cholinomimetic effects. These include nausea, diarrhea, insomnia, vomiting, muscle cramp, fatigue, and anorexia. These adverse events were often of mild intensity and transient, resolving during continued ARICEPT® treatment without the need for dose modification. There is evidence to suggest that the frequency of these common adverse events may be affected by the rate of titration. An open-label study was conducted with 269 patients who received placebo in the 15- and 30-week studies. These patients were titrated to a dose of 10 mg/day over a 6-week period. The rates of common adverse events were lower than those seen in patients titrated to 10 mg/day over one week in the controlled clinical trials and were comparable to those seen in patients on 5 mg/day. See Table 1 for a comparison of the most common adverse events following one week and six week titration regimens. **Adverse Events Reported in Controlled Trials** The events cited reflect experience gained under closely monitored conditions of clinical trials in a highly selected patient population. In actual clinical practice or in other clinical trials, these frequency estimates may not apply, as the conditions of use, reporting behavior, and the kinds of patients treated may differ. Table 2 lists treatment emergent signs and symptoms that were reported in at least 2% of patients in placebo-controlled trials who received ARICEPT® and for which the rate of occurrence was greater for ARICEPT® assigned than placebo assigned patients. In general, adverse events occurred more frequently in female patients and with advancing



**Eisai Inc.**  
 Teaneck, NJ 07666

MADE IN USA



**Pfizer U.S. Pharmaceuticals**  
 New York, NY 10017

# CNS SPECTRUMS

The International Journal of Neuropsychiatric Medicine

## EDITOR

**Eric Hollander, MD**  
Mount Sinai School of Medicine  
New York, NY

## INTERNATIONAL EDITOR

**Joseph Zohar, MD**  
Chaim Sheba Medical Center  
Tel Aviv, Israel

## ASSOCIATE INTERNATIONAL EDITOR

**Donatella Marazziti, MD**  
University of Pisa  
Pisa, Italy

## EDITORIAL DIRECTOR

**James La Rossa Jr.**

## BOARD OF ADVISORS

**Margaret Altemus, MD**  
Cornell University Medical Center  
New York, NY

**Mitchell F. Brin, MD**  
Mount Sinai School of Medicine  
New York, NY

**John Caronna, MD**  
New York Hospital-Cornell  
Medical Center, New York, NY

**Dennis S. Charney, MD**  
Yale University  
New Haven, CT

**Emil F. Coccaro, MD**  
MCP at EPPH  
Philadelphia, PA

**Jeffrey L. Cummings, MD**  
University of California  
Los Angeles, CA

**Dwight L. Evans, MD**  
University of Pennsylvania  
Philadelphia, PA

**Mark George, MD**  
Medical University of South Carolina  
Charleston, SC

**Jack Gorman, MD**  
College of Physicians and  
Surgeons, Columbia University  
New York, NY

**Thomas R. Insel, MD**  
Yerkes Primate Labs  
Emory University School of Medicine  
Atlanta, GA

**Michael A. Jenike, MD**  
Massachusetts General Hospital  
Charlestown, MA

**Lorin M. Koran, MD**  
Stanford University Medical School  
Stanford, CA

**James Leckman, MD**  
Yale University  
New Haven, CT

**Herbert Y. Meltzer, MD**  
Vanderbilt University Medical Center  
Nashville, TN

**Stuart A. Montgomery, MD**  
St. Mary's Hospital Medical School  
London, United Kingdom

**Dennis L. Murphy, MD**  
National Institute of Mental Health  
Bethesda, MD

**Charles B. Nemeroff, MD, PhD**  
Emory University School of Medicine  
Atlanta, GA

**Humberto Nicolini, MD, PhD**  
Instituto Mexicano de Psiquiatria  
Mexico

**Katharine Phillips, MD**  
Brown University  
Providence, RI

**Harold A. Pincus, MD**  
American Psychiatric Association  
Washington, DC

**Scott L. Rauch, MD**  
Massachusetts General Hospital  
Charlestown, MA

**Stanley I. Rapoport, MD**  
National Institute of Mental Health  
Bethesda, MD

**Alan Schatzberg, MD**  
Stanford University Medical School  
Stanford, CA

**Dan J. Stein, MB**  
University of Stellenbosch  
Tygerberg, South Africa

**Norman Sussman, MD**  
New York University Medical School  
New York, NY

**Neal R. Swerdlow, MD, PhD**  
University of California, San Diego  
La Jolla, CA

**Michael R. Trimble, MD**  
National Hospital for Neurology  
and Neurosurgery  
London, United Kingdom

**H. M. van Praag, MD**  
University of Maastricht  
Maastricht, The Netherlands

**Herman G.M. Westenberg, MD**  
University Hospital Utrecht  
Utrecht, The Netherlands

**Richard Wyatt, MD**  
National Institute of Mental Health  
Bethesda, MD

**Stuart Yudofsky, MD**  
Baylor College of Medicine  
Houston, TX

## MBL COMMUNICATIONS

### CEO & PUBLISHER

**James La Rossa Jr.**

### PRESIDENT & ASSOCIATE PUBLISHER

**Darren L. Brodeur**

### MANAGING EDITOR

**Claire R. Roberts**

### ASSOCIATE EDITORIAL DIRECTOR

**Genevieve Romano**

### SECTION EDITOR

**Steven Ovadia**

### SENIOR EDITOR

**Jenny R. Green**

### SPECIAL PROJECTS EDITOR

**Marla K. Lehner**

### ASSOCIATE EDITOR

**Lisa Nicpon**

### PUBLISHING ASSOCIATE

**Imre Balanli**

### ADMINISTRATIVE

#### ASSISTANT

**Leelawatee Ramadhin**

#### ART DIRECTOR

**Anthony J. Korsak**

#### COPY EDITORS

**Lauren A. Cerruto**  
**Michelle Cervone, MD**  
**Clinton Corbett**  
**John Martino**  
**Laura Ninger**  
**Meg Phelan**

### CONTROLLER

**Deborah Policarpio Gomez**

### CORPORATION COUNSEL

**Kevin F. Saer, Esq.**  
Davis, Wright & Tremaine

### OF COUNSEL

**Susan G. La Rossa, Esq.**  
Putney, Twombly, Hall &  
Hirson